

**The stress energy tensor of a locally supersymmetric quantum field on a
curved spacetime**

Dissertation
zur Erlangung des Doktorgrades
des Fachbereichs Physik
der Universität Hamburg

vorgelegt von
Martin Köhler
aus Berlin

Hamburg
1995

DESY 95-080

hep-th/9505014

Gutachter der Dissertation:	Prof. Dr. K. Fredenhagen Prof. Dr. D. Buchholz
Gutachter der Disputation:	Prof. Dr. K. Fredenhagen Prof. Dr. H. Nicolai
Datum der Disputation:	4. April 1995
Sprecher des Fachbereichs Physik und Vorsitzender des Promotionsausschusses:	Prof. Dr. B. Kramer

ABSTRACT. For an analogon of the free Wess-Zumino model on Ricci flat spacetimes, the relation between a conserved ‘supercurrent’ and the point-separated improved energy momentum tensor is investigated and a similar relation as on Minkowski space is established. The expectation value of the latter in any globally Hadamard product state is found to be a priori finite in the coincidence limit if the theory is massive. On arbitrary globally hyperbolic spacetimes the ‘supercurrent’ is shown to be a well defined operator valued distribution on the GNS Hilbertspace of any globally Hadamard product state. Viewed as a new field, all n -point distributions exist, giving a new example for a Wightman field on that manifold. Moreover, it is shown that this field satisfies a new wave front set spectrum condition in a non trivial way.

ZUSAMMENFASSUNG. In dieser Arbeit wird ein Analogon zum freien supersymmetrischen Wess-Zumino Modell auf dem Hintergrund einer Ricci-flachen, global hyperbolischen Raumzeit betrachtet. Die zu diesem Modell gehörenden ‘Superströme’ und deren Beziehungen zum erweiterten Energie Impuls Tensor werden untersucht. Es zeigt sich, daß sich in diesem Modell, wie im Minkowskiraum, der durch ‘Punktetrennung’ regularisierte Energie Impuls Tensor mit Hilfe der ‘Superströme’ erzeugen läßt. Ferner ergibt sich unter der Voraussetzung einer massiven Theorie, daß der Erwartungswert von letzterem auch bei Aufhebung der Regularisierung endlich bleibt. Ein weiteres Resultat dieser Arbeit ist, daß die ‘Superströme’ auf beliebigen global hyperbolischen Raumzeiten operatorwertige Distributionen auf dem GNS-Hilbertraum eines jeden globalen Hadamardproduktzustandes definieren. Diese Distributionen erfüllen die üblichen Wightman Axiome und können deshalb als Beispiele für neue Wightman Felder auf Mannigfaltigkeiten dienen. Insbesondere existieren alle n -Punkt Distributionen und diese erfüllen in nicht trivialer Weise eine neue Wellenfronten-Spektrumsbedingung.

Contents

Introduction	7
Chapter 1. The Wess Zumino model on Minkowski spacetime	11
1. Classical spinors in Minkowski spacetime	11
2. The internal symmetries of the classical free Wess-Zumino Model	15
3. Normal Ordering	18
Chapter 2. Spinors on Lorentz manifolds	21
1. Bundle-formalism for spinors	21
2. The spinor connection	23
3. Čech-cohomology groups	25
4. Stiefel-Whitney classes	25
5. An existence theorem	27
Chapter 3. An analogon of the free Wess-Zumino model	29
1. A locally conserved ‘supercurrent’	31
2. The improved ‘supercurrent’	32
3. Quantization: The algebraic approach to quantum field theory	33
4. Hadamard states	36
5. Wave front sets	37
6. The wave front set spectrum condition	45
Chapter 4. The energy momentum tensor	49
1. New Wightman fields: The ‘supercurrents’ after quantization	49
2. The ‘supercharge’	51
3. The energy momentum tensor in terms of ‘supercurrents’ and ‘supercharges’	72
4. The ‘vacuum’ expectation value	74
5. Two examples: The Schwarzschild and Kerr solution	78
Discussion and outlook	81
Appendix A. Algebraic topology	85
1. Basic Terminology	85
2. Sheaf theory	86
3. Sheaf cohomology	87
Appendix B. The Hadamard recursion relations	89
1. The coincidence limit of the smooth DeWitt-Schwinger term	90
Appendix C. Useful formulas	91
1. γ -matrix identities	91
2. Rearrangement of Majorana spinors	91
Appendix D. Listing of the Maple V programs	93
1. The Schwarzschild spacetime	93
2. The Kerr spacetime	96
Bibliography	101

Introduction

In the last decade there was remarkable progress in the construction of unified theories for the fundamental forces of nature. The unification of the electro-magnetic with the weak interaction summit into the Salam-Weinberg theory, while attempts to incorporate strong interaction, described by quantum chromodynamics, seem to achieve success in the so-called ‘supersymmetric grand-unified theories’ (SUSY-GUTs). Ordinary GUTs, despite of their qualitative and quantitative success, for instance equality of different coupling constants at high energies, disparity of quark-lepton masses at low energy, determination of the electro-weak mixing-angle, etc., unfortunately contain baryon and lepton number violating interactions, which contradict the presently observed stability of the proton (lifetime $> 10^{32}$ years). On the other hand the addition of supersymmetry leaves the standard success of the ordinary GUTs more or less intact and predicts $10^{32\pm 2}$ years for the lifetime of the proton, which is compatible with the lower experimental bound mentioned above. However, an incorporation of gravity is still one of the greatest challenges of modern physics. In the last years there were an increasing number of approaches to this problem which look promising and the question arose whether one can say anything about the influence of gravity on quantum phenomena. In some sense this situation is comparable to the one encountered just before the discovery of quantum electrodynamics. In those days of quantum mechanics, one treated the electro-magnetic field in the calculation as an external classical potential, while the matter was already described quantum mechanically. The resulting theory, which one would call a semiclassical approximation nowadays, already gave some results which were later shown to be in complete agreement with the fully quantized theory, i.e., quantum electrodynamics. On the other hand those calculations showed also that a semiclassical theory leads to inconsistencies if it is stressed too far. One may therefore hope that a similar semiclassical description is also valid for gravity. Quantum field theory on curved spacetimes (QFT on CST) provides such a framework: The spacetime is described classically by a manifold together with a metric field, while all the matter is described as quantized fields, which propagate on that spacetime. The attempt to translate the considerations on the domain of validity from quantum electrodynamics (QED) to QFT on CST indicates that the latter is expected to be huge, namely from the macroscopic scale down to the order of the Planck length ($\approx 10^{-33}$ cm) or Planck time ($\approx 10^{-44}$ s) respectively. This huge domain should leave enough room for a fruitful semiclassical theory. However, there is a subtle problem with such naive comparison of QED and QFT on CST: Due to the equivalence principle, gravitational energy couples equally strong to geometry, as does all other matter. Hence gravity will enter in a non trivial way at *all* scales. Fortunately one may still succeed in a semiclassical description of gravity. To see the argument, recall that in classical general relativity gravitational waves are described as small perturbations of the background metric. These perturbations can be treated as a null fluid and their contribution to the l.h.s. (i.e., to the geometrical side) of the Einstein equation can be separated off from the (geometrical) rest in such a way that they may be treated as part of the source, i.e., as part of T^{ab} . It seems therefore reasonable to suppose that the ‘graviton’ field, representing linearized perturbations of the background spacetime, can be included along with all other fields as part of the matter rather than geometry. This approximation should work so long as one stays clearly away from the Planck scale. Due to the weakness of gravitational effects, possible direct applications of this framework seemed to lie in the early cosmology or in the description of microscopic black holes. This situation changed when Hawking during his studies of quantum black holes in 1975 [Haw75] discovered the by now famous (macroscopic) black hole radiation, opening a broad new area of physics. Although this radiation is still too weak to be detected experimentally (e.g., the temperature of a black hole of a mass ten times that of our sun is approximately 10^{-8} K, which is eight orders of magnitude below the the temperature of the cosmic background radiation), his result is remarkable for at least two reasons. First, it seems to be universal in the sense that it can be established using very general assumptions on the quantum fields involved [FH90] and it has been reproduced in various different ways. Second, it establishes and fixes the close relation between black holes and thermodynamics, which had been suggested already before the application of quantum field theory to black hole physics. Hawking’s radiation corresponds to a continuous energy-flow from the black hole to infinity and, at least due to energy conservation, should be accompanied by a mass-loss of the former, i.e., the black hole should evaporate. For a precise mathematical formulation of this back-reaction problem, a local stress energy tensor for the quantum field is needed. This object, denoted by T^{ab} , is important for another reason, too: Since the work of Unruh [Unr76] it is known that the traditional particle concept of Minkowski quantum field theory is not suitable for quantum fields on curved spacetimes. Instead the physical situation in the latter case should be described in a more operational way, i.e., in terms of detectors and counting rates. Such a description is provided by a local stress tensor. Following Hawking’s paper, in

the mid seventies, there were undertaken enormous efforts on the study of T^{ab} and its expectation value in some states ω ($\langle T^{ab} \rangle_\omega$). Unfortunately the classical expression for the stress energy tensor in the cases of the standard fields (e.g., scalar, Dirac, photon, etc.) yields divergent results after quantization, because products of field operators at the same point occur. One therefore has to subtract out the divergences in some way, i.e., to regularize and renormalize. The various renormalization techniques which were developed in that time all gave a locally conserved expectation value for the stress tensor, which therefore is a good candidate for the r.h.s. of the semiclassical Einstein equation, even if particle creation due to gravitational effects occur. Wald was even able to formulate a set of axioms [Wal77, Wal78] which almost uniquely fixes these renormalized expectation values for every state under consideration. Despite of their success these methods are still burdened with some fundamental difficulties. The first one concerns renormalization. In Minkowski space one agrees on the fact that only energy *differences* are measurable. Hence any energy renormalization should not change the physical content of the theory. On the other hand in classical general relativity the *absolute* value of the energy curves spacetime. This means that energy renormalization could change the physical content of the theory in that case. To avoid renormalization, one could for example use a mathematical framework which allows products of field operators at the same spacetime point to be well defined. The new generalized functions of Colombeau [Col84], which can be extended to manifolds [Köh91], could be used to implement such a setting. However, at the level of interpretation such a scheme is believed to be in fact equivalent to the renormalization prescriptions mentioned above. It follows that a model which has a finite stress energy tensor a priori would be highly desirable. In fact such a model is already discussed by Unruh [Unr76] in 1+1 dimensions, but apparently his ansatz was never extended to the four dimensional case. Meanwhile one did discover on Minkowski space a whole class of models with such a remarkable finiteness property. These models possess, besides the usual Poincaré symmetry, an internal symmetry, the generators of which anti-commute with those of the Poincaré group. The resulting algebra for all generators turned out to be a graded Lie-algebra, which is now called a supersymmetric extension of the Poincaré algebra. Consequently, these models were summarized under the name supersymmetric models. One of their fundamental properties is the fact that they possess equally many bosonic and fermionic degrees of freedom and all particles of one supermultiplett must have equal masses¹. Moreover, the stress energy tensor of these models can be expressed locally in terms of spinor currents [FZ75], a representation which manifestly shows a priori finiteness of the energy. Wess and Zumino [WZ74] gave the first field theoretical example of a global supersymmetric field theory in 1974 and the subject evolved rapidly from thereon. On the other hand it is evident from experiments that (unbroken) supersymmetry is *not* realized in nature. To solve this contradiction, the mechanism of spontaneous symmetry breaking was introduced into the framework, but it must be said that even today there are only hints on supersymmetry, but no experimental verification. It is well known that supersymmetry –together with general covariance– leads unavoidably to supergravity theories [Nie81], hence leaves the frame of QFT on CST. The only known non trivial example of a spacetime which admits a supersymmetric extension of its isometry group is the anti-DeSitter solution of Einstein’s equation [BF82]. Unfortunately this manifold is not globally hyperbolic and this leads to problems during quantization, namely boundary conditions at infinity are needed. On the other hand if one is interested in a cancellation of the divergences of the stress energy tensor only, a model with the same *particle contents* as a ‘real’ supersymmetric model should suffice. Coming back to the renormalization of the stress tensor on a curved spacetime, a second difficulty is a direct implication of one of the major, still unsolved problems of QFT in CST. Namely the characterization of physical admissible states. (Recall that the expectation value of T^{ab} must be calculated in some state ω .) The problem occurs, since field theories have infinite many degrees of freedom and hence there exist inequivalent representations of the canonical (anti) commutation relations. It follows that the usual Hilbertspace approach, where all physical admissible states are identified with positive trace class operators, is insufficient in general. Instead one should use the algebraic approach to quantum field theory, to which Haag’s book [Haa92] provides a good introduction. The basic objects in this approach are the ‘algebra of local observables’ which, roughly speaking, describes elementary events, together with states, which describe the preparation of the physical system or equivalently the probability for subsequent events. It is worth noting that every choice of a state on the algebra of local observables uniquely fixes, via the GNS reconstruction theorem, a Hilbertspace and a ‘vacuum’ vector, on which the observables act as bounded linear operators. For linear scalar and spinor fields on globally hyperbolic spacetimes, the algebra of local observables, satisfying the axioms of algebraic quantum field theory, was successfully constructed by Dimock in the 1980’s [Dim80, Dim82]. In the exceptional case of Minkowski space, there exists an – usually unique– Poincaré invariant state, satisfying the spectrum condition, i.e., the vacuum, which is used to construct all other physical relevant states for particle physics. Unfortunately both, Poincaré invariance and spectrum condition rely on the globally flat structure of Minkowski space; thus they can not be extended naturally to generic manifolds. On the other hand for free fields the class of the so-called Hadamard states seems to be a suitable replacement for the vacuum. Those states are characterized locally by the short distance behavior of their two-point distributions [FSW78]. It was shown in [Ver94] that they satisfy the condition of local definiteness due to Haag, Narnhofer and Stein [HNS84] and that their local von Neumann algebras are factors of type III₁ (see also [Wol92]). Finally Hadamard states are precisely those states for which the expectation value of the stress energy tensor can successfully be renormalized. We should also mention

¹We do assume the supersymmetry to be unbroken.

another class of states, which is believed to be equivalent in the domain of application, namely the adiabatic vacuum states, which were closely investigated by Lüders and Roberts [LR90]. This work is organized as follows: Following this Introduction in chapter 1 the ideas to represent the stress energy tensor in terms of supercurrents is motivated and some basic properties of the free Wess-Zumino model, which is the simplest supersymmetric model on Minkowski space, are reviewed. The classical settings for this model are described and a simple argument is presented, which shows that its energy operator is a priori normal ordered after quantization. In the next chapter the basic notation for the definition of spinors on Lorentz manifolds is established. We expose the relation between topological properties of the spacetime and the existence of classical spinor fields. Chapter 3 describes our analogon of the free Wess-Zumino model. We begin with the classical settings and introduce spinor currents which shall later generate the stress energy tensor. The assumption that these currents should be locally conserved restricts the underlying spacetime to have vanishing scalar curvature. For the convenience of the reader this classical setting is followed by a brief introduction to the algebraic approach to quantum field theory. In a next step the set of admissible states is fixed to consist of Hadamard states only. We give a precise definition of these states due to Kay and Wald [KW91] and extend their definition for scalar fields to fermions. Unfortunately this condition is quite complicated and difficult to check in actual calculations. Moreover, various questions concerning multiplication of Hadamard distributions and their restriction to hypersurfaces are almost impossible to deal with in this setting. Fortunately Hörmander and Duistermaat developed the mathematical theory of Fourier integral operators [Hör71, DH72], in which the concept of wave front sets occur. In terms of wave front sets the questions mentioned above can be answered quite easily. Radzikowski was the first who gave a characterization of Hadamard states by their wave front sets [Rad92]. However, his proof as it stands is only valid in Minkowski space. The gap in his argument is filled in section 5.1 of chapter 3, where the wave front set of any Hadamard state on an arbitrary globally hyperbolic Lorentz manifold is calculated. We close this chapter by proposing a general wave front set spectrum condition, which might serve in the future as a replacement for the usual spectrum condition on manifolds. In chapter 4 the stress energy tensor of our model is calculated. The quantized versions of the spinor currents defined in chapter 3 are shown to yield to new Wightman fields on the spacetime, whose n -point distribution satisfy our new wave front set spectrum condition. Mimicking the situation on Minkowski space, we *define* a candidate for the energy momentum tensor restricted to a spacelike hypersurface $\mathcal{C}$ in terms of the spinor current. The result, which is a distribution *on* $\mathcal{C}$ can be expressed in terms of the basic fields and equals the point-separated ‘true’ stress energy tensor if and only if the spacetime is Ricci flat, i.e., if it is a vacuum solution of the Einstein equation. Using the results of Christensen [Chr76, Chr78], it is shown next that the expectation value of our point-separated stress energy tensor, with respect to *every* Hadamard product state has finite coincidence limit after elementary averaging, i.e., it defines a (locally conserved) tensor field on the spacetime. As a consequence of this a priori normal ordering feature, the stress energy tensor turns out to be a second new Wightman field on the spacetime. We close the work with two examples: On the Schwarzschild and Kerr solutions of the Einstein equation the finite terms of our energy momentum tensor in the DeWitt-Schwinger pseudo-state are calculated.

The Wess Zumino model on Minkowski spacetime

A field theoretical realization of the super-Poincaré algebra yielding its lowest dimensional representation was discovered and studied by J. Wess and B. Zumino in 1974 [WZ74]. This model describes two spin-0 fields (A, B), one of which is scalar (A), the other (B) is pseudo-scalar and a third (Ψ) which is a Majorana spinor field. In this chapter we describe some features of this model, which motivate our subsequent studies on curved spacetimes. To establish the mathematical settings, we start with an introduction to classical spinors on Minkowski space. The classical Lagrangian of the Wess-Zumino model is given next and its internal symmetries are investigated. We close the chapter by showing that the energy operator in the quantized version of this supersymmetric theory is a priori normal ordered and obtain a formula for the energy-momentum tensor in terms of suitable spinor currents.

1. Classical spinors in Minkowski spacetime

Spinors arise most naturally in quantum field theory on Minkowski spacetime. It is even possible to show that all classical (e.g. tensorial) objects can be constructed using spinors and some authors even believe that spinors are the most fundamental objects in physics (See [PR84]). This statement clearly shows the importance of the concept of spinors, even if one does not follow Penrose's and Rindler's extreme point of view. This section starts with a motivation, which is followed by the definition of classical spinor fields on Minkowski space. The notion of Dirac- and charge conjugation is introduced next and the definition of Majorana spinors closes the section. Consider a spacetime (M, g_{ab}) , which we assume to be a four dimensional C^∞ -manifold M together with a Lorentz metric g_{ab} with signature $(+, -, -, -)$. Let $\mathcal{S}$ denote the set of all physical states on this spacetime. At this stage, it is not necessary to specify $\mathcal{S}$ more precisely. However, we assume the elements of $\mathcal{S}$ to be characterized *locally*; the usual tensor fields on M are an example. The principle of 'special covariance', i.e., every physically admissible result of a measurement which is performed by an observer $\mathcal{O}$ must also be a physically admissible result of a measurement performed by a second observer $\tilde{\mathcal{O}}$ who is related to $\mathcal{O}$ by an isometry, induces for every isometry ϱ of the spacetime a map on the physical states. This map $\tilde{\varrho} : \mathcal{S} \rightarrow \mathcal{S}$ is fixed by the requirement that every state $\tilde{\varrho}(\omega)$ ($\omega \in \mathcal{S}$) gives the same results of measurement for the observer $\tilde{\mathcal{O}}$ as ω for $\mathcal{O}$. The group of isometries for Minkowski space $(\mathbb{R}^4, \eta_{ab})$ is the well known extended Poincaré group $(\mathcal{P})$. The experiments of Wu ([W⁺57]) (violation of parity) and Fitch ([Fit81]) (violation of time-reversal if the CPT-Theorem is valid) however indicate that the physical laws on Minkowski space are specially covariant under the restricted Poincaré group $(\mathcal{P}_0)$ only. We therefore restrict ourself to this subgroup. It was shown in 1939 by Wigner [Wig39] that the assumption that $\tilde{\varrho}$ does not change the transition amplitudes for pure states in quantum mechanics¹ implies that $\tilde{\varrho}$ is a projective representation of the proper Poincaré group. Moreover Wigner showed that adding the requirement that the representation should depend continuously on the group elements fixes this representation up to sign. As a next step it was proved by Bargman [Bar54] in 1954 that those representations are in one-to-one correspondence to the true representations of the covering group $\hat{\mathcal{P}}_0$ of the proper Poincaré group. Every unitary representation of $\hat{\mathcal{P}}_0$ can be decomposed into a direct sum (or a direct integral) of irreducible representations. To study the former it is therefore sufficient to classify all irreducible representations. The latter can be classified partially by the mass (m^2) and the sign of the energy (ϵ).² If $m^2 > 0$ and $\epsilon = +1$ one further invariant s , called spin, is needed to fix the representation up to unitary equivalence. It can be shown (see e.g. [B⁺90]) that all irreducible physical representations of $\hat{\mathcal{P}}_0$ can be constructed in the space of spinor wave functions (see below). We remark that this space carries a representation of $\hat{\mathcal{P}}_0$ by construction. Moreover there exists a positive definite $\hat{\mathcal{P}}_0$ invariant form such that the equivalence classes of the spinor wave functions with respect to this form span a pre-Hilbertspace (see [B⁺90, p. 285] for the details). Completing this space yields a Hilbertspace which carries the unitary representation of $\hat{\mathcal{P}}_0$ we are looking for.

REMARK . Note that the Hilbertspace mentioned above is a one-particle Hilbertspace. The quantum statistics will not be implemented until the introduction of a many-particle Hilbertspace (Fockspace).

¹It is assumed in the proof that pure states are represented as one-dimensional projectors on a separable Hilbertspace.

²Let P^μ and $M^{\lambda\nu}$ denote the generator of the representation. The three invariants mass (m^2), sign of the energy (ϵ) and spin (s) are given by: $m^2 = P_\mu P^\mu = P^2$; $\epsilon = \text{sign}(P^0)$; $-m^2(s^2 + s) = W^2$, where $W = \epsilon_{\mu\nu\lambda\rho} P^\nu M^{\lambda\rho}$ is the Pauli-Lubanski vector.

1.1. $SL(2, \mathbb{C})$ -spinors. In this subsection the space of spinor tensor fields is constructed using the two component formulation of Penrose & Rindler [PR84]. This formulation is more in the spirit of the introduction than the usual four component Dirac approach which will be used later on exclusively, since two component $SL(2, \mathbb{C})$ -spinors transform according to an *irreducible* representation of the homogeneous part of $\hat{\mathcal{P}}_0$ and there is a close relation between spinor tensors and spinor wave functions. We note first that $\hat{\mathcal{P}}_0$ is isomorphic to $ISL(2, \mathbb{C})$, the semidirect product of the group of translations with the group of linear transformations with unit determinant acting on a two dimensional complex Hilbertspace. Let (W, ϵ_{xy}) be such a Hilbertspace together with a volume form ϵ_{xy} . Then the homogeneous part of $ISL(2, \mathbb{C})$ in its defining representation leaves the form ϵ_{xy} invariant. The pair (W, ϵ_{xy}) is called a *spinor space*, its elements are called *spinors*. Let (W^*, ϵ^{xy}) be the dual space of W , $(\bar{W}, \bar{\epsilon}_{x'y'})$ the space of all anti-linear maps from W^* into $\mathbb{C}$. The map

$$\begin{aligned} \bar{\cdot} : W &\rightarrow \bar{W} \\ \xi^x &\mapsto \bar{\xi}^{x'} : (\nu_x \mapsto \overline{\xi^x \nu_x}) \end{aligned}$$

is called *complex conjugation* and induces a mapping between ϵ_{xy} and $\bar{\epsilon}_{x'y'}$. To clarify the notation we denote all elements of $\bar{W}$ with upper primed indices. The dual space of $\bar{W}$ is denoted by $(\bar{W}^*, \bar{\epsilon}^{x'y'})$. The elements of W^* ($\bar{W}$, $\bar{W}^*$) are called dual (adjoint, adjoint dual) spinors. The space of spinor tensors $\mathbf{T}$ is the tensor product of these spaces. Note that the relative order of primed and unprimed indices is irrelevant, i.e., $v^{xx'y}$ and $v^{xyx'}$ are equivalent, since $W \otimes \bar{W} \otimes W$ and $W \otimes W \otimes \bar{W}$ are naturally isomorphic. However $v^{xyx'}$ and $v^{yx'x}$ are different in general. Smooth mappings from the Minkowski space into $\mathbf{T}$ are called *spinor tensor fields*. An action of $ISL(2, \mathbb{C})$ on these mappings can be defined by brute force as follows: To any $g \in ISL(2, \mathbb{C})$ we associate the transformation

$$\xi^x(q) \mapsto L^x_y \xi^y[\mathcal{P}^{-1}(q)], \quad (1)$$

where $L^x_y \in SL(2, \mathbb{C})$ is the homogeneous part of g , $\mathcal{P}$ denotes the associated Poincaré element and q is a point in Minkowski space.

REMARK .

- This transformation is a true representation of $\hat{\mathcal{P}}_0$, while for the Poincaré group itself it is a representation up to sign only.
- The representation given by Eqn. (1) is *not* irreducible, since spinor tensors are defined on the configuration space. However it has the advantage that one can identify certain spinor tensors with ordinary tensors. To obtain all *irreducible* representations mentioned above, one has to introduce spinor wave functions, which are defined on momentum space together with an appropriate transformation law: For any non negative integer or half integer j , consider the symmetric tensor product of $2j$ copies of W together with the self representation of $SL(2, \mathbb{C})$ and its complex conjugate. Mappings from the momentum space, which is isomorphic to $\mathbb{R}^4$, into one of these tensor products are spinor wave functions. A transformation law, which is identical to Eqn. (1), up to a factor containing the inhomogeneous part of the corresponding element $g \in ISL(2, \mathbb{C})$, finishes the construction. For more details, e.g., the explicit form of the transformation, for the proof of the claimed irreducibility as well as for the proof of the statement that all irreducible representation of $\hat{\mathcal{P}}_0$ can be obtained in this way, the reader is referred to the book of Bogolubov loc. cit. and the references therein.

We establish now the relation between so called real spinor tensor fields and vector fields on Minkowski space. Let (W, ϵ_{xy}) be a spinor space and let us denote the tensor product of W with $\bar{W}$ by Y . Complex conjugation maps Y into itself. Those elements in Y which are invariant under complex conjugation are called *real spinor tensors* and an analogous definition is valid for spinor tensor fields.

REMARK . The term ‘real’ does not make sense for spinors, i.e., for elements of W . To see this consider the case $W = \mathbb{C}^2$, $z \in \mathbb{C}^2$. The definition $\text{Re}(z) := 1/2(z + \bar{z})$ obviously is *not* invariant under the action of $SL(2, \mathbb{C})$.

The real elements of Y span a (real) four dimensional subspace of Y , which we denote by V . Let σ^x, ι^x be a basis in W with the property $\sigma^x \iota^y \epsilon_{xy} = 1$. Then

$$t^{xx'} = \frac{1}{\sqrt{2}}(\sigma^x \bar{\sigma}^{x'} + \iota^x \bar{\iota}^{x'}) \quad (2)$$

$$x^{xx'} = \frac{1}{\sqrt{2}}(\sigma^x \bar{\iota}^{x'} + \iota^x \bar{\sigma}^{x'}) \quad (3)$$

$$y^{xx'} = \frac{i}{\sqrt{2}}(\sigma^x \bar{\iota}^{x'} - \iota^x \bar{\sigma}^{x'}) \quad (4)$$

$$z^{xx'} = \frac{1}{\sqrt{2}}(\sigma^x \bar{\sigma}^{x'} - \iota^x \bar{\iota}^{x'}) \quad (5)$$

are a basis in V . Moreover $g_{xx'yy'} := \epsilon_{xy}\bar{\epsilon}_{x'y'}$ defines a Lorentz metric on V (For the proof calculate explicitly that (2)–(5) are an orthogonal basis for $g_{xx'yy'}$). Finally consider a tetrad field (t^a, x^a, y^a, z^a) on Minkowski space. The following hybrid vector spinor field

$$\sigma_{xx'}^a := t^a t_{xx'} - x^a x_{xx'} - y^a y_{xx'} - z^a z_{xx'}$$

defines for every point $q \in \mathbb{R}^4$ a vector space isomorphism between the tangential space $T_q \mathbb{R}^4$ at this point and V . Note that we lowered the indices with the metric $g_{xx'yy'}$. This isomorphism respects the metric structure ($\eta^{ab} \equiv \sigma^a_{xx'} \sigma^b_{yy'} g^{xx'yy'}$) and the action of the Poincaré group (See [Wal84]). Thus having chosen the map $\sigma^a_{xx'}$ we can identify real spinor tensor fields with vector fields.

1.2. Dirac spinors. While $SL(2, \mathbb{C})$ -spinors simplify the calculations in most cases since no Dirac γ -matrices occur, the study of uncharged fermions is best done using the Dirac description. Consider the direct sum of two spinor spaces (W, ϵ_{xy}) and $(\bar{W}, \bar{\epsilon}_{x'y'})$. Denoting this space by E , E carries a (reducible) representation of $SL(2, \mathbb{C})$ inherited from W and $\bar{W}$ respectively:

$$\pi : L \mapsto \begin{pmatrix} L & 0 \\ 0 & \bar{L} \end{pmatrix} \equiv S \quad \forall L \in SL(2, \mathbb{C}) \quad (6)$$

Recall that $\bar{W}$ carries the complex conjugate representation of $SL(2, \mathbb{C})$. E is called *Dirac spinor space* and the image of $SL(2, \mathbb{C})$ under π is naturally isomorphic to the group Spin_0 (See below). Using the volume forms ϵ_{xy} and $\bar{\epsilon}_{x'y'}$, the γ -matrices can be written in the following 4×4 matrix representation (See [PR84, p. 460]):

$$\gamma_a{}^A{}_B = \sqrt{2} \begin{pmatrix} 0 & \sigma_a{}^{zz'} \epsilon_z^x \bar{\epsilon}_{z'y'} \\ \sigma_a{}^{zz'} \bar{\epsilon}_{z'x'} \epsilon_{zy} & 0 \end{pmatrix},$$

where $A = x \oplus x'$ and $B = y \oplus y'$, i.e., both indices A and B can take the four values $11'$, $12'$, $21'$ and $22'$ respectively. An elementary but lengthy calculation verifies that the Clifford relations are satisfied:

$$\gamma_a \gamma_b + \gamma_b \gamma_a = 2\eta_{ab} \mathbf{1}, \quad (7)$$

where $\eta_{ab} = \text{diag}(+1, -1, -1, -1)$ is the Minkowski metric and $\mathbf{1}$ is the identity on E . We remark that Spin_0 mentioned above is the connected component of the identity in the group Spin which consists of all 4×4 matrices S satisfying

$$S \gamma^a S^{-1} = \Lambda^a{}_b \gamma^b$$

for some real constant $\Lambda^a{}_b$. The elements of E are called *Dirac spinors*. The elements of the dual vector-space of E , which is called the Dirac cospinor space (E^*), are called *Dirac cospinors*. Finally smooth mappings from Minkowski space into E (E^*), denoted by $\Gamma(E)$ ($\Gamma(E^*)$), are called *Dirac spinor fields* (*Dirac cospinor fields*). By analogy to the previous subsection, an action of $ISL(2, \mathbb{C})$ on these fields is given by the transformation

$$\Psi^A(q) \mapsto \Psi'^A(q) = S^A{}_B \Psi^B(\mathcal{P}^{-1}(q)) \quad \forall g \in ISL(2, \mathbb{C}), \Psi^A \in \Gamma(E), \quad (8)$$

and

$$\bar{\Psi}_A(q) \mapsto \bar{\Psi}'_A(q) = \bar{\Psi}_B(\mathcal{P}^{-1}(q)) S^{-1}{}^B{}_A \quad \forall g \in ISL(2, \mathbb{C}), \bar{\Psi}_A \in \Gamma(E^*), \quad (9)$$

where $S^A{}_B$ is the homogeneous part of g in the representation given by Eqn. (6), $\mathcal{P}$ denotes the Poincaré element associated to g and q is some point in Minkowski space. Since E and E^* are dual to each other, we can equip the vector spaces $\Gamma(E)$ and $\Gamma(E^*)$ with an (antisymmetric) wedge product $\wedge$. More precisely: For $\Psi^A \in \Gamma(E)$ and $\bar{\Psi}_A \in \Gamma(E^*)$ we define

$$\begin{aligned} \bar{\Psi}_B \wedge \Psi^A : (\Gamma(E) \oplus \Gamma(E^*)) \otimes (\Gamma(E) \oplus \Gamma(E^*)) &\mapsto C^\infty(\mathbb{R}^4) \\ \begin{pmatrix} \phi_1^B \\ \bar{\phi}_{1A} \end{pmatrix} \otimes \begin{pmatrix} \phi_2^B \\ \bar{\phi}_{2A} \end{pmatrix} &\mapsto \bar{\Psi}_B \phi_1^B \cdot \Psi^A \bar{\phi}_{2A} \\ &\quad - \bar{\Psi}_B \phi_2^B \cdot \Psi^A \bar{\phi}_{1A} \end{aligned} \quad (10)$$

Moreover,

$$\begin{aligned} \bar{\Psi}_B(q) \wedge \Psi^A(q') : (\Gamma(E) \oplus \Gamma(E^*)) \otimes (\Gamma(E) \oplus \Gamma(E^*)) &\mapsto \mathbb{C} \\ \begin{pmatrix} \phi_1^B \\ \bar{\phi}_{1A} \end{pmatrix} \otimes \begin{pmatrix} \phi_2^B \\ \bar{\phi}_{2A} \end{pmatrix} &\mapsto \bar{\Psi}_B \phi_1^B(q) \cdot \Psi^A \bar{\phi}_{2A}(q') \\ &\quad - \bar{\Psi}_B \phi_2^B(q) \cdot \Psi^A \bar{\phi}_{1A}(q') \end{aligned} \quad (11)$$

is well defined. In the future we will omit the $\wedge$ and simply write $\bar{\Psi}_B(q)\Psi^A(q')$ or even $\bar{\Psi}_B\Psi^{A'}$. The prime reminds of the fact that A' is a Dirac spinor index referring to the point q' . We obviously have

$$\{\bar{\Psi}_B, \Psi^{A'}\}_+ = \{\bar{\Psi}_B, \bar{\Psi}_{B'}\}_+ = \{\Psi^A, \Psi^{A'}\}_+ = 0,$$

where $\{\cdot, \cdot\}_+$ denotes the anticommutator. In this sense classical Dirac spinor- and cospinor fields are *anti-commuting* objects. Note that $\bar{\Psi}_B\Psi^{A'}$ transforms as a cospinor at p and as a spinor at p' . Such quantities are usually called bi-spinors (See [DB60] for further details). Classical Dirac spinors without further assumptions are suitable for models describing charged fermionic fields. We are now going to define neutral spinor fields which we need later for the definition of the Wess-Zumino model. In the literature neutral spinor fields often are called Majorana spinor fields. We begin with the implementation of a charge conjugation. Majorana spinor fields are the elements of $\Gamma(E)$ which are invariant under this mapping. Consider a fixed but arbitrary four dimensional matrix representation of the Dirac γ -matrices. Since the adjoint representation also satisfy Eqn. (7) there exists a unique matrix β with unit determinant³ that intertwines between these. The four maps

$$\begin{aligned} E_\Lambda : \mathbb{R}^4 &\rightarrow E \\ p &\mapsto \omega_\Lambda, \end{aligned}$$

such that for every point p the four elements ω_Λ , $\Lambda = 1, \dots, 4$ form a basis in E , define a *moving frame*. The dual frame $(E^1, \dots, E^4)$ consists of four maps E^Σ from Minkowski space to E^* satisfying $E^\Lambda(E_\Sigma) = \delta^\Lambda_\Sigma$. Every Dirac spinor field Ψ (Dirac cospinor field $\bar{\Psi}$) can be written as $\Psi = \Psi^\Lambda E_\Lambda$ ($\bar{\Psi} = \bar{\Psi}_\Sigma E^\Sigma$), where Ψ^Λ ($\bar{\Psi}_\Sigma$) are complex valued functions on Minkowski space called *components of Ψ* . The Dirac conjugation is given by the following definition:

DEFINITION 1.1. Let $\Psi^A \in \Gamma(E)$ be a Dirac spinor field with components Ψ^Λ . We define the *Dirac conjugate spinor field* $\Psi^+_A \in \Gamma(E^*)$ by setting its components equal to

$$\Psi^+_\Sigma = \overline{\Psi^\Lambda} \beta_{\Lambda\Sigma}. \quad (12)$$

Here $\overline{}$ denotes complex conjugation and $\beta_{\Lambda\Sigma}$ are the matrix elements of β .

DEFINITION 1.2. The Dirac adjoint of a Dirac cospinor field $\Psi^+ \in \Gamma(E^*)$ is defined by

$$(\Psi^+)^+ := (\beta^*)^{-1\Lambda\Sigma} \overline{\Psi^+_\Sigma} E_\Lambda \quad (13)$$

where $(\beta^*)^{-1\Lambda\Sigma}$ are the matrix elements of the adjoint matrix of β^{-1} .

For the proof that Ψ^+ and $(\Psi^+)^+$ defined by Eqn. (12) and (13) respectively are well defined, observe first that

$$\begin{aligned} (\gamma_a)^* &= (S^{-1}\gamma_b\Lambda(S)^b{}_a S)^* \\ &= S^* \beta \gamma_b \beta^{-1} \Lambda(S)^b{}_a (S^{-1})^* \\ &= S^* \beta S \gamma_a S^{-1} \beta^{-1} (S^{-1})^* \\ &\equiv \beta \gamma_a \beta^{-1}, \end{aligned}$$

since $\det(S^*\beta S) = +1$. Hence for all $S \in \pi(SL(2, \mathbb{C}))$ we have $S^*\beta S = \beta$. Under a change of frames which induces a change of components $\Psi^\Lambda \mapsto \Psi'^\Lambda = S^\Lambda_\Sigma \Psi^\Sigma$ one finds

$$\begin{aligned} (\Psi')^+_\Sigma &:= \overline{\Psi'^\Lambda} \beta_{\Lambda\Sigma} \\ &= \overline{S^\Lambda_\Gamma \Psi^\Gamma} \beta_{\Lambda\Sigma} \\ &= \overline{\Psi^\Gamma} \overline{S^T}^\Lambda_\Gamma \beta_{\Lambda\Sigma} = \overline{\Psi^\Gamma} \beta_{\Gamma\Delta} S^{-1\Delta}_\Sigma = \Psi^+_\Delta S^{-1\Delta}_\Sigma \end{aligned}$$

since $(\overline{S^T})^\Lambda_\Gamma = (S^*)^\Lambda_\Gamma$. This shows that $\Psi^+ := \Psi^+_\Sigma E^\Sigma$ is a frame invariant object. For a cospinor field Ψ^+ , locally given by $\Psi^+ = \Psi^+_\Sigma E^\Sigma$, a similar calculation as above shows that $(\Psi^+)^+$ transforms as a Dirac spinor field under a change of frame, which coincides with Ψ . The charge conjugation is defined next: The transposed Gamma matrices γ^T satisfy the anticommutation relation Eqn. (7). Hence there exists a unique 4×4 matrix C with unit determinant such that $C\gamma_a C^{-1} = -\gamma_a^T$. This matrix is called *charge conjugation matrix*. Note that $C^T = -C = C^{-1}$.

DEFINITION 1.3. For any Dirac spinor field $\Psi \in \Gamma(E)$ we define the *charge conjugate Dirac field* ${}^c\Psi$ by:

$$\Gamma(E) \ni {}^c\Psi = C^{\Lambda\Sigma} \Psi^+_\Sigma E_\Lambda$$

where $C^{\Lambda\Sigma}$ are the matrix elements of C and Ψ^+_Σ are the components of the Dirac conjugate spinor field of Ψ .

³In the literature one usually considers only those representations of the Dirac γ -matrices, where β is identical γ^0

For the proof that this definition is sensible, use a similar argument as above to show that for all $S \in \pi(SL(2, \mathbb{C}))$ we have $CS = S^{-1T}C$ or equivalently $SC = C(S^{-1})^T$. Under a change of frames

$$\begin{aligned} ({}^c\Psi')^\Lambda &= C^{\Lambda\Delta}\Psi^+_{\Gamma}S^{-1\Gamma}_{\Delta} \\ &= C^{\Lambda\Delta}(S^{-1T})^{\Gamma}_{\Delta}\Psi^+_{\Gamma} \\ &= S^{\Lambda}_{\Sigma}C^{\Sigma\Delta}\Psi^+_{\Delta} \\ &= S^{\Lambda}_{\Sigma}({}^c\Psi)^{\Sigma} \end{aligned}$$

which shows that ${}^c\Psi := ({}^c\Psi)^\Lambda E_\Lambda \in \Gamma(E)$ is a frame invariant object.

DEFINITION 1.4. Dirac spinor fields which are invariant under charge conjugation are called *Majorana spinor fields*.

In the remaining part of this chapter we will use Majorana spinor fields exclusively. To shorten the notation we will omit the word ‘Majorana’ in the future whenever it appears unnecessary.

2. The internal symmetries of the classical free Wess-Zumino Model

The Wess-Zumino model consists of three fields, two Spin-0 fields (A, B) one of which is scalar (A), the other (B) is pseudo-scalar and the third (Ψ) is a Majorana spinor field. The free Lagrangian density of this model is [WZ74]:

$$\mathcal{L} = \frac{1}{2}(\nabla A)^2 - \frac{1}{2}m^2 A^2 + \frac{1}{2}(\nabla B)^2 - \frac{1}{2}m^2 B^2 + \frac{1}{2}\Psi^+(i\overleftrightarrow{\not{\nabla}} - m)\Psi, \quad (14)$$

where $m > 0$ is the mass of the fields and $+$ denotes Dirac conjugation. Note that all fields have *equal* masses. The equations of motion resulting from (14) are the Klein-Gordon and Dirac equations respectively.

$$(\square + m^2)A = 0 \quad (15a)$$

$$(\square + m^2)B = 0 \quad (15b)$$

$$(i\overleftrightarrow{\not{\nabla}} - m)\Psi = 0 \quad (15c)$$

$$\Psi^+(i\overleftarrow{\not{\nabla}} + m) = 0 \quad (15d)$$

Besides the usual Poincaré symmetry the model possesses an internal symmetry, which is called *supersymmetry*. The latter transforms Bose to Fermi fields and vice versa. To see this explicitly let us consider the following transformations of the basic fields:

$$A \mapsto A' = A + \delta A = A + \epsilon^+ \Psi \quad (16a)$$

$$B \mapsto B' = B + \delta B = B - i\epsilon^+ \gamma^5 \Psi \quad (16b)$$

$$\Psi \mapsto \Psi' = \Psi + \delta \Psi = \Psi - (i\overleftrightarrow{\not{\nabla}} + m)(A - i\gamma^5 B)\epsilon, \quad (16c)$$

where ϵ is a x -independent spinor field. The resulting transformation of $\mathcal{L}$ up to first order

$$\delta\mathcal{L} = \mathcal{L}(A', B', \Psi') - \mathcal{L}(A, B, \Psi) = \nabla^a \left(\nabla_a A \delta A + \nabla_a B \delta B - \frac{1}{2} \Psi^+ i \gamma_a \delta \Psi \right) =: \nabla^a v_a \quad (17)$$

is a pure divergence. This means that the action $S = \int d^4x \mathcal{L}$ is left invariant, hence Eqn. (16a)–(16c) is indeed a symmetry transformation. For the proof simply insert the definitions and use the Fierz rearrangement formulas of Appendix C:

$$\begin{aligned} \delta \mathcal{L} &= \mathcal{L}(A', B', \Psi',) - \mathcal{L}(A, B, \Psi) \\ &= \left(\frac{1}{2} (\nabla(A + \delta A))^2 - \frac{m^2}{2} (A + \delta A)^2 + \frac{1}{2} (\nabla(B + \delta B))^2 - \frac{m^2}{2} (B + \delta B)^2 \right. \\ &\quad \left. + \frac{1}{2} (\Psi + \delta \Psi)^+ (i \overleftrightarrow{\nabla} - m) (\Psi + \delta \Psi) \right) \\ &\quad - \left(\frac{1}{2} (\nabla A)^2 - \frac{m^2}{2} A^2 + \frac{1}{2} (\nabla B)^2 - \frac{m^2}{2} B^2 + \frac{1}{2} \Psi^+ (i \overleftrightarrow{\nabla} - m) \Psi \right) \\ &= \nabla_a A \nabla^a \delta A - m^2 A \delta A + \nabla_a B \nabla^a \delta B - m^2 B \delta B - \frac{1}{2} \Psi^+ (i \overleftrightarrow{\nabla} - m) \delta \Psi + \frac{1}{2} (\delta \Psi)^+ (i \overleftrightarrow{\nabla} - m) \Psi \end{aligned}$$

(1. order)

$$\begin{aligned} &= \nabla^a (\nabla_a A \delta A + \nabla_a B \delta B) \\ &\quad - \square A \delta A - m^2 A \delta A - \square B \delta B - m^2 B \delta B + \frac{1}{2} \Psi^+ (i \overleftrightarrow{\nabla} - m) \delta \Psi - \frac{1}{2} \Psi^+ (i \overleftrightarrow{\nabla} + m) \delta \Psi \end{aligned}$$

by (118d)

$$\begin{aligned} &= \nabla^a (\nabla_a A \delta A + \nabla_a B \delta B - \frac{i}{2} \Psi^+ \gamma_a \delta \Psi) \\ &\quad - (\square + m^2) A \Psi^+ \epsilon + i(\square + m^2) B \Psi^+ \gamma^5 \epsilon + (\square + m^2) A \Psi^+ \epsilon - i(\square + m^2) B \Psi^+ \gamma^5 \epsilon \end{aligned}$$

using (118c) and since $(i \overleftrightarrow{\nabla} - m) \delta \Psi = (\square + m^2)(A - i \gamma^5 B) \epsilon$

$$\begin{aligned} &= \nabla^a (\nabla_a A \delta A + \nabla_a B \delta B - \frac{i}{2} \Psi^+ \gamma_a \delta \Psi) \\ &=: \nabla^a v_a \end{aligned}$$

REMARK . It is a generic feature of supersymmetry transformations, in contrast to other symmetries, e.g., Poincaré invariance, that they produce a nontrivial variation of the Lagrangian density. The reader should note that Eqn. (17) is obtainable from Eqn. (14) without using the field equations (15a)–(15d).

Using Noether's theorem one obtains a conserved current associated with the transformation (16a)–(16c): It is given by:

$$i^a := v^a - \frac{\partial \mathcal{L}}{\partial (\partial_a \Phi)} \delta \Phi \quad \Phi = (A, B, \Psi)$$

Inserting

$$\frac{\partial \mathcal{L}}{\partial (\partial_a A)} = \partial^a A, \quad \frac{\partial \mathcal{L}}{\partial (\partial_a B)} = \partial^a B, \quad \frac{\partial \mathcal{L}}{\partial (\partial_a \Psi)} = \frac{i}{2} \Psi^+ \gamma^a$$

one obtains –using the Fierz rearrangement identities again–:

$$\begin{aligned} i^a &= -i \Psi^+ \gamma^a \delta \Psi \\ &= i \Psi^+ \gamma^a (i \overleftrightarrow{\nabla} + m) (A - i \gamma^5 B) \epsilon \\ &= \epsilon^+ i (A - i \gamma^5 B) (i \overleftrightarrow{\nabla} - m) \gamma^a \Psi \\ &=: \epsilon^+{}_A k^{aA} \end{aligned} \tag{18}$$

The current k^{aA} is usually called *supercurrent*. Note that it carries –besides the usual vector index– an additional spinor index, hence it is a mixed spinor vector field. Its Dirac conjugate is

$$k^{+a}{}_A = i \Psi^+ \gamma^a (i \overleftrightarrow{\nabla} + m) (A - i \gamma^5 B) \tag{19}$$

OBSERVATION .

1. The charge Q associated to the Noether current i^a induces the symmetry transformation Eqn. (16a)–(16c).
2. The symmetry transformation is a realization of the supersymmetry algebra.

PROOF. 1. The classical charge Q_ϵ associated with i^a is defined by

$$Q_\epsilon := \int_\Sigma k^{+a} \epsilon d\sigma_a, \quad (20)$$

where Σ is the $t = 0$ hypersurface. Recall that Q_ϵ is actually independent of t and Σ , since k^a is a conserved current. To see that Q_ϵ generates the symmetry transformations Eqn. (16a) - (16c) we introduce the canonical momenta for the three fields: For $t = t_0$

$$\begin{aligned} \Pi_A &:= \frac{\partial \mathcal{L}}{\partial(\partial_0 A)} = \partial^0 A \\ \Pi_B &:= \frac{\partial \mathcal{L}}{\partial(\partial_0 B)} = \partial^0 B \\ \Pi_\Psi &= i\Psi^+ \gamma^0 \end{aligned}$$

Note that the last equation does *not* read $\Pi_\Psi = \frac{i}{2}\Psi^+ \gamma^0$ (Wrong !), since Ψ is a Majorana spinor field, i.e., Ψ and Ψ^+ are *not* independent. (See for example the discussion in [Rom60, pp. 306]). Inserting these definitions in Eqn. (20) we obtain

$$\begin{aligned} Q_\epsilon(A, \nabla A, B, \underline{\nabla} B, \Psi, \underline{\nabla} \Psi, \Pi_A, \Pi_B, \Pi_\Psi) \\ = \int_\Sigma d^3\sigma \Pi_\Psi (i\gamma^0 \Pi_A + \gamma^0 \gamma^5 \Pi_B + (i\underline{\nabla} + m)(A - i\gamma^5 B)) \epsilon, \end{aligned} \quad (21)$$

where the underscore denotes spatial (i.e. three dimensional) expressions. Note that Q_ϵ is bilinear in it's variables. We now compute the Poisson brackets of Q_ϵ with the fields⁴

$$\begin{aligned} \{Q_\epsilon, A\} &:= \int_\Sigma d^3\sigma \sum_\Phi \frac{\partial Q}{\partial \Phi} \frac{\partial A}{\partial \Pi_\Phi} - \frac{\partial Q}{\partial \Pi_\Phi} \frac{\partial A}{\partial \Phi} \\ &= -\Pi_\Psi i\gamma^0 \epsilon = \Psi^+ \gamma^0 \gamma^0 \epsilon = \epsilon^+ \Psi \equiv \delta A \end{aligned} \quad (22)$$

$$\{Q_\epsilon, B\} = -\Pi_\Psi \gamma^0 \gamma^5 \epsilon = -i\Psi^+ \gamma^5 \epsilon = -i\epsilon^+ \gamma^5 \Psi \equiv \delta B \quad (23)$$

$$\{Q_\epsilon, \Psi\} = -(\underline{\nabla} + m)(A - i\gamma^5 B)\epsilon \equiv \delta \Psi \quad (24)$$

$$(25)$$

This shows that Q_ϵ indeed generates the supersymmetry transformation at the classical level.

2. To establish the second statement, we investigate the algebra of the supersymmetry charge:

$$\{\{Q_{\epsilon_1}, Q_{\epsilon_2}\}, \cdot\}, \quad (26)$$

where the $\cdot$ denotes one of the basic fields. Let us evaluate these expressions explicitly using the Jacobi identity and the Fierz rearrangement formulas. We start with the basic field A :

$$\begin{aligned} \{\{Q_{\epsilon_1}, Q_{\epsilon_2}\}, A\} &= \{Q_{\epsilon_1}, \{Q_{\epsilon_2}, A\}\} - \{Q_{\epsilon_2}, \{Q_{\epsilon_1}, A\}\} \\ &= \{Q_{\epsilon_1}, \epsilon_2^+ \Psi\} - \{Q_{\epsilon_2}, \epsilon_1^+ \Psi\} \\ &= -\epsilon_2^+ (i\underline{\nabla} + m)(A - i\gamma^5 B)\epsilon_1 + \epsilon_1^+ (i\underline{\nabla} + m)(A - i\gamma^5 B)\epsilon_2 \\ &= -2i\epsilon_2^+ \underline{\nabla} \epsilon_1 A \end{aligned}$$

Similarly for B :

$$\{\{Q_{\epsilon_1}, Q_{\epsilon_2}\}, B\} = -2i\epsilon_2^+ \underline{\nabla} \epsilon_1 B$$

Finally let us calculate $\{\{Q_{\epsilon_1}, Q_{\epsilon_2}\}, \Psi\}$:

$$\begin{aligned} \{\{Q_{\epsilon_1}, Q_{\epsilon_2}\}, \Psi\} &= \{Q_{\epsilon_1}, \{Q_{\epsilon_2}, \Psi\}\} - \{Q_{\epsilon_2}, \{Q_{\epsilon_1}, \Psi\}\} \\ &= \{Q_{\epsilon_1}, (-1)(i\underline{\nabla} + m)(A - i\gamma^5 B)\epsilon_2\} \\ &\quad - \{Q_{\epsilon_2}, (-1)(i\underline{\nabla} + m)(A - i\gamma^5 B)\epsilon_1\} \\ &= -(i\underline{\nabla} + m)(\epsilon_1^+ \Psi \epsilon_2 - \gamma^5 \epsilon_1^+ \gamma^5 \Psi \epsilon_2) - (1 \leftrightarrow 2) \end{aligned} \quad (27)$$

To evaluate the last term we use the following identity for products of spinors and arbitrary 4×4 matrices N and M [Wes90]:

$$\Psi_1^+ M \Psi_3 \Psi_4^+ N \Psi_2 = -\frac{1}{4} \sum_R \sigma(R) \Psi_1^+ O_R \Psi_2 \Psi_4^+ N O_R M \Psi_3,$$

⁴For simplicity we consider a representation of the Dirac γ -matrices, where $\beta \equiv \gamma^0$.

where

$$\sigma(R) = \begin{cases} +1 \\ -1 \\ +1 \end{cases} \quad \text{for } O_R = \begin{cases} 1, \gamma^5, \gamma^a \\ \gamma^5 \gamma^a \\ \frac{1}{4}(\gamma^a \gamma^b - \gamma^b \gamma^a) \equiv G^{[ab]} \end{cases}$$

$$\begin{aligned} (27) &= -i\epsilon_1^+ \nabla_a \Psi \gamma^a \epsilon_2 - m\epsilon_1^+ \Psi \epsilon_2 + i\epsilon_1^+ \gamma^5 \nabla_a \Psi \gamma^a \gamma^5 \epsilon_2 + m\epsilon_1^+ \gamma^5 \Psi \gamma^5 \epsilon_2 - (1 \leftrightarrow 2) \\ &= -\frac{1}{4} \sum_R -i\sigma(R) \epsilon_1^+ O_R \epsilon_2 \gamma^a (O_R - \gamma^5 O_R \gamma^5) \nabla_a \Psi \\ &\quad - m\sigma(R) \epsilon_1^+ O_R \epsilon_2 (O_R - \gamma^5 O_R \gamma^5) \Psi - (1 \leftrightarrow 2) \end{aligned}$$

Note that the l.h.s. of Eqn. (27) is antisymmetric in ϵ_1 and ϵ_2 and the only antisymmetric terms on its r.h.s. involving ϵ_1 and ϵ_2 are $\epsilon_1^+ \gamma^a \epsilon_2$ and $\gamma^5 G^{[ab]} \gamma^5$. But $\gamma^5 G^{[ab]} \gamma^5 = G^{[ab]}$ so that the brackets are zero in this case. We get

$$\begin{aligned} (27) &= \sum_b i\epsilon_1^+ \gamma^b \epsilon_2 \gamma^a \gamma^b \nabla_a \Psi + m\epsilon_1^+ \gamma^b \epsilon_2 \gamma^b \Psi \\ &= \sum_b \epsilon_1^+ \gamma^b \epsilon_2 (-\gamma^b ((i\nabla - m)\Psi) + 2i\eta^{ab} \nabla_a \Psi) \\ &= -2i\epsilon_2^+ \nabla \epsilon_1 \Psi \end{aligned}$$

On the other hand the derivative ∇_a is the generator of the translations. Therefore the Poisson brackets of the supercharge generate the translations, i.e., Q_ϵ is the ‘square-root’ of the 4-momentum $P_a = i\nabla_a$ and

$$\{Q_{\epsilon_1}, Q_{\epsilon_2}\} = -2\epsilon_2^+ \gamma^a P_a \epsilon_1 \quad (28)$$

□

3. Normal Ordering

We will show in this section that the momentum operator and the Hamiltonian of the quantized Wess-Zumino model are a priori normal ordered, i.e., that their vacuum expectation values vanish. Moreover the stress energy tensor of the model can be written as an operator valued distribution on the vacuum Hilbertspace in terms of the supercurrent. We start with the supercurrent. It is worth noting that Eqn. (18) is well defined after quantization, since the spectrum condition ensures that products of *different* fields at the same spacetime point exist. For the adjoint of the supercurrent one obtains:

$$\begin{aligned} i_\epsilon^{a*} &= (\epsilon^+ i(A - i\gamma^5 B)(i\nabla - m)\gamma^a \Psi)^* \\ &= \Psi^+ \gamma^a (i\nabla + m)(A - i\gamma^5 B)\epsilon \\ &= k^+{}^a \epsilon \\ &= \epsilon^+ k^a = i_\epsilon^a, \end{aligned}$$

since k^a is a Majorana spinor. Hence j_ϵ^a is a conserved hermitian current. Let $f_R \in C_0^\infty(\mathbb{R}^3)$ and $f_T \in C_0^\infty(\mathbb{R})$ be two smooth function with compact support, such that

$$f_R(x) = \begin{cases} 1 & |x| < R \\ f_R(|x|) & R \leq |x| \leq R + \Lambda \\ 0 & |x| > R + \Lambda \end{cases}$$

for some $\Lambda \in \mathbb{R}$, and

$$f_T : \begin{cases} \int f_T(x_0) dx_0 = 1 \\ f_T(x_0) = 0 & |x_0 - t| > T \\ f_T(x_0) = f_T(-x_0 + 2t) \end{cases}$$

Then $Q_{\epsilon RT} = j_\epsilon^0(f_R f_T)$ is a well defined operator. We have the following two theorems

THEOREM 1.5 (Theorem 6.2.3 of [Lop91]). *Let A_L be a localized operator, then*

$$C_\epsilon(A_L) = [Q_{\epsilon RT}, A_L]_-$$

applied to any state in the set

$$D_\Omega = \{u \in H \mid u \text{ is generated by polynomials of the smeared fields applied to the vacuum } \Omega\}$$

is independent of both f_R and f_T , whenever j_ϵ^a is a conserved current and R is sufficiently large. $C_\epsilon(A_L)$ is a localized operator.

THEOREM 1.6 (Theorem 6.2.8 and 6.2.9 of [Lop91]). *Assume the existence of a mass gap and the local conservation of the current; then the limit $\lim_{R \rightarrow \infty} (u, Q_{\epsilon RT} v) =: Q_\epsilon(u, v)$ exists for all quasi-local vectors u and v and defines a linear operator $\epsilon^+ G$ on the Hilbertspace:*

$$Q_\epsilon(u, v) = (u, \epsilon^+ G v) \quad \forall u, v \in H, v \text{ quasilocal}$$

REMARK .

- The charge $\epsilon^+ G$ is hermitian due to the fact that the corresponding current is. Moreover, since i_ϵ^a is translational covariant, it can be proved that $\epsilon^+ G$ is self-adjoint with a dense invariant domain D_Ω^{ex} , consisting of all states with finite number of particles and smooth momentum space wave-functions having compact support.
- By Theorem 6.2.12 of [Lop91] one finds $(\Omega, [Q_{\epsilon RT}, A_L]_- \Omega) = 0$ for all localized operators and sufficiently large R . It follows that $\epsilon^+ G$ annihilates the vacuum, i.e., supersymmetry is unbroken and on the vacuum $\lim_{R \rightarrow \infty} [Q_{\epsilon RT}, A_L]_- \Omega = [\epsilon^+ G, A_L]_- \Omega = \epsilon^+ G A_L \Omega$. The relation between currents and charges for models with spontaneously broken symmetries is closely investigated by Buchholz, Doplicher, Longo and Roberts [BDLR92].

Let us now take a closer look at the charge operators: Q_ϵ is bilinear in the basic fields. Thus the results for the classical charges

obtained in the previous section translate to the quantum field theoretical case:

$$\begin{aligned} [\epsilon_1^+ G, \epsilon_2^+ G]_- &= [\epsilon_1^+ G, G^+ \epsilon_2]_- && \text{since } G \text{ transforms as a Majorana spinor} \\ &= \epsilon_1^+ [G, G^+]_+ \epsilon_2 && \epsilon_1^+ \text{ and } \epsilon_2 \text{ anticommute} \\ &= 2\epsilon_1^+ \gamma^a P_a \epsilon_2 && \text{by Eqn. (28)} \end{aligned}$$

It is therefore possible to write:

$$\{G^A, G_B^+\}_+ = 2\gamma^{aA}{}_B P_a \quad (29)$$

Now multiply Eqn. (29) by γ^0 and take the trace over the spinor indices:

$$\begin{aligned} \text{Tr} (\{G^A, G_B^+\}_+ \gamma^{0B}{}_C) &= G^A \bar{G}_B^T \gamma^{0B}{}_C \gamma^{0C}{}_A + \bar{G}_B^T \gamma^{0B}{}_C \gamma^{0C}{}_A G^A \\ &= G^A G_A^\dagger + \text{h.c.} \\ &= 8P_0, \end{aligned}$$

where $\text{Tr} (\gamma^a \gamma^b) = 4\eta^{ab}$ and $\gamma^{02} = 1$ were used. As usual † denotes hermitian conjugation. Since $G^A G_A^\dagger + \text{h.c.}$ is a positive operator, we conclude that P_0 –which is the Hamiltonian– is positive too. A similar calculation yields

$$0 = \text{Tr} \langle \Omega | \{G, G^+\}_+ \gamma^a \Omega \rangle = 8 \langle \Omega | P^a \Omega \rangle,$$

which ensures that all components of P^a are normal ordered. We remind the reader that P^a which is the generator of the translations is a well defined self adjoint operator for the massive Wess-Zumino model. Heuristically this normal ordering feature is due to the fact that bosonic and fermionic loops have opposite sign. Their infinite contributions to the vacuum expectation values cancel each other in models with equally many bosonic and fermionic degrees of freedom. In view of Eqn. (29) the energy momentum tensor θ^{ab} of the Wess-Zumino model can be written as (see e.g. [FZ75])

$$\theta^{ab} = +\frac{1}{16} \text{Tr} (\gamma^a \{k^b, G^+\}_+ + \gamma^b \{k^a, G^+\}_+), \quad (30)$$

where the spinor indices on γ^a , k^a and G^+ were suppressed⁵. (Kraus and Landau loc.cit. showed that the anti-commutator in Eqn (30) is well defined on a suitable subset of D_Ω^{ex}). It follows that the vacuum expectation value of θ^{ab} vanishes. Recall that the supersymmetry is not spontaneously broken, i.e., the supercharge G^+ annihilates the vacuum. In remaining parts of this work the question whether this tameness of the vacuum expectation value of the energy momentum tensor of supersymmetric quantum field theories can be extended to curved space times is investigated. Such a priori renormalization is a highly desirable feature from a general relativistic viewpoint; contrary to special relativity where only differences of energy are measurable, we have measurable effects induced by the absolute value of the energy, in the latter, namely

⁵The explicit proof of Eqn. (30) requires some lengthy calculations. Due to the fact that a similar identity on a manifold will be used and proved later in this thesis, the actual proof of (30) is omitted.

energy curves spacetime. It was already stated in the introduction, that supersymmetry on curved spacetime, to be implemented consistently, leads to supergravity theories in general [Nie81]. Therefore we will only mimic the *structure* of the Wess-Zumino model on a curved spacetime. If we restrict ourself to globally hyperbolic manifolds, i.e., to spacetimes where a rigorous quantization of various free fields has been done [Dim80, Dim82, Dim92], there seems to be no known nontrivial example of a manifold which possesses a true supersymmetric extension of its isometry group. Moreover, in the general case a Lorentz manifold does not possess an isometry group at all. Coming back to the *structure* of a supersymmetric theory one observes immediately that fermionic *and* bosonic fields are needed. (Recall that a supersymmetric transformation changes half integer spin fields to integer spin fields and vice versa). The latter exist on all Lorentz manifolds, whereas for the existence of fermionic fields special conditions on the topology of the manifold must be satisfied [Mil63]. On a globally hyperbolic manifold however a Theorem due to Geroch [Ger68, Ger70] ensures the existence of half integer spin fields in four dimensions (see below). Nevertheless let us begin with a closer look on this topological condition preventing the existence of spinor fields, because there are serious attempts to extend the result of Dimock loc. cit. and others to non globally hyperbolic spacetimes (See [Kay92]). Moreover, there are some cosmologically interesting solutions of the Einstein Equations –for example plain waves– which fail to be globally hyperbolic.

Spinors on Lorentz manifolds

For the definition of spinors on Minkowski spacetime it was important that the Poincaré group is the isometry group of this manifold. It was already mentioned above that in the general case there might exist no isometry group at all. Therefore the study of isometries does not seem to be a good starting point for a definition¹ of spinors, since we would restrict ourself to just a few very special cases. Let us therefore return to the principle of ‘special covariance’: Like on Minkowski space, on a general manifold we can identify an observer with her measurement device at a point p with an orthonormal base system (vierbein) at this point. Different observers at the *same* point will be described by different vierbeins. Since a vierbein consists of four linearly independent orthogonal tangent vectors at p , two different vierbeins are connected by a Lorentz transformation². Thus to describe spinors we will start with the proper Lorentz group, not with the whole Poincaré group, as before. We are going to define spinors in such a way that every transformation of a vierbein ($\hat{=}$ change of observer) results in a corresponding spinor transformation up to sign. The sign ambiguity evaporates—as in Minkowski space—as soon as a continuous curve of transformations starting from the identity is specified. Mathematically this idea is described best in terms of fiber bundles: Spinors are sections in a suitable associated fiber bundle, called spinor bundle. The corresponding principal fiber bundle is a special twofold covering of the principal fiber bundle of the vierbeins. The existence of this covering is a topological property of the underlying manifold. In the first section of this chapter for the convenience of the reader the definition of a principal fiber bundle is recalled and we define the spinor bundle. Some facts on the preferred connection on the latter are presented in the next section. For the proofs that were omitted, as well as for further details, the reader is referred to the book of Kobayashi and Nomizu [KN63]. We continue by defining Čech-cohomology groups and Stiefel-Whitney classes according to the book of Nakahara [Nak90]. The latter are shown in section 5 to measure the existence of a spin structure as Milnor already found out in 1963 [Mil63]. We close this chapter by summarizing some other, but also well known conditions on the existence of spin structures which turn out to be more useful for applications.

1. Bundle-formalism for spinors

Let $\mathcal{G}$ denote a Lie-group, P a manifold and $L : \mathcal{G} \times P \rightarrow P$ a smooth mapping.

DEFINITION 2.1. The mapping L is called *left action*: $\Leftrightarrow$

1. $\forall g \in \mathcal{G}$ fixed, $L_g : P \rightarrow P$ is a diffeomorphism.
2. $\forall g_1, g_2 \in \mathcal{G}, L_{g_1} \circ L_{g_2} \equiv L_{g_1 g_2}$.

DEFINITION 2.2. A left action is called *free*: $\Leftrightarrow \forall g \in \mathcal{G}, u \in P, g \neq e$ one has $L_g(u) \neq u$, i.e., L_g has no fix-points.

DEFINITION 2.3. For all points $u \in P$ the set $O = \{L_g(u) \mid g \in \mathcal{G}\}$ is called the *orbit* of u .

REMARK . P can be represented as a disjoint union of orbits.

DEFINITION 2.4. A *principal fiber bundle* $(P, \mathcal{G}, M, L)$ consists of a manifold P , the bundle manifold, a Lie group $\mathcal{G}$ (fiber group), a second manifold M (base manifold) and a free left action L , which fulfills the following conditions.

1. The orbits of L are in one-to-one correspondence to the points of M and the projection

$$\pi : P \rightarrow M$$

is surjective and smooth.

2. For all $x \in M$ there exists an open neighborhood U of x such that there is a diffeomorphism

$$\Psi : P \supset \pi^{-1}(U) \rightarrow \mathcal{G} \times U$$

and the left action of $\mathcal{G}$ on $\pi^{-1}(U)$ equals the left multiplication on $\mathcal{G} \times U$ (Let $\Psi(u) = (g, x)$, then $\Psi(L_{g'}u) = \Psi(g'g, x)$)

¹In view of this E. Cartan suggested in [Car66] that a sensible definition of spinors on manifolds is not possible.

²We continue to restrict ourself to *proper* Lorentz transformations, since there are experimental evidences (see Chapter 1) that our universe is time and space orientable. Moreover, we can distinguish these orientations by performing T and P asymmetry experiments.

Let $(P, \mathcal{G}, M, \mathbf{L})$ be a principal fiber bundle, F another manifold, and $\chi: \mathcal{G} \times F \rightarrow F$ a second left action, which is not necessarily assumed to be free. Then the associated fiber bundle B is defined by:

DEFINITION 2.5. Consider the product manifold $P \times F$. On this manifold

$$\xi_g(u, f) = (\mathbf{L}_g(u), \chi_g(f))$$

defines a left action. Let B denote the set of all orbits of $\mathcal{G}$ on $P \times F$. By the construction there exists a natural projection $\tilde{\pi}$ from E to M

$$\begin{aligned} \tilde{\pi}: E &\rightarrow M \\ y = [u, f]_{\sim} &\mapsto \pi(u), \end{aligned}$$

where $[\cdot, \cdot]_{\sim}$ denotes the orbit of $(u, f) \in P \times F$. We equip B with the structure of a topological manifold by demanding that for any neighborhood U satisfying condition (2) of Definition 2.4, $\tilde{\pi}^{-1}(U) \subset B$ is diffeomorphic to $F \times U$. $B = (P \times F)_{/\mathcal{G}}$ is called the *associated fiber bundle* to $(P, \mathcal{G}, M, \mathbf{L})$.

REMARK . It follows that $\tilde{\pi}^{-1}(x)$ is diffeomorphic to F , that is, B equals $M \times F$ locally.

We have now available all tools for the definition of spinors on a manifold. Consider the principal fiber bundle $FM = (P, \mathcal{L}_0^\uparrow, M, \mathbf{L})$ which is the bundle consisting of all oriented, time oriented orthonormal frames on (M, g_{ab}) together with the proper Lorentz group $\mathcal{L}_0^\uparrow$. Denoting its universal covering group by Spin_0 we have

DEFINITION 2.6. An oriented, time oriented Lorentz manifold (M, g_{ab}) admits a *spin structure*: $\Leftrightarrow$ There exists a Spin_0 -principal fiber bundle $SM = (\hat{P}, \text{Spin}_0, M, \hat{\mathbf{L}})$ over M , called the *spin frame bundle*, and a bundle homomorphism $p: SM \rightarrow FM$ preserving the base points which satisfies

$$p \circ \hat{\mathbf{L}}_S = \mathbf{L}_{\Lambda(S)} \circ p$$

Here $\Lambda(S) \in \mathcal{L}_0^\uparrow$ is the element of $\mathcal{L}_0^\uparrow$ associated to $S \in \text{Spin}_0$.

REMARK . It was already mentioned above that not all manifolds admit a spin structure. Moreover, spin structures are not unique in general. The existence of a spin structure is a topological property of the manifold, i.e., it is independent of the Lorentz metric. We will come back to this point in section 5 of this chapter.

DEFINITION 2.7. The *spinor bundle* is the associated vector bundle $DM = (SM \times \mathbb{C}^4)_{/\text{Spin}_0}$. Spinor fields are sections in DM , i.e., C^∞ -mappings $u: M \rightarrow DM$, such that $u(x) \in D_x M$, where $D_x M$ denotes the fiber over $x \in M$. The dual vector bundle D^*M is called *cospinor bundle*, its sections are called *cospinor fields*.

In order to relate this bundle theoretical formulation with the intuitive ideas of a spinor, consider a local cross section $E: M \supset U \rightarrow SM$ in the spin structure. This section uniquely determines a ‘moving frame’ $(E_1, \dots, E_4)$ in the spinor bundle DM : The mappings $E_\Lambda: U \rightarrow DM$ are local crossections in the spinor bundle given by

$$E_\Lambda(x) = i(E(x), \omega_\Lambda),$$

where $i: SM \times \mathbb{C}^4 \rightarrow DM$ is the natural embedding of $(SM \times \mathbb{C}^4)$ into DM and ω_Λ , $\Lambda = 1, \dots, 4$ denote a basis in $\mathbb{C}^4$. It follows that every spinor field $u \in \Gamma(DM)$ can be written locally as

$$u = \sum_{\Lambda=1}^4 u^\Lambda E_\Lambda$$

with smooth functions u^Λ . Under a change of sections in SM , given by $E \mapsto E' = \hat{\mathbf{L}}_S(E)$, where $\hat{\mathbf{L}}_S$ is the left-action of $S \in \text{Spin}_0$ on SM , the ‘moving frames’ transform as

$$E_\Lambda \mapsto E'_\Lambda = E_\Gamma (S^{-1})^\Gamma_\Lambda.$$

Consequently the components of a spinor field u transform according to

$$u^\Lambda \mapsto u'^\Lambda = (S)^\Lambda_\Gamma u^\Gamma \quad (31)$$

On the other hand by the bundle-homomorphism p every change of section in the spin structure induces a change of sections in the vierbein bundle FM , i.e., the vierbeins in every fiber are undergoing a Lorentz transformation. In this sense we found

A spinor $u \in \Gamma(DM)$ transforms under a change of observers ($\hat{=}$ change of section in the spin structure) according to a (reducible) representation of the group Spin_0 , the covering group of the Lorentz group.

2. The spinor connection

It is well known that on the tangent bundle of a Lorentz manifold there exists a preferred connection, called Levi-Civita connection. In this section it is shown how this connection can be used to define a preferred connection in the spinor bundle. It is worth noting that this result depends crucial on the fact that the fiber groups of both bundles, i.e., $\mathcal{L}_0^\uparrow$ and Spin_0 , have the same Lie-algebra. Let us begin with two different characterizations of the notion of a ‘connection’ in a principal fiber bundle. Let (M, g_{ab}) denote a pseudo-Riemannian manifold of dimension d , $\mathcal{G}$ a Lie-group and $(P, M, \mathcal{G}, \mathbf{R})$ a principal fiber bundle over M .

DEFINITION 2.8. Let u be a point in $(P, M, \mathcal{G}, \mathbf{L})$ and let $T_u P$ denote the tangent space of P at u . A *connection* Ξ in P is an assignment of a subspace Q_u of $T_u P$ for every point $u \in P$, such that

1. $T_u P = Q_u + G_u$ (direct sum), where $G_u \subset T_u P$ is the subspace tangential to the fiber $\pi^{-1}(\pi(u))$; π denotes the projection to the base point.
2. $Q_{au} = (\mathbf{L}_a)_* Q_u$ ($\mathbf{L}_a$ is the left action of $\mathcal{G}$)
3. The mapping $u \mapsto Q_u$ is smooth

Q_u is called *horizontal*, whereas G_u is called *vertical* subspace of $T_u P$, i.e., a connection uniquely assigns to every element $t \in T_u P$ its horizontal and vertical components, denoted by Ht and Vt respectively.

Note that this Definition yields for every vector-field on M a unique lift, i.e., it allows the parallel transport of vectors along curves in M .

THEOREM 2.9. Let $(P, M, \mathcal{G}, \mathbf{L})$ and Ξ be given as above and let $X \in \Gamma(TM)$ be a vector-field on M , then there exists a unique $\mathcal{G}$ invariant lift $\tilde{X} \in \Gamma(TP)$ of X .

For the proof we refer the reader to [KN63, pp. 65]. As an immediate consequence one finds

LEMMA 2.10. Let Ξ be a connection in $(P, M, \mathcal{G}, \mathbf{L})$, $\gamma: [0, 1] \rightarrow M$ a curve in M , then there exists a diffeomorphism

$$T_\gamma: \pi^{-1}(\gamma(0)) \rightarrow \pi^{-1}(\gamma(1)),$$

which is called parallel transport along γ .

PROOF. Let $u \in \pi^{-1}(\gamma(0))$ be a point in the principal fiber bundle with base point $\gamma(0)$. We define $T_\gamma u$ to be the element in the fiber $\pi^{-1}(\gamma(1))$ for which $T_\gamma u = \tilde{\gamma}(1)$; $\tilde{\gamma}$ denotes the unique lift of γ with $\tilde{\gamma}(0) = u$. We remark that the horizontal lift of a curve is the integral curve of the horizontal lift of its tangent vector field. On the other hand the Definition of a parallel transport uniquely fixes the connection: Let γ be a curve in M , $u \in \pi^{-1}(\gamma(0))$ a point in the fiber bundle and $t \in T_u P$ be an element in the tangent space of u , such that

$$d\pi(t) = \dot{\gamma}(0) \equiv \dot{\gamma}(0).$$

Then the horizontal part Ht of t is defined as follows: Let $\tilde{\gamma}$ be the lift of γ with $\tilde{\gamma}(0) = u$. Then by assumption $\tilde{\gamma}$ denotes the parallel transport of u along γ and the horizontal part of t is defined by $Ht := \tilde{\gamma}_*(0)$. This finishes the proof, since such a splitting of the tangent space of P defines a connection. $\square$

On every Lorentz manifold there exists a distinguished connection, the Levi-Civita connection, with respect to which the metric is covariantly constant. In order to define the ‘pull-back’ of this connection to bundles associated to the tangential bundle, we need the notion of the so called ‘connection form’ Υ .

DEFINITION 2.11. Let $(P, M, \mathcal{G}, \mathbf{L})$ be a principal fiber bundle and let $\text{Lie-}\mathcal{G}$ denote the Lie algebra of $\mathcal{G}$, i.e., $\text{Lie-}\mathcal{G}$ is the Lie algebra of the left invariant vector fields over $\mathcal{G}$. For every $u \in P$ let σ_u denote the smooth mapping

$$\begin{aligned} \sigma_u: \mathcal{G} &\rightarrow P \\ g &\mapsto \sigma_u(g) = \mathbf{L}_g(u). \end{aligned}$$

A Lie algebra valued homomorphism λ from $\text{Lie-}\mathcal{G}$ into the Lie-algebra of the vertical³ vector fields over P is given by:

$$\begin{aligned} \lambda: \text{Lie-}\mathcal{G} &\rightarrow \Gamma(TP) \\ X &\mapsto \lambda X: u \mapsto (\sigma_u)_*(X_e), \end{aligned}$$

where X_e denotes the value of X at the identity of $\mathcal{G}$. Recall that $\text{Lie-}\mathcal{G}$ is isomorphic to the tangent space $T_e \mathcal{G}$ of $\mathcal{G}$ at the identity. Given a connection Ξ in $(P, M, \mathcal{G}, \mathbf{L})$, we define a 1-form Υ on P with values in $\text{Lie-}\mathcal{G}$ as follows. For every $t \in T_u P$ let $\Upsilon(t)$ to be the unique $X \in \text{Lie-}\mathcal{G}$, such that $\lambda X(u)$ is equal to the vertical component of t . The form Υ is called the *connection form* of the given connection Ξ .

³Note that the notion of a vertical vector field is independent of a connection.

PROPOSITION 2.12 (Proposition 1.1. of [KN63, p. 64]). *The Lie-algebra valued 1-form Υ is uniquely fixed by the following three requirements.*

1. $\Upsilon(\lambda X(u)) = X$ for all $u \in \mathbf{P}$ and $X \in \text{Lie-}\mathcal{G}$.
2. Υ is equivariant, i.e., $\Upsilon \circ d\mathbf{L}_g = \text{ad}(g^{-1}) \circ \Upsilon$ for all $g \in \mathcal{G}$, where ad denotes the adjoint representation of $\mathcal{G}$ in $\text{Lie-}\mathcal{G}$.
3. Υ is smooth.

For a given connection Ξ , Υ is given by Definition 2.11 above. For the reverse relation let Υ be a connection form. Then

$$Q_u = \{t \in T_u \mathbf{P} \mid \Upsilon(t) = 0\} \quad \text{and} \quad G_u = \{t \in T_u \mathbf{P} \mid t = \lambda \Upsilon(t)\}$$

define the horizontal and vertical subspace of $u \in \mathbf{P}$ respectively. Any connection in a principal fiber bundle defines canonically a connection in all associated fiber bundles: Let $(\mathbf{P}, M, \mathcal{G}, \mathbf{L})$ be a principal fiber bundle with connection Ξ , $\mathbf{B} = (\mathbf{P} \times \mathbf{F})/\mathcal{G}$ an associated fiber bundle with fiber $\mathbf{F}$. A parallel transport in $\mathbf{B}$ along some curve γ in M is defined by

DEFINITION 2.13. Let $\gamma: [0, 1] \rightarrow M$ denote a smooth curve in M , $\tilde{\pi}$ be the base point projection in $\mathbf{B}$ and $b \in \tilde{\pi}^{-1}(\gamma(0))$ be a point in $\mathbf{B}$. Moreover, let $f \in \mathbf{F}$ and $u \in \mathbf{P}$ be such that $\pi(u) = \gamma(0)$, $[u, f]_{\sim} = b$. By assumption there exists precisely one horizontal lift

$$\tilde{\gamma}: [0, 1] \rightarrow \mathbf{P} \quad \text{with} \quad \tilde{\gamma}(0) = u.$$

A lift of γ to $\mathbf{B}$ is defined by

$$\tilde{\gamma}(s) := [\tilde{\gamma}(s), f]_{\sim}. \quad s \in [0, 1]$$

We define a *parallel transport* $\bar{T}_\gamma$ along γ by

$$\bar{T}_\gamma = \hat{u} \circ T_\gamma(p) \circ \hat{u}^{-1},$$

where $\hat{u}$ denotes the mapping $\hat{u}: u \mapsto [u, f]_{\sim}$ from $\mathbf{P}$ to $\mathbf{B}$.

For the details the reader is referred to the book by S. Kobayashi and K. Nomizu loc.cit. We are now able to define the spinor connection:

DEFINITION 2.14. Let Ξ denote the connection form of the Levi-Civita connection in the Lorentz manifold (M, g_{ab}) with spin structure $(SM, \mathbf{p})$. The connection form σ of the spinor connection is the pull-back of Ξ to $(SM, \mathbf{p})$

$$\sigma := (\varpi)^{-1} \circ \mathbf{p}^*(\Xi),$$

where ϖ denotes the Lie-algebra homomorphism between Lie-Spin_0 and $\text{Lie-}\mathcal{L}_0^\dagger$.

EXAMPLE 2.15. Let (M, g_{ab}) a four dimensional orientable, time orientable Lorentz manifold with spinor bundle $(SM, \mathbf{p})$. Its Levi-Civita connection is characterized by the connection form Υ . Alternatively it can be regarded as a covariant derivative $\nabla: \Gamma(TM) \rightarrow \Gamma(T^*M \otimes TM)$: If $e \in \Gamma(TM)$ denotes a local orthonormal frame, i.e., for $\mu = 1, \dots, 4$ the four vector fields $e^{\mu a} \in \Gamma(TM)$ are everywhere linear independent and orthonormal, we have

$$\begin{aligned} \nabla_a e^{\mu b} &= \sum_\nu (\Upsilon^\mu{}_\nu)_a \otimes e^{\nu b} \\ &= \sum_{\lambda, \nu} \omega^{\lambda\mu\nu} e_a^\lambda \otimes e^{\nu b}, \end{aligned}$$

where $(\Upsilon^\mu{}_\nu)_a$ are the matrix elements of the $\text{Lie-}\mathcal{L}_0^\dagger$ valued one form $e^*(\Upsilon)$ on M . Recall that Υ is a $\text{Lie-}\mathcal{L}_0^\dagger$ valued one form on FM and an orthonormal frame e is a mapping $e: M \rightarrow FM$. Note that $\omega^{\lambda\mu\nu}$ is *anti-symmetric* in μ and ν . The relation between $(\Upsilon^\mu{}_\nu)_a$ and the usual Christoffel symbols is given by

$$\Gamma^\mu{}_{\nu\rho} = \Upsilon^\mu{}_\nu \circ de(e_\rho),$$

where e_ρ denotes the ρ ‘leg’ of the frame e . In terms of the partial derivatives of the metric the Christoffel symbols are written as

$$\Gamma^c{}_{ab} = \frac{1}{2} g^{cd} \{ \partial_a g_{bd} + \partial_b g_{ad} - \partial_d g_{ab} \}.$$

Now let γ_a be an arbitrary but fixed representation of the Dirac γ -matrices and let $E \in \Gamma(SM)$ be spin frame with $e = p \circ E$. Then one finds for the matrix elements of the Lie-Spin_0 valued one form $E^*(\sigma)$ on M

$$\sigma_a{}^A{}_B = -\frac{1}{4} \Gamma^b{}_{ac} \gamma^c{}_A \gamma^A{}_B$$

Combining these results, one finds that for any spinor tensor $f^{aA}_B \in \Gamma(TM \otimes DM \otimes D^*M)$ the covariant derivative has coordinate representation

$$\nabla_a f^{bA}_B = \partial_a f^{bA}_B + \Gamma^b_{ac} f^{cA}_B + \sigma_b^A{}_D f^{aD}_B - f^{aA}_D \sigma_b^D{}_B.$$

Further details and proofs can be found in the standard literature, for example [Wal84, BC64, MTW73, KN63].

Let us now investigate the conditions, which prevent the existence of spin structures for a given manifold. We remind the reader that these conditions are well known, but in order to formulate them, we have to deviate.

3. Čech-cohomology groups

Let $\mathbb{Z}_2$ denote the multiplicative group $\{+1, -1\}$. Let $U = (U_i)$ be an open covering of some manifold M .

DEFINITION 2.16. A mapping

$$f(i_0, \dots, i_r): \emptyset \neq U_{i_0} \cap U_{i_1} \cap \dots \cap U_{i_r} \rightarrow \mathbb{Z}_2$$

which is totally antisymmetric in the indices $(i_0, \dots, i_r)$ is called a *r-Čech-cochain*.

The multiplicative group of all r-cochains is denoted by $S^r(U, \mathbb{Z}_2)$. We define a *boundary operator* δ by

$$\delta: S^r(U, \mathbb{Z}_2) \rightarrow S^{r+1}(U, \mathbb{Z}_2)$$

$$f(i_0, \dots, i_r) \mapsto \prod_{j=0}^{r+1} f(i_0, \dots, \hat{i}_j, \dots, i_r),$$

where $\hat{}$ denotes the suppression of the corresponding index. It is easy to see that δ is nil potent⁴. The set of all cocycles $Z^r(U, \mathbb{Z}_2)$ and coboundaries $B^r(U, \mathbb{Z}_2)$ respectively are given by

$$Z^r(U, \mathbb{Z}_2) = \{f \in S^r(U, \mathbb{Z}_2) | \delta f = 1\}$$

and

$$B^r(U, \mathbb{Z}_2) = \{f \in S^r(U, \mathbb{Z}_2) | f = \delta g \text{ for some } g \in S^{r-1}(U, \mathbb{Z}_2)\}$$

The r-Čech-cohomology group $H^r(U, \mathbb{Z}_2)$ are the cocycles modulo the coboundaries.

$$H^r(U, \mathbb{Z}_2) = Z^r(U, \mathbb{Z}_2) / B^r(U, \mathbb{Z}_2) = \ker \delta_r / \text{Im} \delta_{r-1}$$

To achieve independence of the covering U we use an inductive limes process. The result

$$H^r(M, \mathbb{Z}_2) = \lim_{\substack{\longrightarrow \\ U}} \text{ind} H^r(U, \mathbb{Z}_2)$$

is called *r-Čech-cohomology group* of M (compare Appendix A).

4. Stiefel-Whitney classes

Stiefel-Whitney classes are characteristic classes with values in $H^r(M, \mathbb{Z}_2)$. In general these classes are defined for all fiber bundles with an orthogonal group as fiber group. We are interested in the special case of the tangent bundle of a four dimensional Lorentz manifold only, i.e., a fiber bundle with $O(1, 3)$ as fiber group. Let (M, g_{ab}) be as usual a four dimensional Lorentz manifold and $U = (U_i)$ a simple covering of M , that is a covering such that $U_i \cap U_j$ is either empty or simply connected for all pairs (i, j) . Let $\{e_{i\alpha}\}_{(\alpha=1,2,3,4)}$ denote a vierbein field over U_i . Let U_j denote another element of U then

$$e_{i\alpha} = t_{ij} e_{j\alpha},$$

where $t_{ij}: U_i \cap U_j \rightarrow O(3, 1)$ is the transition function for these two vierbein fields. The mapping

$$f: U_i \cap U_j \rightarrow \mathbb{Z}_2$$

$$t_{ij} \mapsto \det(t_{ij}) = \pm 1$$

defines a 1-cochain, since $f(i, j) = f(j, i)$. Moreover, we have

$$(\delta f)(i, j, k) = \det(t_{jk}) \det(t_{ik}) \det(t_{ij}) = \det(t_{jk} t_{ik} t_{ij}) = +1,$$

since for all transition functions t_{ij} in a fiber bundle the consistency condition

$$t_{ij}(p) t_{jk}(p) = t_{ik}(p) \tag{32}$$

must be satisfied. It follows that $f \in Z^1(M, \mathbb{Z}_2)$ is a cocycle defining an element

$$[f] \in H^1(M, \mathbb{Z}_2)$$

in the first cohomology group.

⁴Since $\mathbb{Z}_2$ is a multiplicative group, this means $\delta \delta f \equiv 1$.

LEMMA 2.17. $[f] \in H^1(M, \mathbb{Z}_2)$ is independent of the vierbein field $\{e_{i\alpha}\}$.

PROOF. Let $\{\tilde{e}_{i\alpha}\}$ denote another vierbein field over U_i with $\tilde{e}_{i\alpha} = h_i e_{i\alpha}$, where h_i is a mapping from U_i into $O(1, 3)$. Assume we are given the following transition functions $\tilde{t}_{ij}$, such that $\tilde{e}_{i\alpha} = \tilde{t}_{ij} \tilde{e}_{j\alpha}$ is satisfied. Between the original transition functions t_{ij} and $\tilde{t}_{ij}$ there is the following relation:

$$\tilde{t}_{ij} = h_i t_{ij} h_j^{-1}.$$

Therefore

$$\tilde{f}(i, j) := \det(h_i t_{ij} h_j^{-1}) = f(i, j) \cdot \delta(f_0)$$

where $f_0(i) := \det(h_i)$. Note that $\det(h_i) \equiv \det(h_i^{-1})$ since $f_0 \equiv \pm 1$. This shows that f and $\tilde{f}$ differ by a coboundary only, i.e., they define the same element $[f] \in H^1(M, \mathbb{Z}_2)$. $\square$

DEFINITION 2.18. The set of all elements $H^1(M, \mathbb{Z}_2) \ni [f] = w_1(M)$ is called *first Stiefel-Whitney class of M* .

LEMMA 2.19 ([Nak90]). M is orientable if and only if $w_1(M)$ is trivial.

PROOF. Assume M to be orientable. The structure group of the tangential bundle is reducible to $SO(1, 3)$ in this case and $\det(t_{ij}) \equiv +1$. This means that $w_1(M)$ is trivial. On the other hand assume the triviality of $w_1(M)$. This implies that $f(i, j)$ is a coboundary, i.e., $f(i, j) = (\delta f_0)(i, j)$. Since $f_0(i) = \pm 1$ for all i , we can find suitable elements $h_i: U_i \rightarrow O(1, 3)$, such that $f_0(i) = \det(h_i)$ for all i . Suppose we are given vierbein fields $\{e_{i\alpha}\}$ on U_i . Use the functions h_i to define new vierbein fields via $\tilde{e}_{i\alpha} = h_i e_{i\alpha}$. The corresponding transition functions $\tilde{t}_{ij}$ satisfy the relation $\det(\tilde{t}_{ij}) = +1$ for all pairs of indices (i, j) such that $U_i \cap U_j \neq \emptyset$:

Suppose we had $f(i, j) := \det(t_{ij}) = -1$ for some indices (i, j) . Choosing h_i and h_j such that $\det(h_i) = +1$ and $\det(h_j) = -1$ respectively, we find for the new functions $\tilde{f}(i, j)$:

$$\tilde{f}(i, j) = \det(h_i t_{ij} h_j^{-1}) = +1$$

$\square$

For the Definition of the second Stiefel-Whitney class consider a four dimensional *oriented* Lorentz manifold (M, g_{ab}) . Let TM denote its tangent bundle. Note that TM has $SO(1, 3)$ as fiber group. Let $U = \{U_i\}$ denote a simple covering of M and $t_{ij}: M \rightarrow SO(1, 3)$ are the local transition functions. We will denote the universal (twofold) covering group of $SO(1, 3)$ by Spin . Let $\phi: \text{Spin} \rightarrow SO(1, 3)$ be the corresponding 2:1 group homomorphism. A family $\{\tilde{t}_{ij}\}$, $\tilde{t}_{ij}: M \rightarrow \text{Spin}$ is called a *lift* of t_{ij} , if and only if for all pairs (i, j) such that $U_i \cap U_j \neq \emptyset$

1. $\phi(\tilde{t}_{ij}) = t_{ij}$
2. $\tilde{t}_{ji} = \tilde{t}_{ij}^{-1}$

are satisfied. Such families always exist locally. Since we have the consistency condition Eqn. (32) it follows that

$$\phi(\tilde{t}_{ij} \tilde{t}_{jk} \tilde{t}_{ki}) = \mathbf{1}$$

which implies

$$(\tilde{t}_{ij} \tilde{t}_{jk} \tilde{t}_{ki}) = \pm \mathbf{1}.$$

On the other hand the transition functions $\tilde{t}_{ij} \in \text{Spin}$ are defining a spin structure if and only if they fulfill the consistency condition

$$\tilde{t}_{ij} \tilde{t}_{jk} = \tilde{t}_{ik} \Leftrightarrow (\tilde{t}_{ij} \tilde{t}_{jk} \tilde{t}_{ki}) = +\mathbf{1} \quad (33)$$

We define a 2-Čech-cochain $f: U_i \cap U_j \cap U_k \rightarrow \mathbb{Z}_2$ by

$$(\tilde{t}_{ij} \tilde{t}_{jk} \tilde{t}_{ki}) = f(i, j, k) \cdot \mathbf{1}$$

By Eqn. (33) we have $f(i, j, k) \in S^2(M, \mathbb{Z}_2)$.⁵ We also find

$$\begin{aligned} (\delta f)(i, j, k, l) &= f(j, k, l) f(i, k, l) f(i, j, l) f(i, j, k) \\ &= (\pm 1)^{2 \cdot 6} t_{ij} t_{jk} t_{ki} t_{ij} t_{jl} t_{li} t_{ik} t_{kl} t_{li} t_{jk} t_{kl} t_{li} \\ &= +\mathbf{1}. \end{aligned}$$

⁵It is enough to show the symmetry: By Definition we have $f(i, j, k) = f(i, k, j)^{-1}$. It remains to show that $f(i, j, k) = f(j, k, i)$.

$$\begin{aligned} f(i, j, k) &= \tilde{t}_{ij} \tilde{t}_{jk} \tilde{t}_{ki} = \tilde{t}_{ij} \tilde{t}_{jk} \underbrace{\tilde{t}_{ki} \tilde{t}_{ij}}_{= f(j, k, i) \tilde{t}_{kj}} \tilde{t}_{jk} \tilde{t}_{ki} f(i, j, k) = f(i, j, k) f(j, k, i) f(i, j, k) \\ &= f(j, k, i) \tilde{t}_{kj} \end{aligned}$$

It follows that $f(i, j, k) = f(j, k, i)^{-1} = f(j, k, i)$.

Note that the second equality follows from the fact that we have a freedom of sign in every transition function and all transition function occur twice. Finally, using the consistency condition Eqn. (32) for the functions t_{ij} gives the result. It follows that f defines an element in $H^2(M, \mathbb{Z}_2)$. The set of all elements in $H^2(M, \mathbb{Z}_2) \ni [f] = w_2(M)$ is called 2. *Stiefel-Whitney class* of M . It is easy to show that $[f]$ is independent of the vierbein family $\{e_{i\alpha}\}$ and the ‘lift’.

5. An existence theorem

In this section we prove the following Theorem which ensures the existence of a spin structure for a given manifold. A particular property which prevents the existence of spin structures is described next.

THEOREM 2.20 ([Nak90]). *Let TM denote the tangent bundle of an orientable, time orientable, four dimensional Lorentz manifold M . There exists a spinor bundle over M precisely when the second Stiefel-Whitney class $w_2(TM)$ of TM is trivial.*

It is worth noting that this Theorem involves *global* properties of the manifold and is independent of the metric structure, i.e., the existence of a spin structure is a *topological* property of the manifold.

PROOF. Suppose there exists a spin structure over M , then all transition functions $\tilde{t}_{ij}$ satisfy the compatibility condition Eqn. (33) for all overlapping regions U_i, U_j, U_k . It follows that $w_2(M)$ is trivial. On the other hand assume the triviality of $w_2(M)$, i.e.,

$$f(i, j, k) = (\delta f_1)(i, j, k) = f_1(j, k)f_1(i, k)f_1(i, j),$$

where f_1 is a 1-cochain. Let $\tilde{t}_{ij}$ be some given transition functions. The transition functions

$$\tilde{t}'_{ij} = f_1(i, j)\tilde{t}_{ij}$$

satisfy

$$(\tilde{t}'_{ij}\tilde{t}'_{jk}\tilde{t}'_{ki}) = f_1(i, j)f_1(j, k)f_1(k, i) \cdot \tilde{t}_{ij}\tilde{t}_{jk}\tilde{t}_{ki} = \delta f_1 \delta f_1 = +1,$$

which ensures that $\{\tilde{t}'_{ij}\}$ defines a spin structure over M . □

We do now consider a ‘counterexample’: Let (M, g_{ab}) be a 4-dimensional orientable and time orientable Lorentz manifold. Consider a family $\{l_s\}_{s \in [0,1]}$ of closed curves in M . Suppose further that $l_0 \equiv l_1 \equiv p$, i.e., l is a special embedding of the two-sphere into M . Let $\{e_{i\alpha}\}$ denote an oriented, time oriented vierbein in the tangential space of the point p . The parallel transport of this vierbein along l yields a curve in the proper Lorentz group since both, the starting and the ending point of l coincide:

$$\begin{aligned} h_s &: [0, 1] \rightarrow \mathcal{L}_0^\uparrow \\ s &\mapsto h_s \end{aligned}$$

Moreover, since $h_0 \equiv h_1 \equiv 1$ this curve is closed.

LEMMA 2.21. *If h_1 is not homotop to the trivial curve then there exists no notion of a parallel transport for spinors, i.e., M does not admit a spin structure⁶.*

PROOF BY CONTRADICTION. Assume we have a spin structure and h_1 is *not* null homotop. Consider the parallel transport of some spinor along l_1 . On one hand l_1 is a constant curve, such that the spinor remains unchanged during its transport. On the other hand, since h_1 is homotop to a 2π rotation by assumption we conclude using the continuity of the parallel transport, that the spinor must change sign during the transport. Contradiction. □

Other properties of Lorentz manifolds *without* spin structure are investigated by Ohlmeyer [Ohl92], to which we refer the interested reader. To summarize the results of this section, the following statements are equivalent: Let (M, g_{ab}) denote a 4-dimensional orientable, time orientable Lorentz manifold, then

- I:** M admits a spin structure
- II:** The second Stiefel-Whitney class of M is trivial
- III:** Let $\mathcal{B}$ denote the bundle of all orthonormal bases of M . Its fundamental group can be written as $\pi_1(\mathcal{B}) = \pi_1(M) \times \mathbb{Z}_2$, where $\pi_1(M)$ denote the fundamental group of the manifold M , $\mathbb{Z}_2 = \{-1, +1\}$ and $\times$ denotes the direct product for groups.
- IV:** On every two sphere in M one can find three everywhere linearly independent vector fields.

If M is in addition non compact

- V:** M is parallelizable [Ger70]

⁶The parallel transport on the manifold induces –by a pull back of its connection form (see section 2)– a parallel transport in the spin structure.

Note that ‘**V**’ implies that all 4-dimensional globally hyperbolic manifolds admit a spin structure (See [Ger70]): As is pointed out by Geroch, an orientable or time orientable manifold does admit a spin structure precisely when every of its covering manifolds does. Without loss of generality we may therefore restrict ourselves to the case of an orientable, time orientable, globally hyperbolic manifold. These manifolds can be written as $M = \mathbb{R} \times \Sigma$, where Σ is a smooth spacelike hypersurface. Σ is three dimensional and by assumption orientable; hence parallelizable (see e.g. [Ste51, page 204]). Due to the fact that M is a Cartesian product of $\mathbb{R}$ and Σ this parallelizability can be extended to the whole of M . Using ‘**V**’ this implies the existence of a spin structure.

ON THE PROOF OF THE EQUIVALENCES.

Theorem 2.20.

I $\Leftrightarrow$ III $\Leftarrow$ III: Consider the following covering space $\tilde{B}$ of B . Let $p_0 \in B$ be fixed. Let p denote a curve in B from the point p_0 to $p \in B$. We introduce the following equivalence relation:

$$p \sim \tilde{p} \quad :\Leftrightarrow \quad \begin{array}{l} \text{i) } p \text{ and } \tilde{p} \text{ have the same end points} \\ \text{ii) } p \circ \tilde{p} \text{ belongs to the homotopy class } (g_1, 1) \in \pi_1(B) \text{ in } B, \\ \text{where } g_1 \in \pi_1(M) \text{ is a suitable element of the homotopy group of } M \end{array}$$

It follows that $\tilde{B}$ is a $SL(2, \mathbb{C})$ bundle over M . i.e., a spin structure.

I $\Rightarrow$ III: This implication follows immediately from the Definition of the spin structure: A spin structure is the two-fold covering of the bundle of all orthonormal bases. Thus any simply closed curve⁷ admits precisely two lifts –corresponding to the two elements in $\mathbb{Z}_2$ – in SM . Moreover, traveling along γ twice, yields lifts which are homotop.

I $\Leftrightarrow$ IV : We refer to the book of Penrose [PR84] for the proof of this equivalence⁸.

I $\Leftrightarrow$ V : This equivalence is shown in [Ger70]. We refer the reader to the original work for the proof. Note however that ‘**V** $\Rightarrow$ **I**’ is trivial and follows from the Definition of a spin structure or from: ‘**V** $\Rightarrow$ **IV** and **IV** $\Leftrightarrow$ **I**’.

□

⁷A curve γ is called *simple*, if and only if there exists no closed curve $\tilde{\gamma}$ such that $\tilde{\gamma} \circ \tilde{\gamma} = \gamma$.

⁸N.b. Actually Penrose shows: ‘**IV**’ implies that no situation as described in the ‘counterexample’ Lemma 2.21 above can occur.

An analogon of the free Wess-Zumino model

In this chapter we define an analogon of the free Wess-Zumino model. It was already mentioned above that supersymmetric models do not exist on a general spacetime. Thus we can only mimic the structure of supersymmetric models defined on Minkowski space. Since the Wess-Zumino model is the simplest of those, we mimic its structure, by considering a model which has the same field content. Namely one that consists of two bosonic and one fermionic (Majorana) field. We will refer to this kind of models as *locally supersymmetric* models. It is emphasized that this notion has no relation to supergravity, where local, i.e., spacetime dependent, supersymmetry transformations are considered. In our setting the word ‘local’ should remind of the fact that our model ‘looks’ like a supersymmetric model at every *point* of the spacetime. We begin with the description of the classical settings for the bosonic and fermionic parts, followed by a brief introduction to the algebraic approach to quantum field theory on curved spacetimes. In section 4 we review the global Hadamard condition of Kay and Wald and extend their definition to Fermi fields. In order to describe Hadamard states locally we introduce wave front sets in the next section. The chapter is closed with a proposal of a general wave front set spectrum condition, which might generalize the usual Minkowski spectrum condition to Lorentz manifolds and should be applicable even to models containing interaction. Let M be a four dimensional, smooth, second countable, paracompact, Hausdorff manifold equipped with a metric field g_{ab} with signature $(+, -, -, -)$. The pair (M, g_{ab}) is called a (*curved*) *spacetime*. In the following we will use the timelike sign convention of Landau and Lifshitz which differs from the Misner, Thorne and Wheeler convention ([MTW73]) in the sign of the metric. In particular we define the Riemann tensor by

$$(\nabla_a \nabla_b - \nabla_b \nabla_a) \omega_c =: R_{abc}{}^d \omega_d \quad \forall \text{ dual vector fields } \omega_c$$

and the Ricci tensor by $R_{ab} := R_{acb}{}^c$. Let us assume further that (M, g_{ab}) is orientable, time orientable and globally hyperbolic. Recall that a manifold is called globally hyperbolic if and only if it is of the form $M = \Sigma \times \mathbb{R}$, where Σ is a (spacelike) Cauchy surface. The last assumption guarantees the existence of a spin-structure, which is necessary for the definition of spinor fields (see [Ger68] or Theorem 2.20 above).

REMARK . It should be noted that at least for the metric both sign conventions are used in the literature. This makes the comparison of results sometimes quite difficult, especially if the authors do not explicitly state their conventions. For a recipe to change the formulas between different conventions, the reader is referred to the book of Birrell and Davis [BD82].

Spin 0: A classical neutral massive scalar field $\Phi \in C^\infty(M)$ propagating in a curved spacetime (M, g_{ab}) is described by the action functional

$$S[\Phi] = +\frac{1}{2} \int (\Phi_{;a} \Phi^{;a} - (m^2 + \xi R) \Phi^2) d\mu, \quad (34)$$

where $d\mu$ is the invariant volume measure, $m \geq 0$ is the mass, ξ describes the coupling ($\xi = 0$ minimal, $\xi = \frac{1}{6}$ conformal and $\xi = \frac{1}{4}$ SUSY) and $R \equiv R_a{}^a$ is the scalar curvature. As usual a semicolon denotes covariant differentiation. The field equation is the Klein Gordon equation

$$0 = \frac{\delta S}{\delta \Phi} = (\square + (m^2 + \xi R)) \Phi(x),$$

where $\square = \nabla_\mu \nabla^\mu$ is the D’Alembert operator. The canonical energy momentum tensor is

$$\begin{aligned} \Theta^{ab} &= \frac{\delta \mathcal{L}}{\delta(\frac{\partial \Phi_i}{\partial x^{(a)}})} \frac{\partial \Phi_i}{\partial x^{(b)}} - g^{ab} \mathcal{L} \\ &= \Phi^{;(a} \Phi^{;b)} - \frac{1}{2} g^{ab} (\Phi_{;c} \Phi^{;c} - (m^2 + \xi R) \Phi^2), \end{aligned}$$

where $\mathcal{L}$ denotes the Lagrangian density of Φ and (ab) means symmetrization. However, as potential source term for the Einstein equation the improved energy momentum tensor T^{ab} introduced by Callan et al. [CCJ70] seems more appropriate. This improved energy momentum tensor is obtained by functionally differentiating the action with respect to the metric field. This treats the metric as a dynamical variable which seems natural in view of general relativity. (See the

discussion in the paper by DeWitt and Brehme [DB60] or Fulling's book [Ful89] for further details.) We find that in the SUSY coupled case

$$\begin{aligned} T^{ab} &:= 2g^{-1/2} \frac{\delta S}{\delta g_{ab}} \\ &\equiv \left\{ \frac{1}{4} \{ \Phi^{;a}, \Phi^{;b} \}_+ - \frac{1}{4} \{ \Phi^{;ab}, \Phi \}_+ + \frac{1}{16} g^{ab} \{ \Phi_{;c}^c, \Phi \}_+ - \frac{1}{8} (R^{ab} + \frac{1}{8} R g^{ab}) \{ \Phi, \Phi \}_+ \right. \\ &\quad \left. + \frac{1}{16} m^2 g^{ab} \{ \Phi, \Phi \}_+ \right\} \end{aligned} \quad (35)$$

where $\{ \cdot, \cdot \}_+$ denotes the anticommutator and $g^{1/2}$ is the square root of the determinant of the metric. We now do point splitting: Write each bracket in Eqn. (35) in the form

$$\begin{aligned} \{ \Phi^{;a}, \Phi^{;b} \}_+ &\rightarrow \lim_{x' \rightarrow x} \frac{1}{2} \left(g^a_{c'} \{ \Phi, \Phi' \}_+^{;c'b} + g^b_{c'} \{ \Phi, \Phi' \}_+^{;ac'} \right) \\ \{ \Phi^{;ab}, \Phi \}_+ &\rightarrow \lim_{x' \rightarrow x} \frac{1}{2} \left(\{ \Phi, \Phi' \}_+^{;ab} + g^a_{c'} g^b_{d'} \{ \Phi, \Phi' \}_+^{;c'd'} \right) \\ \{ \Phi, \Phi \}_+ &\rightarrow \lim_{x' \rightarrow x} (\{ \Phi, \Phi' \}_+) \end{aligned}$$

where x' is a point near x , $\Phi' = \Phi(x')$ and $\{ \cdot, \cdot \}_+^{;a'}$ represents the covariant derivative at x' . $g^a_{c'}$ is the bi-tensor of parallel transport defined by the equation $\sigma_{;a} g^a_{c'} = 0$ together with the boundary condition $g^a_c(x, x) = \delta^a_c(x, x)$. The bi-scalar $\sigma(x, x')$ is half the square of the geodesic distance between x and x' . Recall that σ is well defined whenever x and x' are sufficiently close to each other. We end up with

$$\begin{aligned} T^{ab} &= \lim_{x' \rightarrow x} \left\{ \frac{1}{8} \left(g^a_{a'} \{ \Phi, \Phi' \}_+^{;a'b} + g^b_{b'} \{ \Phi, \Phi' \}_+^{;ab'} \right) - \frac{1}{8} \left(\{ \Phi, \Phi' \}_+^{;ab} + g^a_{a'} g^b_{b'} \{ \Phi, \Phi' \}_+^{;a'b'} \right) \right. \\ &\quad \left. + \frac{1}{32} g^{ab} \left(\{ \Phi, \Phi' \}_+^{;c} + g_{cd} g^c_{c'} g^d_{d'} \{ \Phi, \Phi' \}_+^{;c'd'} \right) \right. \\ &\quad \left. - \frac{1}{8} (R^{ab} + \frac{1}{8} R g^{ab}) \{ \Phi, \Phi' \}_+ + \frac{1}{16} m^2 g^{ab} \{ \Phi, \Phi' \}_+ \right\} \end{aligned} \quad (36)$$

Spin 1/2: A classical (anti-commuting!) massive Majorana spinor field Ψ propagating in a spacetime (M, g_{ab}) with spin structure (SM, p) has the action functional (see [DeW65, BD82])

$$S[\Psi] = \frac{1}{2} \int \Psi^+ (i \not{\nabla} \Psi - m \Psi) d\mu.$$

Note the factor 1/2 which is necessary since we assume a Majorana spinor field. The field equation is the Dirac equation

$$(i \not{\nabla} - m) \Psi = 0$$

and for the improved energy momentum tensor one finds ([BD82])

$$T^{ab} = -\frac{i}{8} \left(\Psi^{+;a} \gamma^b \Psi + \Psi^{+;b} \gamma^a \Psi - \Psi^+ \gamma^a \Psi^{;b} - \Psi^+ \gamma^b \Psi^{;a} \right)$$

Since Ψ is a Majorana spinor, we use the identities of Appendix C and write:

$$T^{ab} = \frac{i}{8} \sum_{A,B} \gamma^{(aA}{}_B \left([\Psi^B, \Psi^{+A}]_-^{;b)} - [\Psi^{;b)B}, \Psi^+{}_A]_- \right)$$

where $()$ means symmetrization, $[\cdot, \cdot]_-$ is the commutator and a local spin frame E_A was chosen. Note that T^{ab} transforms as a second rank tensor under changes of frames. Since it is already symmetric we do point splitting by writing

$$T^{ab} = \lim_{x' \rightarrow x} \frac{i}{8} \sum_{A,B} \gamma^{(aA}{}_B \left(g^b_{b'} \mathcal{J}_A^{A'} [\Psi^B, \Psi'^+{}_{A'}]_-^{;b')} - \mathcal{J}_A^{A'} [\Psi^B, \Psi'^+{}_{A'}]_-^{;b)} \right)$$

Here $\mathcal{J}_A^{A'}$ is the bi-spinor of parallel transport which is defined by the equation $\sigma^a \mathcal{J}_A^{B'}{}_{;a} = 0$ together with the boundary condition $\mathcal{J}_A^B(x, x) = \mathbf{1}_A^B(x, x)$. *Local supersymmetry:* A classical locally supersymmetric field theoretical model is a

free field model with two SUSY coupled neutral scalar fields (A, B) together with one (anti-commuting) Majorana spinor

field (Ψ). The action functional of this model is

$$S[A, B, \Psi] = \frac{1}{2} \int (A_{;a} A^{;a} - (m^2 - \xi R) A^2 + B_{;a} B^{;a} - (m^2 - \xi R) B^2 + \Psi^+ (i \overleftarrow{\nabla} - m) \Psi) d\mu.$$

The equation of motion for the three fields obviously decouples, giving the Klein Gordon equation for both A and B and the Dirac equation for Ψ . The already point-separated improved energy momentum tensor is

$$\begin{aligned} T^{ab} = & \lim_{x' \rightarrow x} \left\{ \frac{1}{8} \left(g^a_{a'} \{A, A'\}_+^{;a'b} + g^b_{b'} \{A, A'\}_+^{;ab'} \right) - \frac{1}{8} \left(\{A, A'\}_+^{;ab} + g^a_{a'} g^b_{b'} \{A, A'\}_+^{;a'b'} \right) \right. \\ & + \frac{1}{32} g^{ab} \left(\{A, A'\}_+^{;c} + g_{cd} g^c_{c'} g^d_{d'} \{A, A'\}_+^{;c'd'} \right) \\ & - \frac{1}{8} (R^{ab} + \frac{1}{8} R g^{ab}) \{A, A'\}_+ + \frac{1}{16} m^2 g^{ab} \{A, A'\}_+ + \frac{1}{8} \left(g^a_{a'} \{B, B'\}_+^{;a'b} + g^b_{b'} \{B, B'\}_+^{;ab'} \right) \\ & - \frac{1}{8} \left(\{B, B'\}_+^{;ab} + g^a_{a'} g^b_{b'} \{B, B'\}_+^{;a'b'} \right) + \frac{1}{32} g^{ab} \left(\{B, B'\}_+^{;c} + g_{cd} g^c_{c'} g^d_{d'} \{B, B'\}_+^{;c'd'} \right) \\ & - \frac{1}{8} (R^{ab} + \frac{1}{8} R g^{ab}) \{B, B'\}_+ + \frac{1}{16} m^2 g^{ab} \{B, B'\}_+ \Big\} \\ & + \frac{i}{8} \sum_{A,B} \gamma^{(aA}{}_B \left(g^b_{b'} \mathcal{J}_A{}^{A'} [\Psi^B, \Psi'^+_{A'}]_-^{;b'}) - \mathcal{J}_A{}^{A'} [\Psi^B, \Psi'^+_{A'}]_-^{;b}) \right) \end{aligned} \quad (37)$$

On a Ricci flat spacetime ($R^{ab} \equiv 0$) we obtain,

$$\begin{aligned} T^{ab} = & \lim_{x' \rightarrow x} \left\{ \frac{1}{8} \left(g^a_{c'} \{A, A'\}_+^{;c'b} + g^b_{c'} \{A, A'\}_+^{;ac'} \right) - \frac{1}{8} \left(\{A, A'\}_+^{;ab} + g^a_{c'} g^b_{d'} \{A, A'\}_+^{;c'd'} \right) \right. \\ & + \frac{1}{8} \left(g^a_{c'} \{B, B'\}_+^{;c'b} + g^b_{c'} \{B, B'\}_+^{;ac'} \right) - \frac{1}{8} \left(\{B, B'\}_+^{;ab} + g^a_{c'} g^b_{d'} \{B, B'\}_+^{;c'd'} \right) \\ & + \frac{i}{8} \sum_{A,B} \gamma^{(aA}{}_B \left(g^b_{b'} \mathcal{J}_A{}^{A'} [\Psi^B, \Psi'^+_{A'}]_-^{;b'}) - \mathcal{J}_A{}^{A'} [\Psi^B, \Psi'^+_{A'}]_-^{;b}) \right) \Big\} \quad \text{iff } R^{ab} \equiv 0. \end{aligned}$$

Note that we used the field equations for A and B .

REMARK . For classical fields Eqn. (37) is well defined. However in the quantized version of this model the basic fields are operator valued distributions on some GNS Hilbertspace (see section 3 this chapter below). In this case Eqn. (37) contains –in the limit– the product of the fields in the sense of a product of distributions. As is well known from Minkowski space such products of operator valued distributions do not make sense in general. Thus at this stage Eqn. (37) is only formal for the quantized model. On the other hand Christensen describes in [Chr76, Chr78] a method for renormalizing the expectation values of T^{ab} given by Eqn. (37) for a special class of states, namely for Hadamard states. We will use his results in chapter 4 to show that Eqn. (38) without renormalization remains well defined after quantization.

1. A locally conserved ‘supercurrent’

It is amusing to observe that on a special class of curved spacetimes a locally supersymmetric model possesses due its Bose-Fermi symmetry a locally conserved ‘supercurrent’. In contrast to the Minkowski space, however, this current can *not* be obtained by a Noether method, since on generic spacetimes there exist no covariant constant (i.e. parallel) spinor fields. We remark that there is a tight connection between such parallel spinors and the notion of a Kähler manifold, for which we refer the interested reader to chapter 10.8 in [BT87]. To obtain this current we consider the ‘variable’ part of Eqn. (18).

PROPOSITION 3.1. *The current*

$$k^{aA} = i \left((A - i\gamma^5 B)(i \overleftarrow{\nabla} - m) \gamma^a \Psi \right)^A \quad (38)$$

is locally conserved on spacetimes with vanishing scalar curvature ($R = 0$).

PROOF. By straightforward calculation

$$\begin{aligned} \nabla_a k^{aA} = & - \left((A - i\gamma^5 B)(\overleftarrow{\nabla} \overleftarrow{\nabla} + m^2) \Psi \right)^A - i \left((A - i\gamma^5 B) \overleftarrow{\nabla} (i \overleftarrow{\nabla} - m) \Psi \right)^A \\ = & 0 \quad \text{iff } R = 0. \end{aligned}$$

We used the field equations and the fact that $\nabla \nabla$ applied to a scalar is the D'Alembert operator. Note that the assumption on the scalar curvature of the spacetime enters via the scalar field equations. $\square$

2. The improved 'supercurrent'

In chapter 1 we saw that the (canonical) energy momentum tensor of the free Wess Zumino model can be expressed in terms of the supercurrent k^a (Eqn. (30)). To obtain the *improved* energy momentum tensor in this way it is necessary to use the following 'improved' supercurrent:

$$j^{aA} = k^{aA} - 2\xi(\gamma^a\gamma^b - \gamma^b\gamma^a)^A{}_B \nabla_b ((A - i\gamma^5 B)\Psi)^B \quad (39)$$

REMARK . Ferrara and Zumino introduced the 'improved' supercurrent in 1975 [FZ75] to simplify their studies of the transformation properties of various supercurrents in Minkowski space. They show that the 'improved' supercurrent can be used to construct all generators of the supersymmetry transformations and emphasize that this step is perfectly analogous to the introduction of the 'improved' energy momentum tensor [CCJ70] from which the generators of the Minkowski space symmetries can be constructed.

It is obvious that $\nabla_a j^{aA} = 0$ on Minkowski space since $\nabla_a \nabla_b$ is symmetric in this case. Moreover, since j^{aA} differs from k^{aA} by a pure divergence only, it follows that both of these currents generate the same charge on Minkowski space. On the other hand on a generic spacetime the improved 'supercurrent' is *not* covariantly conserved. Only on spacetimes with vanishing scalar curvature we find in analogy to Proposition 3.1:

PROPOSITION 3.2. *Let (M, g_{ab}) denote a Lorentz manifold with spinor structure. The current*

$$j^{aA} = k^{aA} - 2\xi(\gamma^a\gamma^b - \gamma^b\gamma^a)^A{}_B \nabla_b ((A - i\gamma^5 B)\Psi)^B$$

is covariantly constant if and only if (M, g_{ab}) has vanishing scalar curvature, i.e., $R \equiv 0$.

PROOF. Let us abbreviate $(\gamma^a\gamma^b - \gamma^b\gamma^a)$ by $4G^{[ab]}$, where the square brackets reminds of the fact that $G^{[ab]}$ is antisymmetric in a and b . We have the following useful identity

$$G^{[ab]}G_{[cd]}R^{cd}{}_{ab} = +\frac{1}{8}R,$$

which is valid since

$$G^{[ab]}G_{[cd]}R^{cd}{}_{ab} = \frac{1}{4}\gamma^a\gamma^b\gamma^c\gamma^d R_{abcd},$$

since $R_{abcd} = R_{[ab][cd]}$

$$= \frac{1}{4}\gamma^{(a}\gamma^b\gamma^c)\gamma^d R_{cdab},$$

since $R_{[abc]d} = 0$ by the Bianchi identities

$$\begin{aligned} &= \frac{1}{24} \left(g^{ab}\gamma^c + g^{bc}\gamma^a + \frac{\gamma^a\gamma^b\gamma^c + \gamma^c\gamma^b\gamma^a}{2} \right) \gamma^d R_{cdab} \\ &= \frac{1}{24} (2g^{bc}\gamma^a - g^{ac}\gamma^b) \gamma^d R_{cdab}, \end{aligned}$$

since R_{abcd} is antisymmetric in ab

$$\begin{aligned} &= -\frac{1}{8}\gamma^a\gamma^d R_{bad}{}^b \\ &= +\frac{1}{8}R, \end{aligned}$$

since $R_{ab} = R^c{}_{acb}$ is symmetric in ab

We obtain

$$\begin{aligned}\nabla_a j^{aA} &= \underbrace{\nabla_a k^{aA}}_{=0} - 8\xi G^{[ab]} \nabla_a \nabla_b ((A - i\gamma^5 B)\Psi) \\ &= -8\xi G^{[ab]} \left((\nabla_a \nabla_b (A - i\gamma^5 B))\Psi + [2(\nabla_a (A - i\gamma^5 B))(\nabla_b \Psi)] + (A - i\gamma^5 B)\nabla_a \nabla_b \Psi \right),\end{aligned}$$

where round brackets denote symmetrization

$$\begin{aligned}&= -2\xi (A - i\gamma^5 B) G^{[ab]} G_{[cd]} R^{cd}{}_{ab} \Psi \\ &= -\frac{1}{4} \xi (A - i\gamma^5 B) R \Psi\end{aligned}$$

Here we used the fact that the commutator of two covariant derivations on scalars vanishes whereas on spinors it is given by

$$[\nabla_a, \nabla_b] \Psi = (\nabla_a \nabla_b - \nabla_b \nabla_a) \Psi = 2\nabla_{[a} \nabla_{b]} \Psi = \frac{1}{2} G_{[cd]} R^{cd}{}_{ab} \Psi.$$

It follows that the expression Eqn. (40) vanishes precisely when $R = 0$, i.e., if and only if the spacetime has vanishing scalar curvature. $\square$

REMARK . In particular the improved ‘supercurrent’ is conserved on all spacetimes which are *vacuum solutions* of the Einstein field equations, provided these solutions admit the existence of spinors.

3. Quantization: The algebraic approach to quantum field theory

This section is written for the convenience of the reader and gives a brief introduction into the terminology and notions of the algebraic approach to quantum field theory in curved spacetimes. For further information the reader is referred to the books of Haag [Haa92], Baumgärtel & Wollenberg [BW92] or Horuzhi [Hor90]. Recall that in quantum physics a “basic field” Φ is an operator valued distribution on some Hilbertspace $\mathcal{H}$ satisfying certain conditions which will not be recalled here (see Haag’s book [Haa92] for details and references). These fields can be used to associate to each open region $\mathcal{O}$ of the spacetime an algebra $\mathfrak{A}(\mathcal{O})$ of (unbounded) operators on $\mathcal{H}$, namely the algebra generated by $\Phi(f)$ –the basic field “smeared out” with test functions f with support in $\mathcal{O}$. The elements of $\mathfrak{A}(\mathcal{O})$ can be interpreted as representing physical operations performable in $\mathcal{O}$. Furthermore, once the correspondence

$$\mathcal{O} \rightarrow \mathfrak{A}(\mathcal{O}) \quad (40)$$

is known, one can (in principle) compute all relevant physical quantities, e.g., cross-sections etc. This suggests that all relevant physical information is encoded in the relation (40). For mathematical convenience one restricts to local algebras generated by bounded operators, i.e., to algebras generated by $e^{i\Phi(f)}$ in the notation used above. The corresponding net is called *the net of local algebras* if and only if a generalization of the Haag Kastler axioms are fulfilled (see [Dim80]). However for the purpose of this text it is more convenient to consider a larger net of algebras, namely the net of Borchers Uhlmann algebras. This net, first mentioned by Borchers and Uhlmann ([Bor62, Uhl62]), contains kinematic information only, meaning for example that our model consists of two scalar and one Majorana spinor field. We will denote the net of Borchers Uhlmann algebras by $\mathfrak{A}_U(\mathcal{O})$. A remark concerning the relation between these nets is given at the end of this section. For a scalar field and a given region $\mathcal{O} \subset M$, $\mathfrak{A}_U(\mathcal{O})$ is the tensor algebra of smooth functions with support in $\mathcal{O}$. To be more precise, let $D_m(\mathcal{O})$ denote $\bigotimes^m C_0^\infty(\mathcal{O})$, $m \geq 1$, equipped with the usual topology and define $D_0(\mathcal{O}) \equiv \mathbb{C}$. $D'_m(\mathcal{O})$ denotes $\bigotimes^m [D'(\mathcal{O})]$, the dual space of $D_m(\mathcal{O})$. An element $\lambda^m \in D'_m(\mathcal{O})$ is called a *m-point distribution*. For a family of functions $\{f_m\}_{m \in \mathbb{N}_0}$, where $f_m \in D_m(\mathcal{O})$ with only finitely many f_m non-vanishing, define $f := \bigoplus^m f_m$. Define an involution as $f^* := \bigoplus^m f_m^*$, where $f_m^*(x_1, \dots, x_m) := \bar{f}_m(x_m, \dots, x_1)$, a non-commutative product for f and $g = \bigoplus^m g_m$ by $f \times g = \bigoplus^m (f \times g)_m$, where $(f \times g)_m := \sum_{i=0}^m f_i(x_1, \dots, x_i) g_{m-i}(x_{i+1}, \dots, x_m)$. Together with grade-wise addition the set of such functions f becomes the involutive algebra $\mathfrak{A}_U(\mathcal{O})$. We equip $\mathfrak{A}_U(\mathcal{O})$ with the direct sum topology coming from $D_m(\mathcal{O})$. The dual space of this algebra –denoted by $\mathfrak{A}'(\mathcal{O})$ – consists of a hierarchy of m-point distributions:

$$\mathfrak{A}'(\mathcal{O}) \ni \lambda = (\lambda^1, \dots, \lambda^m, \dots)$$

with $\lambda(f) = \sum_{m=0}^\infty \lambda^m(f_m)$. A state ω on the Borchers Uhlmann algebra $\mathfrak{A}_U(\mathcal{O})$ is a normalized linear form on $\mathfrak{A}_U = \bigcup \mathfrak{A}_U(\mathcal{O})$, i.e.,

$$\begin{aligned}\omega(f^* f) &\geq 0 & \forall f \in \mathfrak{A}_U \\ \omega(\mathbf{1}) &\equiv \omega_0(1) = 1\end{aligned}$$

Such a state is said to satisfy *local commutativity* if and only if for all $m \geq 2$ and all $1 \leq i \leq m-1$ we have

$$\omega^m(f_1 \otimes \cdots \otimes f_i \otimes f_{i+1} \otimes \cdots \otimes f_m) = \omega^m(f_1 \otimes \cdots \otimes f_{i+1} \otimes f_i \otimes \cdots \otimes f_m)$$

for all $f_1, \dots, f_m \in C_0^\infty(\mathcal{O})$ such that $\text{supp } f_i$ and $\text{supp } f_{i+1}$ are spacelike separated. The state is called *quasifree* if and only if its one point distribution vanishes and all higher n -point distributions ω^n are completely determined by its two point distribution ω^2 . The relation between “quantum fields” mentioned at the beginning of this section and the algebraic approach via the Borchers Uhlmann algebra is the GNS-reconstruction theorem, Theorem 3.4 below for which we need the following definition (Wightman axioms).

DEFINITION 3.3 (See, e.g. [Rad92]). A quadruple $(\mathcal{H}, D, \Phi, \Omega)$ is called a neutral scalar quantum field model on a spacetime (M, g_{ab}) , if the following properties are satisfied:

1. $\mathcal{H}$ is a separable Hilbertspace with positive definite inner product $\langle \cdot, \cdot \rangle$.
2. D is a dense subspace of $\mathcal{H}$.
3. Φ is an $\mathcal{H}$ operator valued distribution on $C_0^\infty(M)$.
4. For each $f \in C_0^\infty(M)$ the domain of $\Phi(f)$ contains D , and

$$\Phi(f)D \subset D$$

5. The vector $\Omega \in \mathcal{H}$ is cyclic for Φ and D is the linear span of the set

$$\{\Phi(f_1) \cdots \Phi(f_m)\Omega \in \mathcal{H} | f_1, \dots, f_m \in C_0^\infty(M)\}$$

6. Hermiticity of the field. For each $f \in C_0^\infty(M)$, the domain of the adjoint of $\Phi(f)$ (denoted by $\Phi(f)^*$) contains D and

$$\Phi(f)^* = \Phi(\bar{f}) \quad \text{on } D,$$

where $\bar{}$ means complex conjugation.

7. Local commutativity. For two test-functions $f_1, f_2 \in C_0^\infty(M)$ with spacelike separated support, the commutator $[\Phi(f_1), \Phi(f_2)]_-$ vanishes on D .

THEOREM 3.4 (GNS-Reconstruction). *For every state ω on the Borchers Uhlmann algebra $\mathfrak{A}_U(\mathcal{O})$ satisfying local commutativity there is a scalar quantum field theoretical model $(\mathcal{H}, D, \Phi, \Omega)$, unique up to unitary equivalence, such that for $f_1, \dots, f_m \in C_0^\infty(\mathcal{O})$ we have*

$$\omega^m(f_1 \otimes \cdots \otimes f_m) = \langle \Omega, \Phi(f_1), \dots, \Phi(f_m)\Omega \rangle. \quad (41)$$

Conversely by Eqn. (41) every scalar quantum field theoretical model $(\mathcal{H}, D, \Phi, \Omega)$ defines a state ω on $\mathfrak{A}_U(\mathcal{O})$ satisfying local commutativity.

For a proof of this Theorem see the standard literature (for example [BW92, B⁺90]).

So far we have only considered models with spin zero. For models with higher spin the Borchers Uhlmann algebra has to be modified slightly. For the spin 1/2 case, i.e., fermions, the corresponding definitions are given in the next paragraph. In analogy to the scalar case we define the Borchers Uhlmann algebra for Dirac spinor fields to be the tensor algebra generated by sections with compact support in the spinor and cospinor bundle respectively. To be more precise, let $\hat{D}_0(\mathcal{O}) \equiv \mathbb{C}$ for all open subsets $\mathcal{O} \subset M$ with compact closure. Denote by $\hat{D}_1(\mathcal{O}) = \Gamma(\mathcal{DM}(\mathcal{O}))$ the vector space of all sections in $\mathcal{DM} := \mathcal{DM} \oplus \mathcal{D}^*M$ with support in $\mathcal{O}$ equipped with the topology induced by $D_1(\mathcal{O})$. $\hat{D}_m(\mathcal{O})$ is the m -fold tensor product of the spaces $\hat{D}_1(\mathcal{O})$ ¹ Note that this product is neither commutative nor anti-commutative.

REMARK . Let $\hat{D}'_1(\mathcal{O})$ denote the dual space of $\hat{D}_1(\mathcal{O})$. We sometimes denote by $(f \otimes g)(x, x')$ the integral kernel of the map $f \otimes g : \hat{D}'_1(\mathcal{O}) \times \hat{D}'_1(\mathcal{O}) \mapsto \mathbb{C}$ defined by

$$(f \otimes g)(f^*, g^*) \equiv \int_M \int_M (f \otimes g)(x, x') f^*(x) g^*(x') d\mu d\mu'$$

The transformation properties of the components of $(f \otimes g)(x, x')$ under a change of frames show that $(f \otimes g)$ is a *bi-spinor*. If $f, \hat{f} \in \Gamma(\mathcal{DM}) \xrightarrow{i} \hat{D}_1(\mathcal{O})$ are two spinors with compact support we abbreviate $(f \otimes \hat{f})(x, x')$ by $f^A \hat{f}^{A'}$. The prime on the second index emphasizes the bi-spinor character.

We consider now a family of functions $\{f_m\}_{m \in \mathbb{N}_0}$, where $f_m \in \hat{D}_m(\mathcal{O})$ and with only finite many f_m non-vanishing and define $f := \bigoplus^m f_m$. For those elements f , such that all f_m can be written in the form $f_m = g_1 \otimes \cdots \otimes g_m$ an involution is defined by $f^* := \bigoplus^m f_m^*$ where $f_m^* \equiv (g_1 \otimes \cdots \otimes g_m)^* := g_m^+ \cdots \otimes g_1^+$. $(\cdot)^+$ means Dirac conjugation. For the other elements we define the involution by continuous extension. A product of f and $g = \bigoplus^m g_m$ is defined as

¹Recall that the tensor product of two vector bundles with typical fiber E over some base manifold M is the vector bundle over M with typical fiber $E \otimes E$. Its Fréchet topology is defined analogously to the one of $D_1(\mathcal{O})$, i.e., by the usual family of semi-norms (See, e.g. [Die72] 17.2).

$f \times g = \bigoplus^m (f \times g)_m$ where $(f \times g)_m = \sum_{n=1}^m f_n \otimes g_{m-n}$. With grade-wise addition the set of such f 's becomes the involutive algebra $\hat{\mathfrak{A}}_U(\mathcal{O})$, the Borchers algebra for Dirac Spinors. The topology on $\hat{\mathfrak{A}}_U(\mathcal{O})$ is the one inherited from $\hat{D}_m(\mathcal{O})$. We do not give the definition of the dual space and of states since it is identical to the scalar case (see above).

DEFINITION 3.5. A Dirac quantum field theoretical model is a quadruple $(\mathcal{H}, D, \Psi, \Psi^+, \Omega)$ such that the properties 1-5 of Definition 3.3 –replacing Φ by Ψ and $C_0^\infty(M)$ by $\Gamma(\mathcal{DM})$ – are fulfilled. The remaining two properties are substituted by

1. For each $f \in \Gamma(\mathcal{DM})$, the domain of the adjoint of $\Psi(f)$ –denoted by $\Psi(f)^*$ – and of the Dirac adjoint –denoted by $\Psi^+(f)$ both contain D and

$$\Psi(f)^* = \Psi^+(f^+) \quad \text{on } D,$$

where f^+ denotes the Dirac conjugate of f .

2. Local anti-commutativity. For two test-functions $f_1, f_2 \in \Gamma(\mathcal{DM})$ with spacelike separated support, the anticommutator $[\Psi(f_1), \Psi^+(f_2)]_+$ vanishes on D .

The GNS-reconstruction theorem for Dirac fields is

THEOREM 3.6 (GNS-reconstruction). *For every state ω on the Borchers Uhlmann algebra $\hat{\mathfrak{A}}_U(\mathcal{O})$ for the Dirac field satisfying local anti-commutativity there is a Dirac quantum field theoretical model $(\mathcal{H}, D, \Psi, \Psi^+, \Omega)$, unique up to unitary equivalence, such that for $f_1, \dots, f_m \in \Gamma(\mathcal{DM})$ we have*

$$\omega^m(f_1 \otimes \dots \otimes f_m) = \langle \Omega, \Psi(f_1), \dots, \Psi(f_m) \Omega \rangle. \quad (42)$$

Conversely by Eqn. (42) every Dirac quantum field theoretical model $(\mathcal{H}, D, \Psi, \Psi^+, \Omega)$ defines a state ω on $\hat{\mathfrak{A}}_U(\mathcal{O})$ satisfying local anti-commutativity.

REMARK . $\Psi(f_1) = \Psi \left(\begin{pmatrix} g \\ h^+ \end{pmatrix} \right)$ is identical to $\Psi(g) + \Psi^+(h^+)$ in usual notation.

We get the Borchers Uhlmann algebra for Majorana Spinors by restricting ourself to the Majorana subbundle of $\mathcal{DM}$ and repeating the construction above. To clarify the relation between the net of local observables mentioned in the introduction and the net of Borchers Uhlmann algebras we follow Dimock's construction of the former [Dim80, Dim82]. Recall that states on the Borchers Uhlmann algebra satisfying local (anti) commutativity define a (unique) field theoretical model. Let us denote the corresponding Bose and Fermi fields with Φ and Ψ respectively. Dimock constructs the net of local algebras with these field operators, showing that the result is independent of the state chosen before. Roughly speaking for scalar fields the local algebra $\mathfrak{A}(\mathcal{O})$ is the algebra generated by the exponentiated field operators $(e^{i\Phi(f)})$ smeared with testfunctions with support in $\mathcal{O}$. On the other hand for Fermi fields the corresponding algebra of local observables is generated by $\Psi(h)\Psi^+(h)$, with h having support in $\mathcal{O}$. For a description of models with more than one field things are greatly simplified by the fact that the Borchers Uhlmann algebras are tensor algebras over some spaces of test-function.

DEFINITION 3.7. Let $\mathfrak{A}_1, \dots, \mathfrak{A}_N$ be a finite number of Borchers Uhlmann algebras over spaces of test-functions $\mathcal{A}_1, \dots, \mathcal{A}_N$ respectively. The Borchers Uhlmann algebra over the space $\mathcal{A} = \bigoplus_{i=1}^N \mathcal{A}_i$ is called the *product* of these algebras and is denoted by $\mathfrak{A} := \mathfrak{A}_1 * \dots * \mathfrak{A}_N$, i.e., $\mathfrak{A}$ is the *free* tensor product of the algebras $\mathfrak{A}_1, \dots, \mathfrak{A}_N$.

REMARK . The monomial parts of $\mathfrak{A}$ are $\mathbb{C}, \mathfrak{A}_1 \oplus \dots \oplus \mathfrak{A}_N, (\mathfrak{A}_1 \oplus \dots \oplus \mathfrak{A}_N) \otimes (\mathfrak{A}_1 \oplus \dots \oplus \mathfrak{A}_N), \dots$, the unit element is given by $\mathbf{1} = (1, 0, \dots)$ and multiplication is the usual tensor product. This product –in difference to the usual tensor product of algebras– is not commutative. (Recall that for two algebras $\mathfrak{A}, \mathfrak{B}$, $\mathfrak{A} \otimes \mathfrak{B}$ is well defined. The embeddings are $\mathfrak{A} \xrightarrow{i} \mathfrak{A} \otimes \mathbf{1}, \mathfrak{B} \xrightarrow{i} \mathbf{1} \otimes \mathfrak{B}$. It follows that for $a \in \mathfrak{A}$ and $b \in \mathfrak{B}$ their product is given by $a \cdot b := (a \otimes \mathbf{1})(\mathbf{1} \otimes b) \equiv (a \otimes b) \equiv b \cdot a$).

We take the opportunity to introduce product states on these algebras. These states are used to implement the idea of 'fields not interacting with each other'.

DEFINITION 3.8. Let ω and ω_i be states on $\mathfrak{A} = \mathfrak{A}_1 * \dots * \mathfrak{A}_N$ and $\mathfrak{A}_i$ respectively. ω is called a *product state* if and only if all its m -point distributions ω^m can be written as products of the ω_i , e.g., let $\mathfrak{A} = \mathfrak{A}_1 * \mathfrak{A}_2$ and $\omega = \omega_1 \cdot \omega_2$ be given, then for $f_1 \otimes f_2 \otimes g \in \mathcal{A}_1 \otimes \mathcal{A}_1 \otimes \mathcal{A}_2$ the corresponding three point distribution ω^3 may read

$$\omega^3(f \otimes g_1 \otimes g_2) = \omega_1^2(f_1 \otimes f_2) \cdot \omega_2^1(g).$$

4. Hadamard states

The Hadamard condition for quasifree states of quantum fields on a manifold is believed to be a necessary condition for physical states since the work of DeWitt and Brehme in 1960 [DB60] and was intensively studied since this time by various authors (See the references in Fulling's book [Ful89]). Unfortunately it is a global condition which moreover, can be formulated for free fields only. The first mathematical rigorous formulation of this condition for free scalar fields is due to Kay and Wald [KW91]. We state their definition in this section for the convenience of the reader and extend it, using the ansatz of Najmi and Ottewill [NO84], to the fermionic case. *Scalar case:* Let ω be a quasifree state on the Borchers

Uhlmann algebra for a scalar field on a globally hyperbolic manifold (M, g_{ab}) satisfying the Klein-Gordon equation and local commutativity. Assume that a preferred time orientation has been chosen on (M, g_{ab}) and let T be a global time function increasing towards the future. Recall the definition of a convex normal neighborhood: An open subset $U \subset M$ is called a convex normal neighborhood if for all points $x_1, x_2 \in U$ there exists a unique geodesic *contained* in U connecting x_1 and x_2 . Let $\mathcal{O} \subset M \times M$ be an open neighborhood of the set of causally related points $(x_1, x_2) \in M \times M$, such that $J^+(x_1) \cap J^-(x_2)$ and $J^+(x_2) \cap J^-(x_1)$ are contained within a convex normal neighborhood. As usual $J^\pm(x)$ denotes the causal future (past) of the point x . Then the square of the geodesic distance, σ , is well defined and smooth on $\mathcal{O}$. For each integer p and $\epsilon \geq 0$ define for $(x, x') \in \mathcal{O}$ the complex valued function

$$G_\epsilon^{T,p}(x, x') = \frac{1}{(2\pi)^2} \left(\frac{\Delta^{1/2}(x, x')}{\sigma(x, x') + 2i\epsilon t + \epsilon^2} + v^{(p)}(x, x') \ln[\sigma(x, x') + 2i\epsilon t + \epsilon^2] \right) \quad (43)$$

where $t \equiv T(x) - T(x')$ and $\Delta^{1/2}$ and $v^{(p)}$ are smooth functions uniquely determined by the geometry on M ($\Delta^{1/2}$ is the van Vleck Morette determinant [DeW65] and $v^{(p)}$ is given by the Hadamard recursion relation up to order p (See Appendix B). The branch cut of the logarithm is taken to lie on the negative real axis). Let $\mathcal{C}$ be a Cauchy surface and let N be a neighborhood of $\mathcal{C}$ with the property: For each pair of points $x_1, x_2 \in N$ such that x_1 can be reached by a causal curve emerging from x_2 ($x_1 \in J^+(x_2)$) one can find a convex normal neighborhood in M containing $J^-(x_1) \cap J^+(x_2)$. N is called a causal normal neighborhood of $\mathcal{C}$. Now let $\mathcal{O}'$ be a neighborhood in $N \times N$ such that its closure is contained in $\mathcal{O}$ and choose a smooth real valued function χ on $M \times M$ with the properties $\chi(x, y) = 0$ if $(x, y) \notin \mathcal{O}$ and $\chi(x, y) = 1$ if $(x, y) \in \mathcal{O}'$.

DEFINITION 3.9 (Globally Hadamard states). We say the state ω mentioned above is *globally Hadamard* if and only if its two-point distribution² ω^2 is such that for each integer p there exists a C^p -function $H^{(p)}(x, y)$ on $N \times N$ such that for all $F_1, F_2 \in C_0^\infty(N)$ we have

$$\omega^2(F_1 \otimes F_2) = \lim_{\epsilon \rightarrow 0} \left(\int_{N \times N} \left(\chi(x, y) G_\epsilon^{T,p}(x, y) + H^{(p)}(x, y) \right) F_1(x) F_2(y) d\mu_x d\mu_y \right) \quad (44)$$

REMARK . Kay and Wald prove in [KW91] that this Definition is independent of the Cauchy surface $\mathcal{C}$ and the time function T . This in turn means that the Cauchy evolution preserves the Hadamard structure, i.e., ω^2 restricted to a causal normal neighborhood N of a Cauchy surface $\mathcal{C}$ already fixes ω^2 throughout the whole spacetime.

For a detailed discussion of this definition the reader is referred to the work of Kay and Wald [KW91]. *Fermi case:*

Let ω be a quasifree state on the Borchers Uhlmann algebra for a Majorana spinor field satisfying the Dirac equation on (M, g_{ab}) . In analogy to the scalar case the Hadamard condition is formulated in terms of its two point distribution ω^2 . Since the Dirac operator itself is not a strictly hyperbolic differential operator in the sense of Leray [Ler63] we introduce an auxiliary two point distribution $\tilde{\omega}^2$ such that for all $F_1 \in \Gamma(D^*M(N)) \xrightarrow{i} \hat{D}_1(N)$, $F_2 \in \Gamma(DM(N)) \xrightarrow{i} \hat{D}_1(N)$

$$\omega^2(F_1 \otimes F_2) = (i\mathcal{V}_x + m)\tilde{\omega}^2(F_1 \otimes F_2) \equiv \tilde{\omega}^2((i\mathcal{V} + m)F_1) \otimes F_2$$

where $\mathcal{V}$ is the Dirac operator (minus adjoint Dirac operator) on the spinor (cospinor) part of DM . Since ω^2 is a bi-solution of the Dirac equation it follows that $\tilde{\omega}^2$ is a bi-solution of the SUSY coupled Klein Gordon equation

$$\tilde{\omega}^2 \left(\left(\left[\square + \left\{ m^2 + \frac{1}{4}R \right\} \right] \mathbf{1}_{F_1} \right) \otimes F_2 \right) = \tilde{\omega}^2 \left(F_1 \otimes \left(\left[\square + \left\{ m^2 + \frac{1}{4}R \right\} \right] \mathbf{1}_{F_2} \right) \right) = 0.$$

Having chosen a moving frame E_A we write the components of the kernel of $\tilde{\omega}^2$ as³

$$\omega^2(h \otimes j) = (i\mathcal{V}_x + m\mathbf{1})^B_C \tilde{\omega}^2{}^C_{A'}(x, x') F_{1B}(x) F_2^{A'}(x').$$

²Note that this Definition discusses the Hadamard condition in terms of the (not symmetrized) two point distribution of the state. The symmetric two-point distribution, which is discussed usually in the literature, is the real part of ω^2 .

³For simplicity only the case $F_1 \in \Gamma(D^*M)$ and $F_2 \in \Gamma(DM)$ respectively is considered.

The SUSY coupled Klein Gordon operator is strictly hyperbolic and we define in analogy to the scalar case: The spinor state ω mentioned above is a (*spinor*) *Hadamard state* if and only if it's auxiliary two point distribution $\tilde{\omega}^2$ is a Hadamard distribution. More precisely: Let $(\mathcal{G}_\epsilon^{T,p})^B_{A'}(x, x')$ be the bi-spinor function

$$(\mathcal{G}_\epsilon^{T,p})^B_{A'}(x, x') = \frac{1}{(2\pi)^2} \left(\frac{\Delta^{1/2B}_{A'}(x, x')}{\sigma(x, x') + 2i\epsilon t + \epsilon^2} + v^{(p)B}_{A'}(x, x') \ln[\sigma(x, x') + 2i\epsilon t + \epsilon^2] \right)$$

where $\Delta^{1/2B}_{A'}$ is the spinor van Vleck Morette determinant and $v^{(p)B}_{A'}$ is given by the spinor version of the Hadamard recursion relation (See Appendix B).

DEFINITION 3.10 (Spinor Hadamard states). ω is a *spinor Hadamard state* if and only if it's auxiliary two point distribution $\tilde{\omega}^2$ is such that for each integer p there exist a C^p -function $H^{(p)}(\cdot, \cdot)$ on $\hat{D}_2(N)$ such that for all $F_1 \in \Gamma(D^*M(N)) \xrightarrow{i} \hat{D}_1(N)$, $F_2 \in \Gamma(DM(N)) \xrightarrow{i} \hat{D}_1(N)$ (in a local frame E_A) we have

$$\tilde{\omega}^2(F_1 \otimes F_2) = \lim_{\epsilon \rightarrow 0} \left(\int_{N \times N} \left(\chi(x, y) \mathcal{G}_\epsilon^{T,pB}_{A'}(x, x') \right) F_{1B}(x) F_{2A'}(x') d\mu_x d\mu_{x'} + H^{(p)}(F_1, F_2) \right)$$

REMARK . The reader might wonder why the operator in front of the auxiliary two-point distribution $\tilde{\omega}^2$ is $(i\nabla + m)$ and not $-(i\nabla + m)$; the second being the correct choice for the fundamental solution (See Proposition 4.16 below). To see the reason consider a free massive Majorana spinor field Ψ on Minkowski space. Let Ψ^+ denote its Dirac adjoint and $\Psi^\dagger$ its hermitian conjugate. Then one demands [IZ80]:

$$\{\Psi(x), \Psi^+(y)\}_+ = -iS(x, y) \equiv i(i\not{\partial} + m)E(x, y), \quad (45)$$

where S is the fundamental solution of the Dirac operator $(i\not{\partial} + m)$, and E is the fundamental solution of the Klein Gordon operator. The sign setting is required to ensure the equal time anti-commutation relations

$$\{\Psi(t, \mathbf{x}), \Psi^+(t, \mathbf{y})\}_+ = \gamma^0 \delta^{(3)}(\mathbf{x}, \mathbf{y}),$$

which means that $i\Psi^\dagger$ is conjugate to Ψ . Recall that this result is suggested by the corresponding Lagrangian (compare chapter 3) and that the l.h.s. of Eqn. (45) equals the imaginary part of ω^2 . Comparing Eqn. (45) with the corresponding relation for the scalar field, where one demands the commutator to be $+i$ times the fundamental solution E , one sees that $(i\nabla + m)$ must be used above for consistency.

The Hadamard condition in the formulation of Kay and Wald is a global condition, for example it requires continuity of $H^{(p)}$ on the *whole* spacetime. In his thesis Radzikowski [Rad92] recently gave a *local* characterization of the Hadamard condition in terms of wave front sets. However the proof of his claim is –as it stands– only valid for the rather trivial case of a flat spacetime. The proof presented below (See subsection 5.1) verifies his results in the general case. This wave front set characterization will be extremely useful for all following calculations in this thesis. It also motivates the wave front set spectrum condition, which is presented in the last section of this chapter.

5. Wave front sets

In this section some properties of the wave front set of a distribution are collected. For further details and the proofs which are omitted, we refer the reader to the original literature ([Hör71, DH72]) or to the monographs of Hörmander, Taylor and Reed & Simon [Hör83, Tay81, RS75]. The theory of wave front sets was developed in the seventies by Hörmander together with Duistermaat for their studies of pseudodifferential operators and differential equations on manifolds [Hör71, DH72]. Wave front sets (WF) are refinements of the notion of the singular support (sing supp) of a distribution. One main reason for using them in favor of the sing supp is that they provide a simple characterization for the existence of products of distributions and eliminate the difference between local and global results. It is interesting that Duistermaat and Hörmander even mention relations between their ‘micro-local analysis’ which they used to study solutions of the Klein-Gordon equation on a manifold and quantum field theory.

DEFINITION 3.11. Let $v \in D'(\mathbb{R}^n)$ be a distribution on $\mathbb{R}^n$. The set $\Sigma_z(v)$ is the complement in $\mathbb{R}^n \setminus \{0\}$ of the set of all nonzero $\xi \in \mathbb{R}^n$ for which there is a smooth function ϕ with compact support not vanishing at z and a conic neighborhood C_ξ of ξ such that for all $N \in \mathbb{Z}$ there exists a constant C_N such that for all $\xi' \in C_\xi$

$$(1 + |\xi'|)^N \left| \widehat{\phi u}(\xi') \right| \leq C_N$$

The $\widehat{}$ denotes Fourier transformation.

DEFINITION 3.12. The wave front set of v is defined by

$$\text{WF}(v) = \{(z, \theta) \in T^*\mathbb{R}^n \setminus \{0\} \mid \theta \in \Sigma_z(v)\} \quad (46)$$

All points $\langle x, \xi \rangle \in T^*\mathbb{R}^n$ which are *not* in the wave front set are called *regular directed points*.

REMARK .

1. $\text{WF}(v)$ is a closed subset of $T^*\mathbb{R}^n \setminus \{0\}$ since each regular directed point $\langle x, \xi \rangle \notin \text{WF}(v)$ has by definition an open neighborhood in $T^*\mathbb{R}^n \setminus \{0\}$ consisting of regular directed points, too.
2. The wave front set of a smooth function is the empty set
3. For $v \in D'(\mathbb{R}^n)$ with wave front set $\text{WF}(v)$, the projection of the wave front set to the base point gives the singular support of v .
4. For all smooth functions $\tilde{\phi}$ with compact support $\text{WF}(\tilde{\phi}v) \subset \text{WF}(v)$.
5. For two distributions $v, w \in D'(\mathbb{R}^n)$ with wave front sets $\text{WF}(v)$ and $\text{WF}(w)$ respectively, the wave front set of $(v + w) \in D'(\mathbb{R}^n)$ is contained in $\text{WF}(v) \cup \text{WF}(w)$

In order to extend this definition to manifolds we use the behavior of $\text{WF}(v)$ under diffeomorphisms of $\mathbb{R}^n$ to $\mathbb{R}^n$ (Theorem 3.13 below). Although Definition 3.11 contains a Fourier transform, it turns out that the elements of $\text{WF}(v)$ transform as elements of $T^*\mathbb{R}^n$, the cotangential bundle of $\mathbb{R}^n$. It is therefore possible to define the wave front set of a distribution on a manifold via charts.

THEOREM 3.13 (Theorem IX.44 and Problem 75 in [RS75]). *Let $v \in D'(\mathbb{R}^n)$ be a distribution on $\mathbb{R}^n$, χ be a diffeomorphism of $\mathbb{R}^n$ to $\mathbb{R}^n$ and let $v \circ \chi$ be the distribution⁴*

$$v \circ \chi(f) := v(g^{-1}(f \circ \chi^{-1}))$$

where g is the determinant of the Jacobian matrix $d\chi$. Define $\chi_* : \mathbb{R}^n \times (\mathbb{R}^n \setminus \{0\}) \rightarrow \mathbb{R}^n \times (\mathbb{R}^n \setminus \{0\})$ by

$$\chi_* \langle x, \xi \rangle = \langle \chi(x), d\chi^*(\xi) \rangle$$

where $d\chi^*$ is the adjoint of $d\chi$ with respect to the Euclidean inner product on $\mathbb{R}^n$. Then

$$\text{WF}(v \circ \chi^{-1}) = \chi_*(\text{WF}(v))$$

For the proof of this Theorem, which is only sketched in [RS75], we refer the reader to the monograph of Hörmander [Hör83, p. 265].

DEFINITION 3.14. Let $u \in D'(M)$ be a distribution over some manifold M . Let $\{X_\lambda\}_{\lambda \in \mathbb{N}}$ denote an open covering of M . Choose a compatible partition of unity $\{\Phi_\lambda\}_{\lambda \in \mathbb{N}}$. Without loss of generality we may assume that $\text{supp } \Phi_\lambda$ is contained in a single coordinate patch for every $\lambda \in \mathbb{N}$. The corresponding charts are denoted by $\chi_\lambda : \text{supp } \Phi_\lambda \subset X_\lambda \rightarrow \mathbb{R}^n$. Using the same notation as in Theorem 3.13 above we find as a Corollary

$$\chi_{\lambda*}(\text{WF}(\Phi_\lambda u)) \equiv \text{WF}(\Phi_\lambda u \circ \chi_\lambda^{-1}) \quad (47)$$

where $\Phi_\lambda u \circ \chi_\lambda^{-1}$ obviously is a distribution with compact support over $\mathbb{R}^n$. We define the *wave front set* of u by

$$\text{WF}(u) = \bigcup_{\lambda} \text{WF}(\Phi_\lambda u)$$

REMARK . This is a *local* definition. It follows that the following results which are stated in [RS75] for distributions on $\mathbb{R}^n$, are valid for distributions defined on manifolds, too.

A useful application of wave front sets is the definition of products of distributions. Wave front sets provide a simple characterization for the existence of such products, which furthermore can be extended to manifolds. The following definition of a product and its relation to wave front sets can be found for example in Reed & Simon [RS75, p.90–97].

DEFINITION 3.15. Let $v, w \in D'(\mathbb{R}^n)$. The distribution $T \in D'(\mathbb{R}^n)$ is the *product* of v and w if and only if for all $x \in \mathbb{R}^n$ there exists a smooth function f not vanishing at x such that for all $k \in \mathbb{R}^n$

$$\widehat{f^2 T}(k) = (2\pi)^{-n/2} \int_{\mathbb{R}^n} \widehat{f v}(l) \widehat{f w}(k-l) d^n l \quad (48)$$

where the integral is absolutely convergent.

⁴Note the misprint in [RS75].

REMARK .

- The product is well defined, since if such a T exists, it is unique:
Let $g \in D(\mathbb{R}^n)$ then

$$\widehat{gf^2T} = (2\pi)^{-n/2} \widehat{gf} \widehat{v} * \widehat{f} \widehat{w} = (2\pi)^{-n/2} \widehat{f} \widehat{v} * \widehat{g} \widehat{f} \widehat{w},$$

since the change of variables is legitimate due to the assumption of absolute convergence of (48). Now suppose T_1 and T_2 both fulfill (48), e.g. for all $x \in \mathbb{R}^n$ there exist functions f and g not vanishing at x such that $\widehat{f^2T_1} = \widehat{f} \widehat{v} * \widehat{f} \widehat{w}$ and $\widehat{g^2T_2} = \widehat{g} \widehat{v} * \widehat{g} \widehat{w}$. We conclude $\widehat{f^2g^2T_1} = \widehat{f^2g^2T_2}$, so $(T_1 - T_2)$ vanishes near x for all x , that is it is zero.

- The product of two distributions with disjoint support is zero.
- Since this is a local definition, we can extend it immediately to manifolds, using charts (See [Hör71]).

The following Theorem gives the relation between the wave front sets of two distributions and the existence of their product.

THEOREM 3.16 (Theorem IX.54 of [RS75]). *Let v, w be two distributions on M such that*

$$\text{WF}(v) \oplus \text{WF}(w) \equiv \{ \langle x, k_1 + k_2 \rangle \mid \langle x, k_1 \rangle \in \text{WF}(v), \langle x, k_2 \rangle \in \text{WF}(w) \}$$

does not contain any element of the form $\langle x, 0 \rangle$, then the product $v \cdot w$ exists and has wave front set

$$\text{WF}(v \cdot w) \subset \text{WF}(v) \cup \text{WF}(w) \cup (\text{WF}(v) \oplus \text{WF}(w))$$

Another interesting result, which will be used frequently in this work is the following Theorem of Hörmander [Hör71, Theorem 2.5.11].

THEOREM 3.17. *Let $\mathcal{C}$ be a sub-manifold of M with embedding φ and let*

$$N_\varphi = \{ (\varphi(y), \xi) \in T^*(M) \mid \langle \xi, d\varphi(y)\eta \rangle = 0 \quad \forall \eta \in T(M) \}$$

be the set of all normals of the map φ ($\langle \cdot, \cdot \rangle$ denotes dual pairing). If $u \in D'(M)$ and $\text{WF}(u) \cap N_\varphi = \emptyset$, we can define the restriction (pullback) $u|_{\mathcal{C}} \equiv \varphi^*u$ of u in one and only one way so that it is equal to the composition $u \circ \varphi$ when u is a continuous function and it is a sequentially continuous⁵ function from $D'_\Gamma(M) = \{u \in D'(M) \mid \text{WF}(u) \subseteq \Gamma\}$ to $D'(\mathcal{C})$ for any closed cone $\Gamma \subset T^*(M) \setminus \{0\}$ with $\Gamma \cap N_\varphi = \emptyset$. Moreover,

$$\text{WF}(\varphi^*(u)) \subset \varphi^*\text{WF}(u) = \{ (y, {}^t d\varphi(y)\xi) \mid (\varphi(y), \xi) \in \text{WF}(u) \},$$

where ${}^t d\varphi(y)$ denotes the transposition of $d\varphi(y)$ with respect to the dual pairing.

REMARK .

- The smooth functions of M are sequentially dense in D'_Γ for all admissible $\Gamma \in T^*M \setminus \{0\}$ (See [Hör71, p. 125]).
- Theorem 3.13 above may be viewed as a Corollary to this Theorem setting $\mathcal{C} \equiv M = \mathbb{R}^n$ (Compare [Hör83, p. 265]).

It is worth noting that the restriction mapping φ^* , due to its continuity, can be approximated by a suitable family of test densities:

COROLLARY 3.18. *Let $\mathcal{C}, M, \varphi$ as in Theorem 3.17 above be given and let $u \in D'(M)$ denote a distribution, such that φ^*u exists. Then for all smooth densities f on $\mathcal{C}$ there exists a family of smooth densities f_ρ on M , such that $u(f_\rho) \rightarrow \varphi^*u(f)$ for $\rho \rightarrow 0$.*

PROOF. Let $\{f_\rho\}$ be a family of smooth densities on M , such that for all smooth functions u_l on M $\lim_{\rho \rightarrow 0} u_l(f_\rho) = u \circ \varphi(f) \equiv \varphi^*u_l(f)$. Such families always exist due to the fact that the Dirac measure on M can be approximated by smooth densities (see e.g. [Die72, p. 242]). Now let $\{u_l\}$ denote a Cauchy sequence of smooth functions converging to u , then

$$\begin{aligned} \|\varphi^*u(f) - u(f_\rho)\| &\leq \|\varphi^*u(f) - \varphi^*u_l(f)\| + \|\varphi^*u_l(f) - u_l(f_\rho)\| + \|u_l(f_\rho) - u(f_\rho)\| \\ &\leq \epsilon, \end{aligned}$$

for $\rho \leq \delta$ and $l \geq N_\rho$, since φ^*u_l converges to φ^*u by Theorem 3.17, as well as $u_l(f_\rho)$ and u_l converge to $\varphi^*u_l(f)$ and u respectively. $\square$

In the sequel of this work, Theorem 3.17 above will be used for instance to define restrictions of distributions on $M \times M$ to the diagonal. As a second Corollary one finds:

⁵We refer the reader to Hörmanders original work for the definition of the pseudo topology D'_Γ is equipped with.

COROLLARY 3.19. *Let $f \in C_0^\infty(M \times M)$ be a smooth function on $M \times M$ and let $u \in D'(M)$ be an arbitrary distribution. Then $f \cdot u$ exists as a distribution on $M \times M$ and can be restricted to the diagonal.*

PROOF. Let $\text{WF}(u)$ denote the wave front set of u . Obviously $f \cdot u$ exists as a distribution on $M \times M$ with wave front set

$$\text{WF}(f \cdot u) = \{(x, 0; y, \xi) \in T^*(M \times M) \setminus \{0\} \mid x \in M, (y, \xi) \in \text{WF}(u)\}.$$

Let $\varphi : M \ni x \mapsto (x, x) \in M \times M$ denote the embedding of M onto the diagonal in $M \times M$. φ is smooth and for the set N_φ introduced above one finds:

$$N_\varphi = \{(x, \eta; x, -\eta) \in T^*(M \times M) \mid (x, \eta) \in T^*(M)\}.$$

Thus the assumption of Theorem 3.17 are satisfied, i.e.,

$$\text{WF}(f \cdot u) \cap N_\varphi = \emptyset,$$


which finishes the proof. $\square$

REMARK . We will denote the kernel of this special restriction $\varphi^*(f \cdot u)$ by $\varphi^*(f \cdot u)(y)$, $f(y, y)u(y)$ or $\lim_{x \rightarrow y} f(x, y)u(y)$ respectively.

5.1. The wave front set of a globally Hadamard state. The result of this subsection (Corollary 3.23 below) can already be found in the dissertation of Radzikowski. However the proof he gave is –as it stands– only valid for the rather trivial case of a flat spacetime. The proof presented here verifies his claim in the general case. Let ω^2 w.l.o.g. denote the two point distribution of a globally Hadamard state for a free scalar Klein Gordon field on (M, g_{ab}) . Following Radzikowski's line of argument, consider first the free massive Klein-Gordon field on Minkowski space in the vacuum state. The vacuum is known to fulfil the global Hadamard condition with $\Delta^{1/2} \equiv 1$ and $v^{(p)} \equiv \sum_{n=0}^p 2(\frac{m^2}{2})^{n+1}/n!(n+1)!$ ([CH84]). The wave front set of the corresponding two-point distribution $^{\text{Mk}}\omega^2$ is also well known (see for instance [RS75]):

THEOREM 3.20 (Theorem IX.48 of [RS75]). *The two-point distribution $^{\text{Mk}}\omega^2$ of the free massive scalar field on Minkowski space in the vacuum state has wave front set*

$$\begin{aligned} \text{WF}(^{\text{Mk}}\omega^2) &= \{(x_1, k_1), (x_2, k_2) \in \mathbb{R}^4 \times (\mathbb{R}^4 \setminus \{0\}) \mid x_1 \neq x_2; (x_1 - x_2)^2 = 0; \\ &\quad k_1 \parallel (x_1 - x_2); k_1 + k_2 = 0; \\ &\quad k_1^0 \geq 0\} \\ &\cup \{(x, k_1), (x, k_2) \in \mathbb{R}^4 \times (\mathbb{R}^4 \setminus \{0\}) \mid k_1 + k_2 = 0; k_1^2 = 0; k_1^0 \geq 0\} \end{aligned} \quad (49)$$

“=” 

For a proof see [RS75] or use the following representation of the Fourier transform of $^{\text{Mk}}\omega^2$:

$$\widehat{^{\text{Mk}}\omega^2} = (2\pi)^{-1} \delta(k_1 + k_2) \Theta(k_1^0) \delta(k_1^2 - m^2) \in \mathcal{S}'(\mathbb{R}^4 \times \mathbb{R}^4)$$

Using this representation and Definition 3.12 it is an easy, but lengthly calculation to verify Eqn. (49) explicitly. In order to extend this result to arbitrary (globally hyperbolic) manifolds, let $x \in M$ be a point on a Cauchy surface $\mathcal{C}$ and let U_x be a convex normal neighborhood of x . Denote by $\mathcal{C}_b \subset \mathcal{C}$ an arbitrary (non void) open subset in $\mathcal{C}$, containing x , such that the domain of causal dependence, $\diamond(\mathcal{C}_b)$, is contained in U_x . We are going to calculate the wavefront set of ω^2 for all base points $x_1, x_2 \in D^+(\mathcal{C}_b) := \diamond(\mathcal{C}_b) \cap J^+(\mathcal{C}_b)$ using some properties of Hadamard states on a special smooth deformation $(\tilde{M}, \tilde{g}_{ab})$ of our original spacetime⁶. Finally the propagation of singularities theorem (Theorem 3.21 below) and our knowledge that ω^2 has no singularities for spacelike separated points allows us to extend this result to the whole spacetime (M, g_{ab}) . Given $\mathcal{C}$, $x \in \mathcal{C}$ and $\mathcal{C}_b$ it will be shown below that there exists a globally hyperbolic spacetime $(\tilde{M}, \tilde{g}_{ab})$ with the following properties:

- (i) A neighborhood U of $\mathcal{C}$ in M is isometrically isomorphic to a neighborhood $\tilde{U}$ in $\tilde{M}$ and the isometry ϱ is also an isometry between $\mathcal{C}$ and $\tilde{\mathcal{C}}$; e. g. $(\tilde{M}, \tilde{g}_{ab})$ is a smooth deformation of (M, g_{ab}) .
- (ii) For all points $x_1, x_2 \in D^+(\mathcal{C}_b)$ there exists a Cauchy surface $\tilde{S}$ with neighborhood $\tilde{U}$ in $\tilde{M}$, such that the metric $\tilde{g}_{ab}$ restricted to $\tilde{U}$ is flat (i.e. Minkowskian) and $\diamond(\varrho(\mathcal{C}_b)) \subset \diamond(\tilde{U})$.

⁶The author thanks R. Verch for calling his attention to the deformation argument of Fulling, Narcowich and Wald [FNW81]. Note that the metric g_{ab} restricted to $D^+(\mathcal{C}_b)$ is *not* flat in general; the latter was assumed implicitly in Radzikowski's argument at some point, making the following modification of his proof necessary.

Now ω^2 induces canonically a Hadamard distribution $\tilde{\omega}^2$ on $(\tilde{M}, \tilde{g}_{ab})$, since it does so on $\tilde{U}$: Being a Hadamard distribution on $\tilde{U}$ implies, by the Remark following Definition 3.9, that $\tilde{\omega}^2$ is a Hadamard distribution throughout $(\tilde{M}, \tilde{g}_{ab})$. Moreover, the wave front set of $\tilde{\omega}^2$ on $\tilde{U}$ determines that of ω at U and vice versa by Theorem 3.13. On the other hand, using the propagation of singularities theorem and the smoothness of $\tilde{\omega}^2$ for spacelike separated points, the wave front set of $\tilde{\omega}^2$ at $\tilde{x}_1 = \varrho(x_1)$ and $\tilde{x}_2 = \varrho(x_2)$ is already fixed by the wave front set of that distribution at all points in $\hat{U}$ (See below and note that $\tilde{x}_1$ and $\tilde{x}_2 \in \diamond(\hat{U})$). To obtain the wave front set of ω^2 at x_1 and x_2 it is therefore sufficient to calculate the wave front set of $\tilde{\omega}^2$ for all points in $\hat{U}$. The following computations are performed entirely in the flat part of $\tilde{M}$ and were partly sketched in [Rad92]. Note first that Eqn. (44) is valid in $(\tilde{M}, \tilde{g}_{ab})$, too. To keep the notation simple we omit the $\sim$ in what follows. Choose a point $x \in S \equiv \hat{C}(t_1)$ and a convex normal neighborhood $U_x \subset \hat{U}$ of x . Using normal coordinates and an adapted time function T the distribution $G_\epsilon^{T,p}$, when restricted to $U_x \times U_x$ is

$$\begin{aligned} G^p &:= \lim_{\epsilon \rightarrow 0} \chi G_\epsilon^p \\ &= \lim_{\epsilon \rightarrow 0} \frac{\Delta^{1/2}(x_1, x_2)}{-(x_1 - x_2)^2 + 2i\epsilon(x_1^0 - x_2^0) + \epsilon^2} + v^{(p)}(x_1, x_2) \ln(-(x_1 - x_2)^2 + 2i\epsilon(x_1^0 - x_2^0) + \epsilon^2) \end{aligned}$$

The metric g_{ab} is flat on U_x and hence $2\sigma(x, y) \equiv -(x_1 - x_2)^2$ in these coordinates. Moreover the van Vleck Morette determinant $\Delta^{1/2}$ and $v^{(p)}$ are identical to the corresponding functions on Minkowski space, since only *local* properties of the underlying geometry enter into the Hadamard recursion relations. We conclude that $\text{WF}^{(\text{Mk}, \omega_{m=0}^2)} = \text{WF}(G^p)$ in normal coordinates on $U'_x \times U'_x$. Using Definition 3.14 this result can be pulled back to the manifold:

$$\begin{aligned} \text{WF}(G^p) &= \{(x_1, k_1), (x_2, k_2) \in (T^*M \times T^*M) \setminus \{0\} | \\ &\quad (x_1, k_1) \sim (x_2, -k_2); \quad k_1^0 \geq 0\} \quad \text{on } U'_x \times U'_x \end{aligned}$$

where $(x_1, k_1) \sim (x_2, k_2)$ means (i) x_1 and x_2 can be joined by a null geodesic γ , (ii) $k_1 (= k_{1\nu})$ is a cotangent vector such that $k_1^\mu = k_{1\nu} g^{\mu\nu}$ is tangential to γ , (iii) The parallel transport of k_1 along γ yields k_2 or (i) $x_1 = x_2$, (ii) $k_1^2 = 0$ and (iii) $k_1 = k_2$. Now $H^{(p)} \in C^p(M \times M)$ which implies for $\phi \in C_0^\infty(M)$ and $\phi(x) \neq 0$

$$((\phi \otimes \phi)H^{(p)})^\wedge(k_1, k_2) \leq C_p(1 + |k|)^{-p}$$

for a constant C_p and all $k = (k_1, k_2)$. Since $\omega^2 = G^p + H^{(p)}$ for all p we conclude that the wave front set of ω^2 restricted to $U'_x \times U'_x$ is

$$\begin{aligned} \text{WF}(\omega^2) &= \{(x_1, k_1), (x_2, k_2) \in (T^*M \times T^*M) \setminus \{0\} | \\ &\quad (x_1, k_1) \sim (x_2, -k_2); \quad k_1^0 \geq 0\} \quad \text{on } U'_x \times U'_x \quad (50) \end{aligned}$$

As a Corollary to the following Theorem of Hörmander this extends to all of $\tilde{M}$ (Compare section 2.2 in [Rad92]).

THEOREM 3.21 (Corollary to Theorem 6.1.1 of [DH72]). *Let $P = \mathbf{1} \otimes (\square + \nabla_\mu V^\mu + b)$ be a pseudodifferential operator on a globally hyperbolic manifold (M, g_{ab}) . $\square$ denotes the D'Alembert operator and V^μ and b are a smooth vector field and a smooth function on M respectively. The principal symbol of P is denoted by p and is given here by*

$$\begin{aligned} p : (T^*M \times T^*M) \setminus \{0\} &\rightarrow \mathbb{R} \\ (x_1, k_1; x_2, k_2) &\mapsto g^{\mu\nu}(x_2)k_{2\mu}k_{2\nu} \end{aligned}$$

If $u \in D'(M \times M)$ is a weak solution of $Pu = 0$ with wave front set $\text{WF}(u)$, it follows that

- a) $\text{WF}(u) \subseteq p^{-1}(0)$ and
- b) $\text{WF}(u)$ is invariant under the (Hamiltonian) vector field H_p given by

$$H_p = \sum_{i=1}^{2n} \frac{\partial p(x, k)}{\partial x_i} \frac{\partial}{\partial k_i} - \frac{\partial p(x, k)}{\partial k_i} \frac{\partial}{\partial x_i}$$

in local coordinates.

DEFINITION 3.22. The bi-characteristic strips of P are the curves on the sub-manifold $p^{-1}(0) \subset (T^*M \times T^*M) \setminus \{0\}$ which are generated by H_p .

Note that b) means: If $(x_1, k_1; x_2, k_2)$ and $(x'_1, k'_1; x'_2, k'_2)$ are on the same bi-characteristic strip, denoted by $(x_1, k_1; x_2, k_2) \approx (x'_1, k'_1; x'_2, k'_2)$ and $(x_1, k_1, x_2, k_2) \in \text{WF}(u)$, then $(x'_1, k'_1; x'_2, k'_2) \in \text{WF}(u)$. For $P = \mathbf{1} \otimes (\square + \nabla_\mu V^\mu + b)$ one finds (See Proposition 2.8 in [Rad92])

$$(x_1, k_1; x_2, k_2) \approx (x'_1, k'_1; x'_2, k'_2) \Leftrightarrow (x_1, k_1) = (x'_1, k'_1) \quad \text{and} \quad (x_2, k_2) \sim (x'_2, k'_2)$$

COROLLARY 3.23 (See Theorem 2.6 in [Rad92]). *The two-point distribution ω^2 of a free massive Klein-Gordon field on a globally hyperbolic spacetime in a globally Hadamard state has wave front set*

$$\text{WF}(\omega^2) = \{(x_1, k_1; x_2, -k_2) \in (T^*M \times T^*M) \setminus \{0\} \mid (x_1, k_1) \sim (x_2, k_2); \quad k_1^0 \geq 0\}$$

“=”  (51)

PROOF. Since ω^2 is a bisolution of the Klein-Gordon operator, we can apply Theorem 3.21 with $P_1 = \mathbf{1} \otimes (\square + m^2)$ or $P_2 = (\square + m^2) \otimes \mathbf{1}$. In the first case we conclude by a) that

$$\text{WF}(\omega^2) \subseteq \{(x_1, k_1; x_2, k_2) \in (T^*M \times T^*M) \setminus \{0\} \mid \quad k_2^2 = 0\}$$

and –using b)–

$$((x_1, k_1; x_2, k_2) \in \text{WF}(\omega^2) \wedge (x_2, k_2) \sim (x'_2, k'_2)) \Rightarrow ((x_1, k_1; x'_2, k'_2) \in \text{WF}(\omega^2)) \quad (52)$$

Obviously the second case leads to

$$\text{WF}(\omega^2) \subseteq \{(x_1, k_1; x_2, k_2) \in (T^*M \times T^*M) \setminus \{0\} \mid \quad k_1^2 = 0\}$$

and

$$((x_1, k_1; x_2, k_2) \in \text{WF}(\omega^2) \wedge (x_1, k_1) \sim (x'_1, k'_1)) \Rightarrow ((x'_1, k'_1; x_2, k_2) \in \text{WF}(\omega^2)) \quad (53)$$

To decide whether $(x_1, k_1; x_2, k_2)$ with $k_1^2 = k_2^2 = 0$ is in the wave front set of ω^2 , one checks first whether x_1 and x_2 are spacelike separated. If this is true, then $(x_1, k_1; x_2, k_2) \notin \text{WF}(\omega^2)$, since the projection of the wave front set of a distribution to the base point gives the singular support of that distribution, but ω^2 is assumed to have no singularities at spacelike separated points. If they are causally related, we use Eqn. (52) and (53) to propagate (x_i, k_i) along the null geodesic γ_i with tangent vector k_i at x_i to the Cauchy surface $\mathcal{C}$. Note that due to the global hyperbolicity of M this is always possible. If the points on $\mathcal{C}$ do not coincide, we have $(x_1, k_1; x_2, k_2) \notin \text{WF}(\omega^2)$, since different points on a Cauchy surface are spacelike separated. Should they coincide at say $x \in \mathcal{C}$, we check whether this particular combination $(x, k_1; x, k_2)$ is in the wave front set using Eqn. (50). $\square$




FIGURE 1. Spacetime Deformation

To finish the argument, we have to construct a spacetime $(\tilde{M}, \tilde{g}_{ab})$ that satisfies the properties (i) and (ii). This can be done by using the methods of Appendix C in [FNW81] (Compare Figure 1 and see also [Ver93]). Let $\diamond(\mathcal{C}_b) \subset V \subset N$ be a causal normal neighborhood of $\mathcal{C}$, such that $\mathcal{C}$ is also a Cauchy surface for V . V is, by the normal exponential map ϱ of $\mathcal{C}$, diffeomorphic to an open neighborhood $\tilde{V} \subset \mathbb{R} \times \mathcal{C}$ of $\tilde{\mathcal{C}} = \{0\} \times \mathcal{C} = \varrho(\mathcal{C})$. Using normal coordinates $(t, \mathbf{x})$ around some point in $\tilde{\mathcal{C}}$, the metric ϱ_*g on $\tilde{V}$ takes the form

$$dt^2 - h_{ij}(t, \mathbf{x}) dx^i dx^j$$

Choose $t_1 < t_2 < t_3 < 0$ such that

$$\hat{\mathcal{C}}(t_i) \cap J^-(\tilde{\mathcal{C}}_b) \subset \varrho(U_x) \quad i = 1, 2, 3,$$

where $\hat{\mathcal{C}}(t) := \{t\} \times \mathcal{C}$, $t \in \mathbb{R}$ and $\tilde{\mathcal{C}}_b := \varrho(\mathcal{C}_b)$ (As a consequence of the global hyperbolicity and due to the assumption $\diamond(\mathcal{C}_b) \subset U_x$ such a choice is always possible). Recall that $\varrho(U_x)$ can be covered with a single coordinate patch by assumption. Next choose a neighborhood $\tilde{U}$ of $\tilde{\mathcal{C}}$ such that $\tilde{U} \subset \tilde{V} \cap \text{int}(J^+(\hat{\mathcal{C}}(t_3)))$. Due to the fact that $t_1 < t_2$ there exists a neighborhood $\hat{U}$ of $J^-(\tilde{\mathcal{C}}_b) \cap \hat{\mathcal{C}}(t_1)$ in $\tilde{M}$ such that (i) $\tilde{U} \subset \varrho(U_x)$ and (ii) $\tilde{U} \cap J^+(\hat{\mathcal{C}}(t_2)) = \emptyset$. Let $f \in C^\infty(\mathbb{R} \times \mathcal{C}, \mathbb{R})$ be a smooth function $0 \leq f \leq 1$, $f \equiv 0$ on $\tilde{U}$ and $f \equiv 1$ outside the closure of $\tilde{V}$ or in the past of $J^-(\hat{\mathcal{C}}(t_2))$. Let $\tilde{h}$ be a complete Riemannian metric on $\mathcal{C}$ being flat on the spatial component of $\hat{U} \cap (\hat{\mathcal{C}}(t_3))$. Note that the existence of such a $\tilde{h}$ follows from the fact that $\tilde{U}$ can be covered by a single coordinate patch (i.e. $\varrho(U_x)$). Let $\beta \in C^\infty(\mathbb{R} \times \mathcal{C}, \mathbb{R}^+)$ be a function equal to one on $\tilde{U}$ and on $(-\infty, t_2)$. Define a Lorentzian metric $\tilde{g}_{ab}$ on $\mathbb{R} \times \mathcal{C}$ by setting in the coordinates above the coordinate expression of $\tilde{g}_{ab}$ equal to

$$\beta(t, \mathbf{x}) dt^2 - ((1 - f(t, \mathbf{x})) h_{ij}(t, \mathbf{x}) + f(t, \mathbf{x}) \tilde{h}_{ij}(t, \mathbf{x})) dx^i dx^j$$

By choosing β sufficiently small outside the region where it is demanded to be one⁷, we can ensure that $(\tilde{M} := \mathbb{R} \times \mathcal{C}, \tilde{g}_{ab})$ is globally hyperbolic. Setting $U := \varrho^{-1}(\tilde{U})$ and $\tilde{S} := \varrho^{-1}(\hat{\mathcal{C}}(t_1))$ finishes the construction.

A first consequence of Corollary 3.23 is

COROLLARY 3.24. *Let ω^n be the n -point distribution arising from a quasifree Hadamard state of a massive Klein-Gordon field propagating on a globally hyperbolic spacetime (M, g_{ab}) . Then all finite powers of ω^n exist as products of distributions.*

PROOF. Assuming all odd n -point distributions to vanish, we have, since ω^n arises from a quasifree state

$$\omega^n(x_1, \dots, x_n) = \sum_P \prod_r \omega^2(x_{(r,1)}, x_{(r,2)}),$$

where P denotes a partition of the set of points $\{x_i\}$ into subsets which are pairings of points, labeled by r . Note that the ordering of the points in ω^2 is preserved, e.g. $(r, 1) < (r, 2)$ and no two arguments are identical. The latter fact ensures the existence of the product $\prod_r$ whenever $\omega^2(x_i, x_j)$ are distributions. For the wave front set of ω^n one finds using Theorem 3.16

$$\begin{aligned} \text{WF}(\omega^n) &= \bigcup_{(x_1, \dots, x_n) \in M^n} \text{WF}(\omega^n(x_1, \dots, x_n)) \\ &\subseteq \bigcup_{(x_1, \dots, x_n) \in M^n} \left(\bigcup_P \bigcup_p \bigoplus_{r_p} [M^{n-2} \times \text{WF}(\omega^2(x_{(r_p,1)}, x_{(r_p,2)}))]\right), \end{aligned} \tag{54}$$

where p denotes a subset of P , r_p labels the elements of p and

$$\begin{aligned} &[M^{n-2} \times \text{WF}(\omega^2(x_{(r_p,1)}, x_{(r_p,2)}))] \\ &= \{(x_1, 0; \dots; x_{(r_p,1)}, k_{(r_p,1)}; \dots; x_{(r_p,2)}, k_{(r_p,2)}; \dots; x_n, 0) \in (T^*M)^n \setminus \{0\} | \\ &\quad (x_{(r_p,1)}, k_{(r_p,1)}; x_{(r_p,2)}, k_{(r_p,2)}) \in \text{WF}(\omega^2(x_{(r_p,1)}, x_{(r_p,2)}))\}. \end{aligned}$$

(Compare Example 3.25 below, which demonstrates the notation on the four point function). To prove the Corollary it is –by Theorem 3.16– necessary and sufficient to show that finite sums of $\text{WF}(\omega_n)$ do not contain zero. We know the wave front sets of all distributions in Eqn. 54 explicitly (See Corollary 3.23). Consider the following two cases:

1. The direction in $\text{WF}(\omega_n)$ associated to the first variable is not zero. Then its time component is strictly positive by Corollary 3.23. Since the time components of all directions associated to the first variable of $\text{WF}(\omega_n)$ are always greater or equal to zero, the sum of all these directions can not vanish and the Corollary is proved.
2. The direction in $\text{WF}(\omega_n)$ associated to the first variable is zero for all summands. Then all directions associated to the second variable must have time components greater or equal to zero and we apply the same argumentation as for the first variable. Note that it is excluded by the definition of the wave front set that all directions of all variables vanish simultaneously. Therefore there exists a variable (x_i) $i < n$ such that the analogon of (1) holds.

□

EXAMPLE 3.25. Consider the four-point distribution of a quasifree state:

$$\omega^4(x_1, x_2, x_3, x_4) = \omega^2(x_1, x_2)\omega^2(x_3, x_4) + \omega^2(x_1, x_3)\omega^2(x_2, x_4) + \omega^2(x_1, x_4)\omega^2(x_2, x_3)$$

We have

$$P \in \{\{(x_1, x_2), (x_3, x_4)\}, \{(x_1, x_3), (x_2, x_4)\}, \{(x_1, x_4), (x_2, x_3)\}\}$$

⁷I.e. we “close” up the light cone.

Let $P = \{(x_1, x_3), (x_2, x_4)\}$, then $p \in \{\{(x_1, x_3)\}, \{(x_2, x_4)\}, \{(x_1, x_3), (x_2, x_4)\}\}$. Assume $p = \{(x_1, x_3)\}$, then $r_p = (x_1, x_3)$, i.e. $x_{(r_p, 1)} = x_1$ and $x_{(r_p, 2)} = x_3$. Therefore

$$\begin{aligned}
& \text{WF}(\omega^4) \\
& \subseteq \bigcup_{(x_1, \dots, x_4) \in M^4} \left([M^2 \times \text{WF}(\omega^2(x_1, x_2))] \cup [M^2 \times \text{WF}(\omega^2(x_3, x_4))] \right. \\
& \quad \left. \cup [(M^2 \times \text{WF}(\omega^2(x_1, x_2))) \oplus (M^2 \times \text{WF}(\omega^2(x_3, x_4)))] \right. \\
& \quad \left. \cup 2 \leftrightarrow 3 \right. \\
& \quad \left. \cup 2 \leftrightarrow 4 \right) \\
& \equiv \{(x_1, k_1; x_2, k_2; x_3, 0; x_4, 0) \mid (x_1, k_1; x_2, k_2) \in \text{WF}(\omega^2)\} \\
& \quad \cup \{(x_1, 0; x_2, 0; x_3, k_3; x_4, k_4) \mid (x_3, k_3; x_4, k_4) \in \text{WF}(\omega^2)\} \\
& \quad \cup \{(x_1, k_1; x_2, k_2; x_3, k_3; x_4, k_4) \mid (x_1, k_1; x_2, k_2) \in \text{WF}(\omega^2); (x_3, k_3; x_4, k_4) \in \text{WF}(\omega^2)\} \\
& \quad \cup \dots
\end{aligned}$$

We can extend Corollary 3.23 and Corollary 3.24 immediately to the fermionic case, since both, the spinor van Vleck Morette determinant $\Delta^{1/2 B}_{A'}$ and $v^{(p) B}_{A'}$ are smooth near the diagonal for all spinor indices A', B and the Dirac operator, which acts as a differential operator, does not enlarge the wave front set.

COROLLARY 3.26. *The two point distribution $\omega^2_{A^{A'}}$ of a free Majorana spinor field in a globally Hadamard state has wave front set contained in*

$$\text{WF}(\omega^2_{A^{A'}}) \subseteq \{(x_1, k_1), (x_2, -k_2) \in T^*M \setminus \{0\} \times T^*M \setminus \{0\} \mid (x_1, k_1) \sim (x_2, k_2); \quad k_1^0 \geq 0\} \quad (55)$$

The extension of Corollary 3.24 above to the fermionic case reads:

COROLLARY 3.27. *Let $\omega^n_{A^{A'}}$ be the n -point distribution arising from a quasifree Hadamard of a massive Majorana spinor field propagating on a globally hyperbolic spacetime (M, g_{ab}) . Then all finite powers of $\omega^n_{A^{A'}}$ exist as product of distributions.*

PROOF. For fixed Spinor indices the wave front set of the fermionic two-point distribution is contained in the wave front set for the scalar field (See Eqn. (55)). Applying the same argumentation as in the proof of Corollary 3.24 finishes the proof. $\square$

REMARK . It is worth noting that for the wave front set of all finite powers of all Hadamard two point distributions ω^2 satisfies

$$\text{WF}(\omega^2 \dots \omega^2) \subseteq (\text{WF}(\omega^2) \oplus \text{WF}(\omega^2)) \cup \text{WF}(\omega^2). \quad (56)$$

This relation follows by Theorem 3.16 together with the observation that the first summand in Eqn. (56) is invariant under further additions of $\text{WF}(\omega^2)$, i.e.,

$$\text{WF}(\omega^2) \oplus \text{WF}(\omega^2) \oplus \text{WF}(\omega^2) = \text{WF}(\omega^2) \oplus \text{WF}(\omega^2)$$

5.2. A local characterization of globally Hadamard states. One of the main motivations for using wave front sets in quantum field theory is the fact that they allow the specification of global properties locally. The following Theorem, which is one of the main results of Radzikowski's dissertation, gives a local characterization of globally Hadamard states.

THEOREM 3.28 (Theorem 2.6 of [Rad92]). *Let ω^2 be the two-point distribution arising from a state of a massive Klein-Gordon field propagating on a globally hyperbolic spacetime. If ω^2 has wave front set as in Corollary 3.23 above then ω^2 is globally Hadamard.*

REMARK . Our assumption implies that ω^2 fulfils the Klein-Gordon equation and has i times the commutator distribution of $\square + m^2$ as its antisymmetric part. The Corollary 3.23 above may be viewed as the converse of this Theorem.

The proof of this Theorem is rather long; we refer the reader to the dissertation of Radzikowski for the details. Radzikowski's Theorem 2.6 states the equivalence of the wave front set assumption and the globally Hadamard condition $\text{mod } C^\infty$. To prove it, he introduces the Feynman two-point distribution ω_F defined by $\omega_F := i\omega^2 + E^+$, where E^+ is the advanced fundamental solution of $\square + m^2$. The assumption on the wave front set of ω^2 uniquely fixes this distribution (up to C^∞) to be the distinguished Feynman parametrix defined in [DH72]. This in turn implies that ω^2 is globally Hadamard $\text{mod } C^\infty$.

6. The wave front set spectrum condition

In this section we propose a condition on the wave front sets of states of a quantum field on a manifold M , which might generalize the usual Minkowski spectrum condition to Lorentz manifolds. The idea to use wave front sets for a formulation of some kind of spectrum condition is due to Radzikowski [Rad92]. For a motivation of this idea recall that the wave front set of a state can be used to characterize this state globally. However Equation (51) restricts the singular support of $\omega_2(x_1, x_2)$ to points x_1 and x_2 which are null related; hence ω_2 is smooth for spacelike related points. This smoothness is known to be true for reasonable quantum field theories on Minkowski space satisfying the true spectrum condition by the Bargmann-Hall-Wightman Theorem. For timelike related points however a similar general prediction on the smoothness does not exist. In order to include possible singularities at timelike related points, Radzikowski extended in [Rad92] the right hand side of Eqn. (51) to all causally related points. He proposed that the wave front set of the two-point distributions of any physically reasonable state should be contained in this extended set and called this proposal the 'wave front set spectrum condition' (WFSSC). He also proposed a WFSSC for higher n -point distributions and showed that both of his Definitions are compatible to the usual spectrum condition: Each ω^n fulfilling his WFSSC satisfies the true spectrum condition on Minkowski space $\text{mod } C^\infty$ (Theorem 4.10 of [Rad92]) and vice versa. He gives further evidence on the legitimacy of his Definitions by linking them to the scaling limit condition of Fredenhagen and Haag [FH87]: Both Definitions imply the true spectrum condition in the scaling limit if this limit exists (Theorem 4.11 of [Rad92]). Unfortunately it can be shown that the n -point distributions for $n > 2$ associated to a quasifree Hadamard state of a scalar field on a globally hyperbolic spacetime do not satisfy his WFSSC in general. Moreover, the Wick products defined below give rise to counterexamples even for his two point WFSSC. Thus his original WFSSC needs to be modified. The counterexample mentioned above and the wish to include fields which are Wick products of 'simple' fields, leads us to propose the following 'conic' WFSSC⁸. Let $\mathcal{G}_n$ denote the set of all undirected finite graphs⁹ with vertices $\{1, \dots, n\}$,




FIGURE 2. An immersion of a graph

such that for every element $G \in \mathcal{G}$ all edges occur in both admissible directions. An *immersion* of a graph $G \in \mathcal{G}_n$ into some manifold M is an assignment of the vertices of G to points in M , $\nu \rightarrow x(\nu)$, and of the edges of G to causal curves in M , $e \rightarrow \gamma(e)$, together with a covariantly constant causal covector field k_e on γ , such that

1. If e is an edge from ν to ν' then $\gamma(e)$ connects $x(\nu)$ and $x(\nu')$,
2. If e^{-1} denotes the edge with opposite direction as e , then the corresponding curve $\gamma(e^{-1})$ is the inverse of $\gamma(e)$,
3. For every edge e from ν to ν' , k_e is directed towards the future whenever $\nu < \nu'$,
4. $k_{e^{-1}} = -k_e$.

(Compare Figure 2)

⁸I.e., $(x, k_1; \dots; y, l_1), (x, k_2; \dots; y, l_2) \in \text{WF}(\omega_n) \Rightarrow (x, \lambda k_1 + \mu k_2; \dots; y, \lambda l_1 + \mu l_2) \in \text{WF}(\omega_n) \quad \forall \lambda, \mu > 0$.

⁹We allow each two vertices of a graph to be connected by multiple edges.


DEFINITION 3.29 (GWFSSC). A state ω with n -point distributions ω^n is said to satisfy the *general wave front set spectrum condition* (GWFSSC) if and only if

$$\text{WF}(\omega^n)$$

$$\subseteq \left\{ (x_1, k_1; \dots; x_n, k_n) \in T^*M^n \setminus \{0\} \mid \exists G \in \mathcal{G}_n \text{ and an immersion } (x, \gamma, k) \text{ of } G, \right.$$

such that (i) $x_i = x(i) \quad \forall i = 1, \dots, n$

$$\left. \text{(ii) } k_i = \sum_{\substack{e \\ i \text{ is starting} \\ \text{point of } e}} k_e(x_i) \right\}$$



REMARK . For every set of base points $(x_1, \dots, x_n) \in \text{sing supp } (\omega^n)$ the first non zero direction k_l in the wave front set is future directed.

As an important consequence of this Definition we have the following

PROPOSITION 3.31. *Let ω be a state which satisfies the GWFSSC, then all finite powers of its n -point distributions exist.*

PROOF. To prove the Proposition it is –by Theorem 3.16– sufficient to show that finite sums of $\text{WF}(\omega^n)$ do not contain zero. Consider the following two cases:

1. The direction in $\text{WF}(\omega^n)$ associated to the first variable is not zero. Then its time component is strictly positive since the sum of future directed covectors is again future directed. Thus the sum of *all* directions associated to the first variable can not vanish and the Proposition is proved.
2. The direction in $\text{WF}(\omega^n)$ associated to the first variable is zero for all summands. Then all directions associated to the second variable must have time components greater or equal to zero and we apply the same argumentation as for the first variable. Note that it is excluded by the definition of the wave front set that all directions of all variables vanish simultaneously. Therefore there exists a variable $(x_i) \ i < n$ such that the analogon of (1) holds.

□

To show that there exists non trivial states which verify our Definition we prove the following

LEMMA 3.32. *Let ω denote a quasifree Hadamard state for the Klein-Gordon field on a globally hyperbolic manifold (M, g_{ab}) , then ω satisfies the general wave front set spectrum condition.*

PROOF. Note first that all odd n -point distributions vanish by assumption, hence ω^n satisfies the GWFSSC trivially for odd n . Consider the two point distribution of ω . Its wave front set is explicitly known (see Proposition 3.23) and obviously satisfies the GWFSSC. For a general n -point distribution, with n even, consider a point $p = (x_1, k_1; \dots; x_n, k_n) \in \text{WF}(\omega_n)$. Using the same notation as in Definition 3.24 and Eqn. (54) this point can be written as

$$p = (x_1, \sum_{i=1}^{n/2} k_{1_i}; \dots; x_n, \sum_{j=1}^{n/2} k_{n_j}),$$

where the directions k_{l_m} are suitable chosen with the constraint $(x_l, k_{l_i}; x_m, k_{m_j}) \in \text{WF}(\omega^2)$ or $(x_l, k_{l_i}; x_m, k_{m_j}) = (x_l, 0; x_m, 0)$. We remark that this representation follows trivially from Eqn. (54). Now by Eqn. (51), whenever $k_{l_i}, k_{m_j} \neq 0$, $(x_l, k_{l_i}) \sim (x_m, -k_{m_j})$, i.e., x_l and x_m are immersed vertices ν_l and ν_m of some graph G , such that the corresponding edge e connecting ν_l and ν_m has an immersion $\gamma(e)$, which is the lightlike geodesic connecting x_l and x_m . Moreover,

there exists a covector field k_e such that $k_e(x_l) = k_{l_i}$ and $k_{e^{-1}}(x_m) = -k_{m_j}$. Note that k_{l_i} is directed towards the future whenever $l < m$. Thus G with this immersion (x, γ, k) satisfy (i) and (ii) in the Definition of the GWFSSC. $\square$

We close this chapter by giving our definition of the quantized version of our locally supersymmetric model using the algebraic approach: A scalar quantum field theoretical model $(\mathcal{H}, D, \Phi, \Omega)$ on a spacetime (M, g_{ab}) which we assume to be orientable, time orientable and globally hyperbolic, is said to satisfy the Klein Gordon equation (with coupling ξ) if and only if

$$\text{For all } f \in C_0^\infty(M): \quad \Phi(K_\xi f) = 0,$$

where $K_\xi := (\square + (m^2 + \xi R))$ is the generic Klein Gordon operator associated to g_{ab} . A Majorana quantum field theoretical model $(\mathcal{H}, D, \Psi, \Psi^+, \Omega)$ on the same spacetime (M, g_{ab}) with spin structure $(SM, \mathbf{p})$ is said to satisfy the Dirac equation if and only if

$$\text{For all } f \in \Gamma(DM(\mathcal{O})) \xrightarrow{i} \hat{D}_1(\mathcal{O}): \quad \Psi(\not{D}f) = 0$$

and

$$\text{For all } f^+ \in \Gamma(D^*M(\mathcal{O})) \xrightarrow{i} \hat{D}_1(\mathcal{O}): \quad \Psi^+(\not{D}^+ f^+) = 0$$

where $\not{D} := (i\nabla - m)$ ($\not{D}^+ := (-i\nabla - m)$) is the generic Dirac (adjoint Dirac) operator on the spinor (cospinor) bundle. The states of these models fulfill the same equation of motion as the corresponding fields. Let now $\mathfrak{A}_1, \mathfrak{A}_2$ be two scalar Borchers Uhlmann algebras and $\hat{\mathfrak{A}}_3$ be a Majorana Borchers Uhlmann algebra. Denote by $\mathfrak{A}$ the product $\mathfrak{A}_1 * \mathfrak{A}_2 * \hat{\mathfrak{A}}_3$ of these three algebras. $\mathfrak{A}$ is called a *locally supersymmetric* Borchers Uhlmann algebra. A product state $\omega = \omega_1 \cdot \omega_2 \cdot \omega_3$ on $\mathfrak{A}$ such that each $\omega_{1,2}$ defines a scalar quantum field theoretical model satisfying the Klein Gordon equation with $\xi = 1/4$, whereas ω_3 gives a Majorana quantum field theoretical model satisfying the Dirac equation, defines a *locally supersymmetric quantum field theoretical model* on the spacetime (M, g_{ab}) . For convenience such a state will be called a local SUSY product state (on the local SUSY Borchers Uhlmann algebra $\mathfrak{A}$).

The energy momentum tensor

It was shown in chapter 1 that there is a close relation between the (conserved) supercurrent and the energy momentum tensor in the free Wess-Zumino model on Minkowski space. We expect a similar relation on a curved spacetime, too. It seems therefore reasonable to *define* a candidate for the energy momentum tensor of a locally supersymmetric model on a curved spacetime by an analogon of Eqn. (30) using the improved ‘supercurrent’, i.e., by inserting the improved ‘supercurrent’ (Eqn. 39) into Eqn. (30). It is worth noting that this substitution is necessary in order to obtain the *improved* stress energy tensor instead of the canonical one. Unfortunately there are some conceptual and mathematical difficulties with this approach which will be discussed in the sequel:

1. In classical general relativity the energy momentum tensor T^{ab} of the matter fields has to be locally conserved. Recall that this assumption reflects the well established fundamental law of energy conservation. Moreover, it is required as a consistency condition in the Einstein equations, where T^{ab} acts as a source term¹. It is easy to see –at least in Minkowski space–, that only a *locally conserved* supercurrent leads to a conserved energy momentum tensor. The fact that k^{aA} is locally conserved on Lorentz manifolds with vanishing scalar curvature only, indicates that our Definition is applicable at most to those special spacetimes.
2. A mathematical difficulty comes from the fact that on a general Lorentz manifold the integration of a spinor vector current, such as j^{aA} , over some spacelike hypersurface $\mathcal{C}$ with normal covector field n_a does not make sense, since the parallel transport of spinors is path dependent. Recall that such an integration was necessary on Minkowski space for the definition of the spinor supercharge Q^A . However locally, i.e., for a sufficiently small region $\mathcal{O}_{\mathcal{C}}$ of $\mathcal{C}$ there exists an obvious solution to this problem: The expression $j^{aA}(x)\mathcal{J}_A^{A'}(x, x')$, where $\mathcal{J}_A^{A'}(x, x')$ denotes the bi-spinor of parallel transport introduced in chapter 3 transforms as a cospinor at x and as a spinor at x' , hence the integral

$$\int_{\mathcal{O}_{\mathcal{C}}} j^{aA} \mathcal{J}_A^{A'} n_a d\mathcal{C} \quad (57)$$

does make sense (at least at the classical level). Moreover, since $\lim_{x \rightarrow x'} (\nabla_a \mathcal{J}_A^{A'}) = 0$, we have “asymptotic conservation” of $j^{aA} \mathcal{J}_A^{A'}$, i.e., the coincidence limit $\lim_{x \rightarrow x'} (\nabla_a (j^{aA} \mathcal{J}_A^{A'}))$ vanishes whenever j^{aA} is locally conserved. Note however that the integral (57) obviously depends on x' .

In the remaining part of this work we will roughly speaking investigate the expression

$$+ \frac{1}{8} \left(\gamma^{(a} \{j^{b)}, Q^+\}_{+} \right), \quad (58)$$

for a quantized version of our locally supersymmetric model. Q^+ is the ‘supercharge’ associated to k^{aA} and the point separated version of Eqn. (58) is regarded as an operator valued distribution on a suitable Hilbertspace (See below for the actual definitions).

REMARK . The use of k^{aA} instead of j^{aA} for the definition of the ‘supercharge’ Q^+ is motivated by the observation that in the Wess-Zumino model the divergence, by which j^{aA} and k^{aA} differ, does not contribute to the supercharge G .

1. New Wightman fields: The ‘supercurrents’ after quantization

Consider the locally supersymmetric Borchers Uhlmann algebra $\mathfrak{A}$ on (M, g_{ab}) together with a locally SUSY Hadamard product state ω , i.e., $\omega = \omega_1 \cdot \omega_2 \cdot \omega_3$ is a local SUSY product state, such that ω_1, ω_2 and ω_3 are three Hadamard states. Let $(\mathcal{H}, \mathcal{D}, \mathcal{A}, \mathcal{B}, \Psi, \Psi^+, \Omega)$ denote the corresponding quantum field theoretical model, where A and B are scalar and pseudo-scalar fields respectively and Ψ denotes a Majorana spinor field. Recall that $\mathcal{H}$ is the GNS Hilbertspace of ω and that Ω denotes the corresponding cyclic vector. Let us study the improved ‘supercurrent’

$$\begin{aligned} j^{aA} = & i(A - i\gamma^5 B)(i\overleftarrow{\not{\nabla}} - m)\gamma^a \Psi \\ & - 2\xi(\gamma^a \gamma^b - \gamma^b \gamma^a) \nabla_b ((A - i\gamma^5 B)\Psi), \end{aligned} \quad (59)$$

¹For a discussion of this assumption see e.g. [HE73].

which involves the pointwise product of the basic fields. We find

PROPOSITION 4.1. j^{aA} is well defined and gives a new Wightman field on the GNS Hilbertspace $(\mathcal{H}, A, B, \Psi, \Omega)$ of every quasifree globally Hadamard product state ω .

PROOF. We show that j^{aA} is well defined and fulfils the following four Wightman axioms:

1. For all $f \in \Gamma_0(D^*M \times T^*M)$, j^{aA} is a well defined operator valued distribution on the GNS Hilbertspace (H, Ω, A, B, Ψ) with domain

$$D = \text{Span}\{\Phi(f_1) \cdots \Phi(f_n)\Omega \mid \Phi \in (A, B, \Psi, j^{aA}), f_1, \dots, f_n \in (C_0^\infty(M), \Gamma_0(D^*M), \Gamma_0(D^*M \otimes T^*M))\}$$

By $\Gamma_0(\cdot)$ we denote sections with compact support.

2. D is dense in H .
3. j^{aA} leaves D invariant.
4. j^{aA} is anti-local (e.g. fermionic).

To simplify the notation, these four Wightman axioms are actually shown for the k^{aA} part of j^{aA} only. However the proof below is obviously equally valid for j^{aA} itself.

1. Let $D' = \text{Span}\{\Psi = k(f_1) \cdots k(f_n)\Omega \mid f_i \in \Gamma_0(D^*M \otimes T^*M)\}$. We remark that the definition of D' is only formal at this stage. To show that k^{aA} is well defined on D' , it is sufficient to show that

$$\|k(f)\Psi\| < \infty \quad \forall f \in \Gamma_0(D^*M \times T^*M), \quad \Psi = k(f_1) \cdots k(f_n)\Omega \in D' \quad (60)$$

Choose a moving frame in the spinor bundle (E_A) which induces moving frames in the cospinor bundle (E^A) , in the tangent bundle (e_μ) and in the cotangent bundle (e^μ) . Inserting the definitions of k and Ψ in Equation (60) we obtain in these local coordinates:

$$\begin{aligned} & \|k(f)\Psi\|^2 \\ &= \langle k(f)k(f_1) \cdots k(f_n)\Omega \mid k(f)k(f_1) \cdots k(f_n)\Omega \rangle \\ &= \langle \Omega \mid k^+(f_n^+) \cdots k^+(f_1^+)k^+(f^+)k(f)k(f_1) \cdots k(f_n)\Omega \rangle \\ &= \int \cdots \int d\mu_{x_n} \cdots d\mu_{x_1} d\mu_x d\mu_y d\mu_{y_1} \cdots d\mu_{y_n} \\ & \quad \Psi_{\omega_{2n+2} A_n \cdots A_1 A}^{A' A'_1 \cdots A'_n} (x_n, \dots, x_1, x, y, y_1, \dots, y_n) \gamma^{a_n A_n}_{B_n} \cdots \gamma^{a_1 A_1}_{B_1} \gamma^{a A}_{B'} \\ & \quad \left((-i \overrightarrow{\nabla}_{x_n} - m)^{B_n}_{C_n} \cdots (-i \overrightarrow{\nabla}_x - m)^B_C (i \overrightarrow{\nabla}_y - m)^{C'}_{B'} \cdots (i \overrightarrow{\nabla}_{y_n} - m)^{C'_n}_{B'_n} \right. \\ & \quad \left. {}^A \omega_{2n+2}(x_n, \dots, x_1, x, y, y_1, \dots, y_n) \right. \\ & \quad \left. - ((-i \overrightarrow{\nabla}_{x_n} - m) \gamma^5)^{B_n}_{C_n} \cdots ((-i \overrightarrow{\nabla}_x - m) \gamma^5)^B_C (\gamma^5 (i \overrightarrow{\nabla}_y - m))^{B'}_{C'} \cdots (\gamma^5 (i \overrightarrow{\nabla}_{y_n} - m))^{B'_n}_{C'_n} \right. \\ & \quad \left. {}^B \omega_{2n+2}(x_n, \dots, x_1, x, y, y_1, \dots, y_n) \right) \\ & \quad \gamma^{a' B'}_{A'} \gamma^{a'_1 B'_1}_{A'_1} \cdots \gamma^{a'_n B'_n}_{A'_n} f_{a_n}^{+ C_n}(x_n) \cdots f_{a_1}^{+ C_1}(x_1) f_a^{+ C}(x) f_{a' C'}(y) f_{a'_1 C'_1}(y_1) \cdots f_{a'_n C'_n}(y_n) \end{aligned} \quad (61)$$

The cross $+$ denotes Dirac conjugation and the equation separates since we are dealing with a product state.

$\Phi_{\omega_{2n+2}}$ denotes the n -point distribution of the basic field Φ and the r.h.s of Eqn. (61) defines $\|k(f)\Psi\|^2$. All n -point distributions above arise from quasifree states. Hence they decay into the sum of products of two-point distributions. Using Corollary 3.24 and 3.27 one sees that arbitrary (finite) products of these distributions with each other exist. This ensures the finiteness of Eqn. (61). Now suppose we replace one or more $k(f_i)$ in Eqn. (61) by $\Phi(g_1, \dots, g_k, \tilde{g}_1, \dots, \tilde{g}_l, h_1, \dots, h_m) \equiv A(g_1) \cdots A(g_k) B(\tilde{g}_1) \cdots B(\tilde{g}_l) \Psi(h_1) \cdots \Psi(h_m)$ with $g_1, \dots, \tilde{g}_l \in C_0^\infty(M)$ and $h_1, \dots, h_m \in \Gamma_0(D^*M)$. Redoing the calculation in Eqn. (61) one ends up again with various products of two-point distributions. However all two-point distributions which include parts of the “ Φ ’s” are in fact smooth functions (for example ${}^A \omega_2(g_i, x) \in C^\infty(M)$; ${}^A \omega_2(g_i, g_j) \in \mathbb{C}$). Thus replacing an operator $k(f_i)$ by $\Phi(g_1, \dots, h_m)$ does not change the finiteness of the whole equation. Moreover since the products in Eqn. (61) are again *distributions*, it follows that $k_{\Psi\Psi'} : \Gamma(D^*M \otimes T^*M) \ni f \mapsto \langle \Psi \mid k(f)\Psi' \rangle$ is well defined and continuous for all $\Psi, \Psi' \in D$. This proves the assertion. Note that the relation between k and its adjoint k^* , i.e. $k^*(f) = k^+(f^+)$, follows by Corollary 3.18 from the corresponding properties of the basic fields.

2. is obvious since $\mathcal{H}$ is the GNS Hilbertspace of the basic fields, i.e., the “vacuum” Ω is cyclic for A, B and Ψ .

3. follows directly from our definition of D .
4. Note that A, B and Ψ mutually commute,

$$[A(f), A(g)]_- = [B(f), B(g)]_- = 0 \quad \text{if } \text{supp}(f) \subset \text{supp}(g)^c,$$

whereas two Ψ 's anti-commute for causally disjoint arguments. $\text{supp}(g)^c$ denotes the causal complement of the support of g . Therefore

$$\{k(f), k(g)\}_+ = 0 \quad \text{if } \text{supp}(f) \subset \text{supp}(g)^c,$$

□

REMARK .

- D' is not dense in H , since for example $\Psi = A(f)|\Omega\rangle$ can not be approximated by elements in D' : $\forall \Phi \in D'$ we have $\|\Psi - \Phi\|^2 = \|\Psi\|^2 + \|\Phi\|^2$ since $\langle \Psi, \Phi \rangle = 0$.
- $|\Omega\rangle$ is not cyclic for the fields j^{aA} .

It is worth noting that all n -point distributions of k^{aA} do satisfy our new GWFSSC in a non trivial way. The proof of this statement which is completely analogous to the considerations of Lemma 3.32 is omitted.

COROLLARY 4.2. *Let a spacelike hypersurface $\mathcal{C}$ and $f, g \in \Gamma(D^*M \otimes T^*M)$ be given. It follows that the wave front set of the distribution $(f, g) \mapsto \langle \Omega | j^+(f^+) j(g) \Omega \rangle$ does contain directions normal to $\mathcal{C} \times \mathcal{C}$, i.e., it violates the assumption of Theorem 3.17.*

PROOF. Consider $\langle \Omega | j^+(g^+) j(f) \Omega \rangle$. By Eqn. (61) this expression contains sums of the product of two 2-point distributions, hence the wave front set of this distribution is contained (and in fact equals) the set

$$W \equiv (\text{WF}(\omega^2) \oplus \text{WF}(\omega^2)) \cup \text{WF}(\omega^2).$$

However, due to the fact that in the first term the wave front sets are added, W contains –associated to the first entrance– directions in the whole forward lightcone, that is it contains all timelike direction. On the other hand the normals of $\mathcal{C}$ are timelike by definition, which proves the Corollary. □

REMARK . Unfortunately the assumptions of Theorem 3.17 are only sufficient for the existence of a restriction. Therefore we may not use the results of Corollary 4.2 to conclude that the restriction of j^{aA} to a spacelike hypersurface does not exist. On the other hand it is well known that on Minkowski space the product of free fields can not be restricted to spacelike hypersurfaces. Thus one might conjecture that a similar statement holds on a manifold.

2. The ‘supercharge’

We saw above that we can not apply Theorem 3.17 to restrict the ‘supercurrent’ to a spacelike hypersurface. However, we will show below that the (anti) commutator of the ‘supercurrent’ with a basic field gives an operator valued distribution which can be restricted to any spacelike hypersurface $\mathcal{C}$. Hence it is possible to define terms like $[Q_A^+, A(x)]_- \upharpoonright_{\mathcal{C}}$ rigorously. However, for this definition we need some more notation which is related to the splitting of the spacetime into space and time (See [Wal84, pp. 255]).

2.1. Splitting spacetime into space and time. Let (M, g_{ab}) be a Lorentz manifold and let $\mathcal{C}$ denote a smooth spacelike hypersurface in M with unit normal covector field $\tilde{n}_a$. Then there exists a covector field n_a near $\mathcal{C}$ on M , such that n_a restricted to $\mathcal{C}$ is $\tilde{n}_a$ and such that the integral curves of $n^a = g^{ab}n_b$ are (timelike) geodesics.

DEFINITION 4.3. The spacetime metric g_{ab} induces a spatial, i.e., a three dimensional, Riemannian² metric on $\mathcal{C}$ via the formula

$$h_{ab} := g_{ab} - n_a n_b.$$

The Levi-Civita connection D_a associated to this spatial metric is given by

LEMMA 4.4 (Lemma 10.2.1 of [Wal84]). *The Levi-Civita connection of h_{ab} is given by*

$$D_c T^{a_1 \dots a_l}_{b_1 \dots b_m} = h^{a_1}_{d_1} \dots h^{a_l}_{d_l} h^{b_m}_{e_m} h_c^f \nabla_f T^{d_1 \dots d_l}_{e_1 \dots e_m}, \quad (62)$$

where ∇_a is the Levi-Civita connection of g_{ab} .

DEFINITION 4.5. The extrinsic curvature of $\mathcal{C}$ in M , K_{ab} , is defined by

$$K_{ab} = h_a^c \nabla_c n_b.$$

²Note that the signature of h_{ab} is $(-, -, -)$.

Note that K_{ab} is *symmetric* in its indices due to the fact that it is half the Lie derivative of h_{ab} with respect to the vector field n (See [Wal84]). Moreover, since we assumed n_a to be a geodesic covector field, we have

$$K_{ab} = \nabla_a n_b \quad \text{if } n_a \text{ is geodesic,} \quad (63)$$

since $n_a n^c \nabla_c n_b$ vanishes by assumption. Another useful identity is $\nabla_a(v^a) = D_a v^a + n^a n_b \nabla_a v^b$ which is valid for all vector-fields tangential to $\mathcal{C}$.

DEFINITION 4.6. The trace of K_{ab} , $K^a{}_a$, is denoted by K –the extrinsic curvature scalar– and a hypersurface with $K = 0$ is called *maximal*.

REMARK .

- It is not known whether an arbitrary spacetime admits maximal hypersurfaces *globally*, since there are various constraints. For example the extrinsic curvature and the full Ricci tensor are connected by the Gauss-Codazzi relations. For a special class of spacetimes this difficult problem is investigated in [CW94]. Fortunately the situation in this work is far less complicated, since we need maximal spacelike hypersurfaces *locally* only, i.e., for every point p in the spacetime there should exist a neighborhood U and a spacelike hypersurface $\mathcal{C}$ such that $U \cap \mathcal{C}$ has extrinsic curvature zero. Using the following line of argument together with a result of Bartnik [Bar84], it should be possible to show that *all* spacetimes admit maximal hypersurfaces locally through every point. Bartnik shows in [Bar84] (Theorem 5.4) that all asymptotically flat spacetimes satisfying an interior condition possess maximal spacelike hypersurfaces globally. It is worth noting that his interior condition demands certain quantities which vanish on Minkowski space (equipped with the usual time function) to be uniformly bounded throughout the spacetime, hence a spacetime which differ only slightly from Minkowski space, e.g., on a compact set, does satisfy his interior condition automatically. Consider now a point p in our spacetime with three sufficiently small open neighborhoods U , U' and U'' , such that the closure $\bar{U}$ of U is contained in U' , $\bar{U}' \subset U''$ and U'' can be covered by a single chart. It follows that there exists a deformed spacetime (M', g'_{ab}) together with a global time function, such that M' is contained in $U'' \subset M$, g' equals g on U and g' is flat outside of U' . Note that (M', g'_{ab}) can be constructed using a chart ϱ to pull back the metric g to $\mathbb{R}^4$. Then one deforms this metric in the region $\varrho(U' \setminus \bar{U})$ to the Minkowski metric. The time function t is chosen, such that it is the usual time Minkowski time function outside of $\varrho(U'')$ and some arbitrary time function inside $\varrho(U)$. We interpolate t between $\varrho(U)$ and $\varrho(U')$ such that $\nabla_a t$ is nonzero and everywhere timelike. Finally, using a conformal mapping, which does not change the metric inside $\varrho(U)$, the resulting spacetime is mapped into $\varrho(U'')$. Then (M', g'_{ab}) together with t satisfy the interior condition of Bartnik's Theorem 5.4, which ensures that a maximal spacelike hypersurface $\mathcal{C}$ in $(M', g'_{ab}) \xrightarrow{i} (M, g_{ab})$ passing through p exists. Thus $U \cap \mathcal{C}$ is a spacelike hypersurface with extrinsic curvature zero in (M, g_{ab}) .
- It is worth noting that for a geodesic covector field n_a , which is in addition a Killing covector field, $K_{ab} = 0$ automatically.

2.2. (Anti-) commutators for the ‘supercharge’. In this subsection we will explicitly calculate the (anti-) commutators of the ‘supercharge’ with the basic fields and their derivatives. The former is obtained by smearing the distributional ‘supercurrent’ with a suitable testdensity. We remind the reader that we define distributions according to the notation of Hörmander loc.cit., i.e., distributions are extensions of smooth functions. It follows that the former correspond to the continuous linear forms on the smooth *densities* with compact support (See [Hör83, p. 145]). It is worth noting that the latter can be identified on every spacetime with smooth functions with compact support by using the natural volume element on this Lorentz manifold.

DEFINITION 4.7. Let k^{aA} denote the ‘supercurrent’, $\mathcal{C}$ a spacelike hypersurface, $n^a \in \Gamma(TM)$ a unit timelike geodesic vector field which, when restricted to $\mathcal{C}$ is normal to this surface. Fix a point $x_0 \in \mathcal{C} \subset M$ and let χ be a smooth function with compact support in a normal neighborhood of x_0 , which can be identified with the density $\chi d\mu$, where $d\mu$ denotes the natural volume element on M . The *regularized ‘supercharge’* ${}^{\chi d\mu}Q^{A'}(x_0)$ is defined by

$${}^{\chi d\mu}Q^{A'}(x_0) := \int_M k^{aA}(x) \mathcal{J}_A{}^{A'}(x, x_0) n_a(x) \chi(x) d\mu_x. \quad (64)$$

Note that ${}^{\chi d\mu}Q^{A'}(x_0)$ as well as its Dirac adjoint ${}^{\chi d\mu}Q_{A'}^+(x_0)$ are well defined operators on the GNS Hilbertspace which transform as a spinor (cospinors respectively) at the point x_0 .

PROPOSITION 4.8. *Let $\mathcal{C}$, n^a and x_0 be the same as in Definition 4.7. Then the six operator valued distributions*

$$\begin{aligned} D'(M \times M) \ni {}^A V_{x_0} : (\chi d\mu, h) &\mapsto [\chi^{d\mu} Q_{A'}^+(x_0), A(h)]_- \\ D'(M \times M) \ni {}^B V_{x_0} : (\chi d\mu, h) &\mapsto [\chi^{d\mu} Q_{A'}^+(x_0), B(h)]_- \\ D'(M \times DM) \ni {}^\Psi V_{x_0} : (\chi d\mu, h_B) &\mapsto \{\Psi^B(h_B), \chi^{d\mu f} Q_{A'}^+(x_0)\}_+ \\ D'(M \times T^*M) \ni \nabla_a {}^A V_{x_0} : (\chi d\mu, g^a) &\mapsto [\chi^{d\mu} Q_{A'}^+(x_0), \nabla_a A(g^a)]_- \\ D'(M \times T^*M) \ni \nabla_a {}^B V_{x_0} : (\chi d\mu, g^a) &\mapsto [\chi^{d\mu} Q_{A'}^+(x_0), \nabla_a B(g^a)]_- \\ D'(M \times DM \otimes T^*M) \ni \nabla_a {}^\Psi V_{x_0} : (\chi d\mu, g_B^a) &\mapsto \{\nabla_a \Psi^B(g_B^a), \chi^{d\mu} Q_{A'}^+(x_0)\}_+ \end{aligned}$$

have unique restrictions to the spacelike hypersurface $\mathcal{C}$.

PROOF. Let (H, Ω, A, B, Ψ) denote the GNS-Hilbertspace of a globally Hadamard product-state ω and let D be the usual dense invariant domain for the three fields. We prove first of all that the distribution

$$< {}^A V_{x_0}(\cdot, \cdot) \Omega | {}^A V_{x_0}(\cdot, \cdot) \Omega > \in D'(M^4)$$

on M^4 can be restricted to $\mathcal{C}^4$. Inserting the definitions we obtain:

$$\begin{aligned} & \| {}^A V_{x_0}(\chi d\mu, h d\mu) \Omega \|^2 \\ &= \int_{M^4} d\mu_x d\mu_y d\mu_z d\mu_v \\ & \quad < \Omega | -i[k^{+a}{}_A(x) \mathcal{J}^A_Z(x, x_0) n_a(x) \chi(x), A(y)]_- \cdot \\ & \quad i[k^{a''B''}{}_{(z)} \mathcal{J}_{B''}{}^Y(z, x_0) n_{a''}(z) \chi(z), A(v)]_- h(v) \Omega > \\ &= \int_{M^4} d\mu_x d\mu_y d\mu_z d\mu_v < \Omega | \Psi^+_A(x) \Psi^{A''}(z) \Omega > \gamma^{aA}{}_B \\ & \quad \left(\left(-i \overline{\nabla}_x - m \right)_C^B < \Omega | [A(x), A(y)]_- \Omega > \left(i \overline{\nabla}_z - m \right)_{C''}^{B''} < \Omega | [A(z), A(v)]_- \Omega > \right. \\ & \quad - \left((-i \overline{\nabla}_x - m) \gamma^5 \right)_C^B < \Omega | [B(x), B(y)]_- \Omega > \\ & \quad \left. \left((i \overline{\nabla}_z - m) \gamma^5 \right)_{C''}^{B''} < \Omega | [B(z), B(v)]_- \Omega > \right) \\ & \quad \mathcal{J}^C_Z(x, x_0) n_a \chi(x) \gamma^{a''C''}{}_{A''} \mathcal{J}_{B''}{}^Y(z, x_0) n_{a''}(z) \chi(v) \\ &= \int_{M^4} d\mu_x d\mu_y d\mu_z d\mu_v (-1)^\Psi \omega^{2A'}(x, z) \gamma^{aA}{}_B \\ & \quad \left(\left(-i \overline{\nabla}_x - m \right)_C^B E(x, y) \left(i \overline{\nabla}_z - m \right)_{C''}^{B''} E(z, v) \right. \\ & \quad - \left((-i \overline{\nabla}_x - m) \gamma^5 \right)_C^B E(x, y) \left((i \overline{\nabla}_z - m) \gamma^5 \right)_{C''}^{B''} E(z, v) \Big) \\ & \quad \mathcal{J}^C_Z(x, x_0) n_a \chi(x) \gamma^{a''C''}{}_{A''} \mathcal{J}_{B''}{}^Y(z, x_0) n_{a''}(z) \chi(v) \end{aligned} \tag{65}$$

We used the fact that Ω is a quasifree product state and introduced the following abbreviations for the vacuum expectation values:

$$\begin{aligned} iE(x, y) &:= < \Omega | [A(x), A(y)]_- \Omega > \\ iE(x, y) &:= < \Omega | [B(x), B(y)]_- \Omega > \\ \Psi \omega^{2A'}(x, y) &:= < \Omega | \Psi^+_A(x) \Psi^{A'}(y) \Omega > \end{aligned}$$

Note that $< \Omega | [A(x), A(y)]_- [A(z), A(v)]_- \Omega > = iE(x, y) iE(z, v)$, since we are dealing with free fields. For our claim it is sufficient to show that the following product of distributions

$$\tilde{V} \equiv (E(y, x) \otimes \mathbf{1} \otimes \mathbf{1}) \cdot (\mathbf{1} \otimes \Psi \omega^{2A'}(x, z) \otimes \mathbf{1}) \cdot (\mathbf{1} \otimes \mathbf{1} \otimes E(z, v))$$

can be restricted to $\mathcal{C}$, since (65) consists of terms of this type together with derivatives of it, multiplied by smooth functions, only. For the verification of this statement we calculate the wave front set of the distribution $\tilde{V}$. Note that for a Distribution $u \in D'(M)$ with wave front set $\text{WF}(u)$ the wave front set of $\mathbf{1} \otimes u$ is contained in the set $(M, 0) \times \text{WF}(u)$ (See for example [Hör71] p. 130) and recall that differentiation does not enlarge the wave front set. For the wave front sets of E , which is the difference of the advanced and retarded fundamental solutions of the Klein Gordon equation (see below) one finds (see for example [Hör71]):

$$\text{WF}(E) = \{(x, \xi_1, y, \xi_2) \in T^*M \times T^*M \setminus \{0\} | (x, \xi_1) \sim (y, \xi_2); \xi_1^2 = 0\}, \quad (66)$$

whereas for the wave front set of $\Psi \omega^{2^A}_{A'}$ we already found in Corollary 3.26

$$\text{WF}(\Psi \omega^{2^A}_{A'}) \subseteq \{(x, \xi_1, y, \xi_2) \in T^*M \times T^*M \setminus \{0\} | (x, \xi_1) \sim (y, \xi_2); \xi_1^2 = 0; \xi_1^0 > 0\}. \quad (67)$$

By Theorem 3.16 the wavefront set of $\tilde{V}$ and all its derivatives is contained in the following set:

$$\begin{aligned} \text{WF}(\tilde{V}) &\subseteq S \\ &:= \underbrace{\text{WF}(E \otimes \mathbf{1} \otimes \mathbf{1})}_{=: \Gamma_1} + \underbrace{\text{WF}(\mathbf{1} \otimes E \otimes \mathbf{1})}_{=: \Gamma_2} + \underbrace{\text{WF}(\mathbf{1} \otimes \mathbf{1} \otimes E)}_{=: \Gamma_3} \\ &\cup \Gamma_1 \cup \Gamma_2 \cup \Gamma_3 \cup (\Gamma_1 \oplus \Gamma_2) \cup (\Gamma_1 \oplus \Gamma_3) \cup (\Gamma_2 \oplus \Gamma_3) \\ &\subseteq \{(y, \xi_1, x, -\xi'_1, z, 0, v, 0) \in T^*M^4 \setminus \{0\} | (y, \xi_1) \sim (x, \xi'_1); \xi_1^2 = 0\} \\ &\cup \{(y, 0, x, \xi_2, z, -\xi'_2, v, 0) \in T^*M^4 \setminus \{0\} | (x, \xi_2) \sim (z, \xi'_2); \xi_2^2 = 0; \xi_2^0 > 0\} \\ &\cup \{(y, 0, x, 0, z, \xi_3, v, -\xi'_3) \in T^*M^4 \setminus \{0\} | (z, \xi_3) \sim (v, \xi'_3); \xi_1^2 = 0\} \\ &\cup \{(y, \xi_1, x, \xi_2 - \xi'_1, z, -\xi'_2, v, 0) \in T^*M^4 \setminus \{0\} | (y, \xi_1) \sim (x, \xi'_1); (x, \xi_2) \sim (z, \xi'_2); \\ &\quad \xi_1^2 = \xi_2^2 = 0; \xi_2^0 > 0\} \\ &\cup \{(y, \xi_1, x, -\xi'_1, z, \xi'_3, v, -\xi_3) \in T^*M^4 \setminus \{0\} | (y, \xi_1) \sim (x, \xi'_1); (z, \xi_3) \sim (v, \xi'_3); \\ &\quad \xi_1^2 = \xi_3^2 = 0\} \\ &\cup \{(y, 0, x, \xi_2, z, \xi_3 - \xi'_2, v, -\xi'_3) \in T^*M^4 \setminus \{0\} | (x, \xi_2) \sim (z, \xi'_2); (z, \xi_3) \sim (v, \xi'_3); \\ &\quad \xi_2^2 = \xi_3^2 = 0; \xi_2^0 > 0\} \\ &\cup \{(y, \xi_1, x, \xi_2 - \xi'_1, z, \xi_3 - \xi'_2, v, -\xi'_3) \in T^*M^4 \setminus \{0\} | \\ &\quad (y, \xi_1) \sim (x, \xi'_1); (x, \xi_2) \sim (z, \xi'_2); (z, \xi_3) \sim (v, \xi'_3); \\ &\quad \xi_1^2 = \xi_2^2 = \xi_3^2 = 0; \xi_2^0 > 0\}, \end{aligned}$$

where we used the fact that the wave front set of $\Psi \omega^{2^A}_{A'}$ is contained in that of E . Now let $\tilde{\varphi}$ denote the natural embedding of $\mathcal{C}^4$ into M^4 and $N_{\tilde{\varphi}}$ the associated set of normals of $\tilde{\varphi}$. Then the two sets S and $N_{\tilde{\varphi}}$ do not intersect, hence the restriction of $\tilde{V}$ as well as that of $\|{}^A V_{x_0}(\cdot, \cdot)\Omega\|$ to $\mathcal{C}^4$ exists by Theorem 3.17. Now replace the vector Ω in Eqn. (65) by an arbitrary element $\Phi \in D \subset H$. Since Ω is quasifree one ends up with a sum of terms which are tensor products of $\tilde{V}$ and its derivatives with two-point distributions of the basic fields or fundamental solutions. Since the latter can be restricted³ to $\mathcal{C} \times \mathcal{C}$, it follows that the tensor products mentioned above can be restricted to $\mathcal{C} \times \mathcal{C}$, too. Especially the restriction to $\mathcal{C}$ for the internal variables, i.e., the variables associated to ${}^A V_{x_0}$, exists, which proves the claim for the first operator valued distribution ${}^A V_{x_0}$. The five remaining cases follow completely analogous. The corresponding proofs are omitted in order to avoid repeating the argument verbatim. $\square$

We have the following Theorem.

THEOREM 4.9. *Using the same notation as in Proposition 4.8: For all smooth testdensities on $\mathcal{C}$ with support sufficiently close to x_0 , the resulting six distributions on $\mathcal{C} \times \mathcal{C}$ are independent of the density $\chi d\mathcal{C}$, provided $\chi \equiv 1$ in a neighborhood of the support of the second testdensity. $d\mathcal{C}$ denotes the induced volume form on $\mathcal{C}$. More precisely: Let $\tilde{\varphi}$*

³Recall that their wave front sets contain lightlike singular directions only.

and φ denote the embedding of $\mathcal{C} \times \mathcal{C}$ and $\mathcal{C}$ into $M \times M$ and M respectively. Then

$$\begin{aligned}
D'(\mathcal{C}) \ni \tilde{\varphi}^*({}^A V_{x_0})(\chi d\mathcal{C}, \cdot): \mathbf{h} &\mapsto [{}^C G_{A''}^+(x_0), A(\mathbf{h})]_- \\
&:= [{}^{\chi d\mathcal{C}} Q_{A''}^+(x_0), A(\mathbf{h})]_- \\
&= \varphi^* \left(+i \Psi_A^+ \mathcal{J}_{A''}^A(x_0) \right) (\mathbf{h}) \\
D'(\mathcal{C}) \ni \tilde{\varphi}^*({}^B V_{x_0})(\chi d\mathcal{C}, \cdot): \mathbf{h} &\mapsto [{}^C G_{A''}^+(x_0), B(\mathbf{h})]_- \\
&:= [{}^{\chi d\mathcal{C}} Q_{A''}^+(x_0), B(\mathbf{h})]_- \\
&= \varphi^* \left(+(\Psi^+ \gamma^5)_A \mathcal{J}_{A''}^A(x_0) \right) (\mathbf{h}) \\
D'(D_C M) \ni \tilde{\varphi}^*({}^\Psi V_{x_0})(\chi d\mathcal{C}, \cdot): \mathbf{h}_B &\mapsto \{ \Psi^B(\mathbf{h}_B), {}^C G_{A''}^+(x_0) \}_+ \\
&:= \{ \Psi^B(\mathbf{h}_B), {}^{\chi d\mathcal{C}} Q_{A''}^+(x_0) \}_+ \\
&= \varphi^* \left(+i \left(((i\nabla + m)(A - i\gamma^5 B)) \mathcal{J}(x_0) \right)^B_{A''} \right) (\mathbf{h}_B) \\
D'(T^* \mathcal{C}) \ni \tilde{\varphi}^*(\nabla_a {}^A V_{x_0})(\chi d\mathcal{C}, \cdot): \mathbf{g}^a &\mapsto [{}^C G_{A''}^+(x_0), \nabla_a A(\mathbf{g}^a)]_- \\
&:= [{}^{\chi d\mathcal{C}} Q_{A''}^+(x_0), \nabla_a A(\mathbf{g}^a)]_- \\
&= \varphi^* \left(i D_a(\Psi^+ \mathcal{J})_{A''} + i n_a D_b(\Psi^+ \not{p} \gamma^b \mathcal{J})_{A''} \right. \\
&\quad \left. - m n_a(\Psi^+ \not{p} \mathcal{J})_{A''} \right) (\mathbf{g}^a) \\
D'(T^* \mathcal{C}) \ni \tilde{\varphi}^*(\nabla_a {}^B V_{x_0})(\chi d\mathcal{C}, \cdot): \mathbf{g}^a &\mapsto [{}^C G_{A''}^+(x_0), \nabla_a B(\mathbf{g}^a)]_- \\
&:= [{}^{\chi d\mathcal{C}} Q_{A''}^+(x_0), \nabla_a B(\mathbf{g}^a)]_- \\
&= -i \varphi^* \left(i D_a(\Psi^+ \gamma^5 \mathcal{J})_{A''} + i n_a D_b(\Psi^+ \not{p} \gamma^b \gamma^5 \mathcal{J})_{A''} \right. \\
&\quad \left. - m n_a(\Psi^+ \not{p} \gamma^5 \mathcal{J})_{A''} \right) (\mathbf{g}^a) \\
D'(D_C M \otimes T^* \mathcal{C}) \ni \tilde{\varphi}^*(\nabla_a {}^\Psi V_{x_0})(\chi d\mathcal{C}, \cdot): \mathbf{g}_B^a &\mapsto \{ \nabla_a \Psi^B(\mathbf{g}_B^a), {}^C G_{A''}^+(x_0) \}_+ \\
&:= \{ \nabla_a \Psi^B(\mathbf{g}_B^a), {}^{\chi d\mathcal{C}} Q_{A''}^+(x_0) \}_+ \\
&= \varphi^* \left(-i h_a {}^c \nabla_c \left(((-i\nabla - m)(A - i\gamma^5 B)) \mathcal{J} \right)^B_{A''} \right. \\
&\quad \left. - i n_a \gamma^b h_b {}^c \nabla_c (\not{p}((-i\nabla - m)(A - i\gamma^5 B)) \mathcal{J})^B_{A''} \right. \\
&\quad \left. - m n_a(\not{p}((-i\nabla - m)(A - i\gamma^5 B)) \mathcal{J})^B_{A''} \right) (\mathbf{g}_B^a),
\end{aligned}$$

provided $\chi \equiv 1$ on the support of the testdensities. Here $D_C M$ denotes all elements in DM with basepoint in $\mathcal{C}$.

Note that we used boldface type, e.g., χ and $\mathbf{h}$, to indicate smooth functions and densities respectively on $\mathcal{C}$. The notion ${}^C G_{A''}^+(x_0)$ was introduced to emphasize that the result does not depend on χ . It is obvious that the six operator valued distributions in Theorem 4.9 smoothly depend on x_0 , hence they have a natural extension to operator valued distributions on a neighborhood of $U \subset \mathcal{C} \times \mathcal{C}$ of the diagonal in $\mathcal{C} \times \mathcal{C}$.

COROLLARY 4.10. *The six distributions of Theorem 4.9, when extended to $U \subset \mathcal{C} \times \mathcal{C}$ can be restricted to the diagonal in $\mathcal{C} \times \mathcal{C}$, i.e., they uniquely define six operator valued distributions on $\mathcal{C}$. For the corresponding kernels one*

finds:

$$\begin{aligned}
[{}^c G_A^+, A(x)]_- &= \lim_{x_0 \rightarrow x} [\mathfrak{x} Q_{A'}^+(x_0), A(x)]_- = +i\Psi_A^+(x) \\
[{}^c G_A^+, B(x)]_- &= \lim_{x_0 \rightarrow x} [\mathfrak{x} Q_{A'}^+(x_0), B(x)]_- = (\Psi^+ \gamma^5)_A(x) \\
\{\Psi^B(x), {}^c G_A^+\}_+ &= \lim_{x_0 \rightarrow x} \{\Psi^B(x), \mathfrak{x} Q_{A'}^+(x_0)\}_+ = +i((i\nabla + m)(A - i\gamma^5 B))^B_A \\
[{}^c G_A^+, \nabla_a A(x)]_- &= \lim_{x_0 \rightarrow x} [\mathfrak{x} Q_{A'}^+(x_0), \nabla_a A(x)]_- \\
&= ih_a^b \nabla_b \Psi_A^+ + in_a h_b^c \nabla_c (\Psi^+ \not{p} \gamma^b)_A \\
&\quad - mn_a (\Psi^+ \not{p})_A \\
[{}^c G_A^+, \nabla_a B]_- &= \lim_{x_0 \rightarrow x} [\mathfrak{x} Q_{A'}^+(x_0), \nabla_a B(x)]_- \\
&= h_a^b \nabla_b (\Psi^+ \gamma^5)_A + n_a h_b^c \nabla_c (\Psi^+ \not{p} \gamma^b \gamma^5)_A \\
&\quad - mn_a (\Psi^+ \not{p} \gamma^5)_A \\
\{\nabla_a \Psi^B(x), {}^c G_A^+\}_+ &= \lim_{x_0 \rightarrow x} \{\nabla_a \Psi^B(x), \mathfrak{x}^f Q_{A'}^+(x_0)\}_+ \\
&= -ih_a^b \nabla_b ((-i\nabla - m)(A - i\gamma^5 B))^B_A \\
&\quad - in_a \{\gamma^d h_d^c \nabla_c (\not{p}(-i\nabla - m)(A - i\gamma^5 B))\}^B_A \\
&\quad - mn_a (\not{p}(-i\nabla - m)(A - i\gamma^5 B))^B_A.
\end{aligned}$$

Corollary 4.10 is obvious, since we may use the fact that the pointwise product of a smooth function $g \in C^\infty(M)$ with a section in the spinorbundle $h^A \in \Gamma(DM)$, $g(x)h^A(x)$, is again a section in the spinorbundle. Similarly the pointwise product of $g_A \in \Gamma(D^*M)$ with $h^A \in \Gamma(DM)$, $g_A(x)h^A(x)$, defines a smooth function on M by dual pairing and the claim follows immediately using Corollary 3.19.

REMARK . The kernel of ${}^c Q_{A'}^+$ is not covariantly constant. Thus the results of Proposition 4.8 and Theorem 4.9 depend on the hypersurface $\mathcal{C}$. On the other hand the results of Corollary 4.10 –as long as no derivatives occur– have a natural extension to all of M independent of $\mathcal{C}$.

For the proof of the Theorem we need some properties of the fundamental solutions of the Klein-Gordon and Dirac equation respectively. We begin with the scalar case.

2.3. The fundamental solution of the Klein Gordon equation. In this subsection some useful identities for the fundamental solution of the Klein Gordon equation are established. Let (M, g_{ab}) denote a globally hyperbolic Lorentz manifold with Cauchy surface $\mathcal{C}$. For the sake of simplicity we identify in the sequel of this work test densities and test functions using the natural measure $d\mu = \sqrt{|g_{ab}|}d^4x$ on M and denote the space of testdensities on M with compact support by $D(M)$. Let P be a pseudodifferential operator on M which has the form $P = \square + a$, where $\square$ denotes the D'Alembert operator associated to g_{ab} and a is some smooth function on M . The ‘squared’ Dirac operator and the Klein Gordon operator are examples for P . As a consequence of the global hyperbolicity, there exist global fundamental solutions for P uniquely determined by their support properties (see Leray [Ler63], Choquet-Bruhat [CB77]). That is there exist unique distributions $E^\pm \in D'(M \times M)$, viewed as linear mappings $\Delta^\pm: C_0^\infty(M) \rightarrow C^\infty(M)$ by the identification

$$\int_M f_1 \Delta^\pm(f_2) d\mu = E^\pm(f_1 d\mu \otimes f_2 d\mu) \quad \text{for all } f_1, f_2 \in C_0^\infty(M),$$

which are called advanced and retarded fundamental solutions respectively. They are uniquely determined by the following two properties:

1.

$$P\Delta^\pm = \Delta^\pm P = \mathbf{1}, \tag{68}$$

where $\mathbf{1}$ denotes the identity on $C_0^\infty(M)$.

2. If $f \in C_0^\infty(M)$ then $\text{supp } \Delta^\pm(f)$ is compact with respect to the future (+) or past (−).

Here, a set $U \subset M$ is called compact with respect to the future (past) if and only if for any point $x \in M$, $J^+(x) \cap J^-(U)$ ($J^-(x) \cap J^+(U)$) is compact. Recall that $J^+(x)$ ($J^-(x)$) denotes the set of all points $y \in M$ which can be connected

to x with a future (past) directed causal curve from x to y . We define the fundamental solutions $E = E^+ - E^-$ and $\Delta = \Delta^+ - \Delta^-$. Given Δ it is easy to solve the Cauchy problem for the differential equation $Pu = 0$, $u \in C_0^\infty(M)$. The result is:

PROPOSITION 4.11 (Huygens principle (smooth form)). *Let ρ_0 (ρ_1) denote the restriction (normal restriction) of any smooth function on M to the Cauchy surface $\mathcal{C}$. Then for any smooth solution u of $Pu = 0$ on M and any $f \in C_0^\infty(M)$*

$$\int_M u f \, d\mu = \int_{\mathcal{C}} u_0 \rho_1(\Delta f) - u_1 \rho_0(\Delta f) \, d\mathcal{C}, \quad (69)$$

where $u_0 \equiv \rho_0(u) := u|_{\mathcal{C}}$, $u_1 \equiv \rho_1(u) := (n_\mu \nabla^\mu u)|_{\mathcal{C}}$ and $d\mathcal{C}$ denotes the natural volume element on $\mathcal{C}$ determined by the induced metric h_{ab} (compare subsection 2.1 of this chapter).

PROOF. For the proof, which is merely an application of Greens formula, the reader is referred to Dimock's paper [Dim80] Lemma A.1. $\square$

Now let u be a distributional solution of $Pu = 0$. Then u admits a restriction to $\mathcal{C}$ according to Theorem 3.17, since by Theorem 3.21 its wave front set contains at most lightlike directions. Thus Eqn. (69) can be extended by continuity and we proved

PROPOSITION 4.12. *Let $\hat{\varphi}: (M \times \mathcal{C}) \ni (p_1, p_2) \mapsto (p_1, p_2) \in (M \times M)$ be the natural embedding of $M \times \mathcal{C}$ into $M \times M$. Then the following identity holds for all $f \in D(M)$:*

$$\begin{aligned} u(f) &= \hat{\varphi}^* \{ (1 \otimes u) \cdot ((1 \otimes n_a \nabla^a) E) \} (f \otimes 1 d\mathcal{C}) \\ &\quad - \hat{\varphi}^* \{ (1 \otimes n_a \nabla^a u) \cdot E \} (f \otimes 1 d\mathcal{C}), \\ &\quad \text{"} \equiv \text{"} \int_M \int_{\mathcal{C}} u(z) n_{a'} \nabla^{a'} E(x, z) f(x) - (n_{a'} \nabla^{a'} u(z)) E(x, z) f(x) d\mathcal{C}_z d\mu_x \end{aligned} \quad (70)$$

where $\cdot \otimes \cdot$ is the usual tensor product of distributions, 1 being the function 1 and 1 denotes the restriction of 1 to $\mathcal{C}$. Note that Eqn. (70) is well defined, since $(J^+(\text{supp}(f)) \cup J^+(\text{supp}(f))) \cap \mathcal{C}$ is compact.

It is worth noting that the fundamental solution E , being a distributional bisolution of the Klein Gordon equation, can be restricted to $\mathcal{C}$. For the restriction of E and its normal derivative one finds:

PROPOSITION 4.13. *Let E be the (unique) fundamental solution of the scalar Klein Gordon equation.*

- *The restriction of E to any Cauchy surface $\mathcal{C}$ vanishes, i.e., let $\tilde{\varphi}$ denote the embedding of $\mathcal{C} \times \mathcal{C}$ into $M \times M$, then*

$$\tilde{\varphi}^*(E) = 0. \quad (71)$$

- *The normal derivative of E with respect to the first entry restricted to $\mathcal{C} \times \mathcal{C}$ is minus the Dirac delta distribution, i.e.,*

$$\tilde{\varphi}^*(n^a \nabla_a E)(x, y) = -\delta(x, y). \quad (72)$$

REMARK . These two statements can equivalently be formulated in terms of the operator Δ . They state that Δ , when restricted to $\mathcal{C} \times \mathcal{C}$ vanishes, whereas the restriction of its normal derivative with respect to the first entry is minus the identity on $C_0^\infty(\mathcal{C})$.

PROOF. For $y \in \mathcal{C}$ let $u_y \in D'(M)$ be the unique distributional solution of $(\square + m^2)u_y = 0$ with initial data

$$(u_y)|_{\mathcal{C}} = 0 \quad (n^a \nabla_a u_y)|_{\mathcal{C}} = -\delta(\cdot, y),$$

where $\delta(\cdot, y)$ denotes Dirac delta distribution at $y \in \mathcal{C}$, i.e., for $g d\mathcal{C} \in D(\mathcal{C})$, $\delta(g d\mathcal{C}, y) = g(y)$. Inserting these initial datas in Eqn. (70) we obtain for all $f \in D(M)$ –using the notation of Proposition 4.12–

$$\begin{aligned} u_y(f) &= 0 - \hat{\varphi}^* \{ 1 \otimes n^a \nabla_a u_y \} \hat{\varphi}^* \{ E \} (f \otimes 1 d\mathcal{C}) \\ &= (1 \otimes \delta(\cdot, y)) \cdot \hat{\varphi}^* \{ E \} (f \otimes 1 d\mathcal{C}) \\ &= +\hat{\varphi}^* \{ E \} (f \otimes y) \\ &= E(f, y). \end{aligned}$$

This proves $u_y = E(\cdot, y)$ for $y \in \mathcal{C}$. Thus $E(\cdot, y)|_{\mathcal{C}} = 0$ together with $(n^a \nabla_a E(\cdot, y))|_{\mathcal{C}} = -\delta(\cdot, y)$ for all $y \in \mathcal{C}$, which was claimed in the Corollary. $\square$

Note that $E(f, y)$ is a smooth function on M for all $f \in C_0^\infty(M)$, since Δ applied to a smooth function with compact support is smooth. Hence $E|_{\mathcal{C} \times \mathcal{C}}(\cdot, y) \equiv E(\cdot, y)|_{\mathcal{C}}$ if $y \in \mathcal{C}$. The next Lemma states that derivatives of distributions in direction tangential to $\mathcal{C}$ commute with the restriction to $\mathcal{C}$.

LEMMA 4.14. *Let $\mathcal{C}$ be a spacelike hypersurface and let $u \in D'(M)$ be a distribution on M such that $\varphi^*(u)$, the restriction of u to $\mathcal{C}$, exists. Let v^a be a smooth vector field tangential to $\mathcal{C}$ and let h_{ab} , D_a denote the induced metric and Levi-Civita connection respectively on $\mathcal{C}$ (compare Definition 4.3 and Lemma 4.4). Then*

$$\varphi^*(v^a \nabla_a u) = v^a D_a \varphi^*(u).$$

PROOF. Suppose u is given by a smooth function, then

$$\begin{aligned} \varphi^*(v^a \nabla_a u)(f) &:= \int_{\mathcal{C}} (v^a \nabla_a u(x)) f(x) d\mathcal{C} \\ &= \int_{\mathcal{C}} (v^a D_a u(x)) f(x) d\mathcal{C}, \end{aligned}$$

since v^a is tangential on $\mathcal{C}$

$$= - \int_{\mathcal{C}} u(x) D_a (v_a f)(x) d\mathcal{C},$$

by Stokes Theorem, i.e., integration by parts, which is valid, since D_a is the induced Levi-Civita connection on $\mathcal{C}$

$$=: (v^a D_a \varphi^*(u))(f)$$

By continuity this extends to $D'(M)$. □

COROLLARY 4.15. *Let E be the fundamental solution of the Klein Gordon equation, v^a denote a smooth vector field and $\tilde{\varphi}$ is the embedding of $\mathcal{C} \times \mathcal{C}$ into $M \times M$. Then*

$$\tilde{\varphi}^*(v^a \nabla_a E) = -v^a n_a \delta.$$

PROOF. Split the vectorfield v^a in its normal and tangential components and use the linearity of $\tilde{\varphi}^*$.

$$\tilde{\varphi}^*(v^a \nabla_a E) = \tilde{\varphi}^*(v^a h_a^b D_b E) + \tilde{\varphi}^*(v^a n_a n^b \nabla_b E)$$

By Lemma 4.14 the first summand is equal to $v^a h_a^b D_b \tilde{\varphi}^*(E)$, which vanishes by Proposition 4.13. By Eqn. (72) the second summand is equal to $-v^a n_a \delta$, which finishes the proof. □

We now establish analogous properties for the fundamental solution of the Dirac equation.

2.4. The fundamental solution of the Dirac equation. Let (M, g_{ab}) be a globally hyperbolic Lorentz manifold with Majorana spinor (cospinor) bundle DM (D^*M). Let $\mathcal{C}$ be a Cauchy surface in M . We remind the reader that functions and densities on M and $\mathcal{C}$ can be identified using the appropriate metric. The Dirac operator $\not{D} := (i\not{\nabla} - m)$ fulfils the following important identity (See e.g. [BT87])

$$(i\not{\nabla} - m)(-i\not{\nabla} - m) = \left(\square + (m^2 + \frac{1}{4}R) \right) \mathbf{1},$$

where $\mathbf{1}$ denotes the identity on the spinor bundle. Note that $(\square + (m^2 + \frac{1}{4}R))$ is the SUSY coupled Klein Gordon operator, which is strictly hyperbolic in the sense of Leray. It follows that

$$\left(\square + (m^2 + \frac{1}{4}R) \right) u = 0$$

possesses unique advanced and retarded fundamental solutions viewed either as distributions in

$D'(D^*M \times DM)$, denoted by $E_A^{\pm A'}$, or as linear operators from $\Gamma_0(DM)$ to $\Gamma(DM)$, denoted by $\Delta_A^{\pm A'}$. Their difference will be written as $E_A^{A'} := E_A^{+A'} - E_A^{-A'}$ and $\Delta_A^{A'} := \Delta_A^{+A'} - \Delta_A^{-A'}$ respectively. The existence of a unique fundamental solution for the Dirac equation $\not{D}u = 0$ is a consequence of the following Theorem of Dimock [Dim82].

PROPOSITION 4.16.

- The operator $\not{D}$ on $\Gamma_0(DM)$ has a unique fundamental solutions

$$\Xi^{\pm} : \Gamma_0(DM) \rightarrow \Gamma(DM)$$

given by

$$\Xi^{\pm} := (-i\not{\nabla} - m)\Delta^{\pm}$$

- Its Dirac adjoint, $\not{D}^+$, which is an operator on $\Gamma_0(D^*M)$, has unique fundamental solutions

$$\hat{\Xi}^{\pm} : \Gamma_0(D^*M) \rightarrow \Gamma(D^*M)$$

given by

$$\hat{\Xi}^{\pm} := (i\not{\nabla} - m)\Delta^{\pm},$$

where $\Delta^\pm$ are the fundamental solutions of $(\square + (m^2 + \frac{1}{4}R)) \mathbf{1}$ uniquely determined by their support properties. $\mathbf{1}$ denotes the identity on the spinor (cospinor) bundle and the spinor indices have been suppressed.

We may identify $\Xi^\pm$ ($\hat{\Xi}^\pm$) with distributions denoted by $S^\pm \in D'(D^*M \times DM)$ ($\hat{S}^\pm \in D'(DM D^*M)$). Their difference is written as Ξ ($\hat{\Xi}$) and S ($\hat{S}$) respectively. For the formulation of Huygens principle it is necessary to define products of distributions in these bundles. Let us call $D'(DM)$ ($D'(D^*M)$) the space of spinor (cospinor) valued distributions. Using a similar notation as in the previous subsection, we denote the testdensities for $D'(DM)$ ($D'(D^*M)$) by $D(D^*M)$ ($D(DM)$). It is worth noting that the smooth sections in the corresponding bundles are dense in these spaces. Moreover, on a globally hyperbolic manifold things are greatly simplified by the fact that every spin structure SM over M necessarily is the trivial bundle [Ger68]. It follows that all calculation can be performed in 'coordinates': Pick a global spin frame $E: M \rightarrow SM$. We denote the corresponding moving frames by $(E_1, \dots, E_4)$ and their duals by $(E^1, \dots, E^4)$ respectively. Then every element $u^A \in D'(DM)$, $v_A \in D'(D^*M)$, $f^A \in D(D^*M)$, $g_A \in D(DM)$ can uniquely be written as

$$\begin{aligned} u &= \sum u^\Lambda E_\Lambda & v &= \sum v_\Lambda E^\Lambda \\ f &= \sum f^\Sigma E_\Sigma & g &= \sum g_\Sigma E^\Sigma, \end{aligned}$$

where u^Λ and v_Λ , $\Lambda = 1, \dots, 4$ are distributions on M and f^Σ , g_Σ , $\Sigma = 1, \dots, 4$ are densities with compact support. Note that we use an upper index for $f^A \in D(D^*M)$, since this density can be identified with a *spinor* field. A *distributional product* of u^A and v_A is well defined if and only if $\sum_\Lambda u^\Lambda \cdot v_\Lambda$ exists, where each term in the sum is the usual product of distributions. It is obvious that in the special case $u^A \in \Gamma(DM) \subset D'(DM)$ and $v_A \in \Gamma(D^*M) \subset D'(D^*M)$ this definition is frame independent. Due to the continuity of the product and since $\Gamma(DM)$ and $\Gamma(D^*M)$ are dense in $D'(DM)$ and $D'(D^*M)$ respectively, this independence can be lifted and $u^A \cdot v_A \in D'(M)$ defines a distribution on M if the product exists. We state now the distributional version of Huygens principle for the 'squared' Dirac operator on the spinor bundle. Recall that all distributional solutions of $(\square + (m^2 + \frac{1}{4}R)) u = 0$ can be restricted to $\mathcal{C}$, since by Theorem 3.21 their wave front sets contain at most lightlike directions.

PROPOSITION 4.17 (Huygens principle (distributional form)). *Let $\hat{\varphi}$ denote the natural embedding of $M \times \mathcal{C}$ into $M \times M$. For any distributional solution $u_A \in D'(D^*M)$ of $(\square + (m^2 + \frac{1}{4}R)) u = 0$, the following identity holds for all $f^A \in D(D^*M)$:*

$$\begin{aligned} u_A(f^A) &= \hat{\varphi}^* \left\{ (\mathbf{1}_{A'}^{B'} \otimes u_A) \cdot \left((\mathbf{1}_{B'}^{C'} \otimes n_a \nabla_a) E_{C'}^A \right) \right\} (f^{A'} \otimes \mathbf{1} d\mathcal{C}) \\ &\quad - \hat{\varphi}^* \left\{ (\mathbf{1}_{A'}^{B'} \otimes n_a \nabla^a u_A) \cdot E_{B'}^A \right\} (f^{A'} \otimes \mathbf{1} d\mathcal{C}), \end{aligned} \quad (73)$$

where $\cdot \otimes \cdot$ is the usual tensor product of distributions, $\mathbf{1}_{A'}^{B'}$ denotes the identity on DM , whereas $\mathbf{1}$ denotes the restriction of the function 1 to $\mathcal{C}$. Note that Eqn. (73) is well defined, since $(J^+(\text{supp}(f^A)) \cup J^+(\text{supp}(f))) \cap \mathcal{C}$ is compact.

The fundamental solution S of the Dirac equation can be restricted to $\mathcal{C}$, too, since its wave front set is contained in the wave front set of E . The result is:

PROPOSITION 4.18. *Let $S_A^{A'} \in D'(D^*M \times DM)$ be the unique fundamental solution of the Dirac operator. The restriction of $S_A^{A'}$ to the Cauchy surface $\mathcal{C}$ with normal covectorfield n_a and embedding $\tilde{\varphi}: \mathcal{C} \times \mathcal{C} \rightarrow M \times M$ is well defined and given by:*

For all $f^A d\mathcal{C} \in D(D_C^*M)$, $g_B d\mathcal{C} \in D(D_C M)$

$$\tilde{\varphi}^* \left(S_A^{B'} \right) (f^A, g_{B'}) = \int_{\mathcal{C}} f^A (i\not{n})_A^B g_B d\mathcal{C}, \quad (74)$$

where $D_C M$ ($D_C^* M$) denotes the restriction of DM ($D^* M$) to $\mathcal{C}$.

REMARK . The result of the Proposition is equivalent to the statement that Ξ restricted to $\mathcal{C}$ is $(i\not{n})$ times the identity on $\Gamma_0(D_C M)$.

For the proof of the Proposition we need the following two Lemmata.

LEMMA 4.19. *Let $E_A^{A'} \in D'(D^*M \times DM)$ be the unique fundamental solution of the 'squared' Dirac operator $(\square + (m^2 + \frac{1}{4}R))$.*

- *The restriction of $E_A^{A'}$ to any smooth Cauchy surface $\mathcal{C}$ vanishes, i.e., let $\tilde{\varphi}$ be the embedding of $\mathcal{C} \times \mathcal{C}$ into $M \times M$, then*

$$\tilde{\varphi}^*(E_A^{A'}) = 0 \quad (75)$$

- The restriction of its normal derivative with respect to the first entry gives $-\mathcal{J}_A^{A'}$ times the Dirac delta distribution, i.e.,

$$\tilde{\varphi}^*(n^a \nabla_a E_A^{A'})(x, y) = -\mathcal{J}_A^{A'} \delta(x, y). \quad (76)$$

Recall that $\mathcal{J}$ denotes the bi-spinor of parallel transport and δ is defined as in Proposition 4.13.

PROOF. Let $u_{yA}^{B'} \in D'(D^*M)$ be the unique solution of $(\square + (m^2 + \frac{1}{4}R))u_{yA}^{A'} = 0$ with initial data

$$(u_{yA}^{A'})|_{\mathcal{C}}(x) = 0 \quad ((n^a \nabla_a u_{yA}^{A'})|_{\mathcal{C}}(x) = -\mathcal{J}_A^{A'}(x, y)\delta(x, y),$$

where $\delta(x, y)$ denotes the Dirac delta distribution at $y \in \mathcal{C}$. Using the distributional form of Huygens principle, we obtain for all $f^A \in D(D^*M)$ —using the notation of Proposition 4.17—

$$\begin{aligned} u_{yA}^{B'}(f^A) &= 0 - \varphi^* \left\{ (\mathbf{1}_C^A \otimes n^a \nabla_a u_{yB''}^{B'}) \cdot E_A^{B''} \right\} (f^C \otimes \mathbf{1} d\mathcal{C}) \\ &= \left((\mathbf{1}_C^A \otimes \mathcal{J}_{B''}^{B'} \delta(\cdot, y)) \varphi^*(E_A^{B''}) \right) (f^C \otimes \mathbf{1} d\mathcal{C}) \\ &= E_A^{B'}(f^A, y) \end{aligned} \quad (77)$$

This proves $u_{yA}^{B'} = E_A^{B'}(\cdot, y)$ for all points $y \in \mathcal{C}$. Note that $E_A^{B'}(f^A, y)$ is smooth in y , since we are dealing with the fundamental solution of the ‘squared’ Dirac equation (recall the results for the scalar case). We conclude: $E_A^{B'}(\cdot, y)|_{\mathcal{C}} = 0$ and $(n^a \nabla_a E_A^{A'}(\cdot, y))|_{\mathcal{C}} = -\mathcal{J}_A^{A'} \delta(\cdot, y)$ for all $y \in \mathcal{C}$, which was claimed. $\square$

The second Lemma is the analogon of Lemma 4.14 above.

LEMMA 4.20. Let $\mathcal{C}$ be a spacelike hypersurface and let $u^A \in D'(DM)$ be a spinor valued distribution on M such that $\varphi^*(u^A)$, the restriction of u^A to $\mathcal{C}$, exists. Let v^a be a smooth vector field tangential to $\mathcal{C}$ and let h_{ab} , D_a denote the induced metric and Levi-Civita connection respectively⁴ (See Definition 4.3 and Lemma 4.4). Then

$$\varphi^*(v^a \nabla_a u^A) = v^a D_a \varphi^*(u^A).$$

PROOF. Suppose u^A is given by a smooth section with compact support in DM , then for any smooth section f_A in the cospinor bundle $v^a u^A f_A$ is a smooth vector field and the integral of $D_a(v^a u^A f_A)$ over $\mathcal{C}$ vanishes by Stokes Theorem. We can apply the latter, since D_a is the Levi Civita connection on $\mathcal{C}$. We obtain the following chain of equation:

$$\begin{aligned} 0 &= \int_{\mathcal{C}} D_a(v^a u^A f_A) d\mathcal{C} \\ &= \int_{\mathcal{C}} (D_a v^a) u^A f_A + v^a ((\nabla_a u^A) f_A + u^A (\nabla_a f_A)) d\mathcal{C} \end{aligned}$$

since D_a is a derivation and on scalars $v^a D_a \equiv v^a \nabla_a$

$$= \int_{\mathcal{C}} u^A D_a(v^a f_A) + v^a (\nabla_a u^A) f_A d\mathcal{C}$$

since D_a is a derivation and $v^a D_a \equiv v^a \nabla_a$ for pure (co)spinors

$$=: (-v^a D_a \varphi^*(u^A))(f_A) + \varphi^*(v^a \nabla_a u^A)(f_A)$$

Note that the last equality is obtained by using the definition of the covariant derivative for distributions. Moreover we used the metric to identify functions and densities. By continuity this extends to all of $D'(M)$, which proves the Lemma. $\square$

REMARK . The previous Lemma is not a complete analogon of Lemma 4.14. In the former everything is independent of the fact that $\mathcal{C}$ is embedded into M , e.g., all tensorial objects are tangential to $\mathcal{C}$ and covariant differentiation does not change this fact. On the other hand in the spinorial case, the embedding enters via the spinor bundle, i.e., spinors on $\mathcal{C}$ are elements in the bundle $D_{\mathcal{C}}M$ consisting of all spinors in DM , whose base points are elements of $\mathcal{C}$. It follows that $D_{\mathcal{C}}M$ can not be constructed without specifying the embedding $\mathcal{C} \xrightarrow{i} M$, i.e., the elements of $D_{\mathcal{C}}M$ do not lie in some subbundle ‘tangential’ to $\mathcal{C}$. It is worth noting that such a subbundle actually can be defined (See [NO84]). However such a treatment yields Spin(3)-spinors, which is undesirable from our point of view.

⁴We extend Eqn. (62) without modification to spinors, i.e., for a smooth section f^A in the spinor bundle

$$D_a f^A \equiv h_a{}^b \nabla_b f^A.$$

Note that we do not project on the spinor indices.

PROOF OF PROPOSITION 4.18. Using Proposition 4.16 above we rewrite the fundamental solution S (suppressing all spinor indices)

$$S = (-i\nabla - m)E = (-i\gamma_a h^{ab}\nabla_b - i\not{n}^a\nabla_a - m)E \quad (78)$$

For the restriction of S to $\mathcal{C} \times \mathcal{C}$ we obtain by Lemma 4.19

$$\tilde{\varphi}^*(S)(x, y) = \tilde{\varphi}^*(-(i\gamma_a h^{ab}\nabla_b E))(x, y) + i\not{n}\mathcal{J}\delta(x, y) - m\tilde{\varphi}^*(E)(x, y) \quad (79)$$

The first summand vanishes by Lemma 4.20, since $\gamma_a h^{ab}$ is tangential to $\mathcal{C}$, the third term is zero by Lemma 4.19. The remaining term in the middle is the result that was claimed. $\square$

PROOF OF THEOREM 4.9. To prove the Theorem, we calculate the (anti) commutators explicitly. It was already shown in Proposition 4.8 that the restrictions of all six operator valued distributions V_{x_0} yield operator valued distributions on $\mathcal{C}$. Consider $\tilde{\varphi}^*({}^A V_{x_0}) \in D'(\mathcal{C} \times \mathcal{C})$. Inserting the definitions and writing all distributions as integrals we obtain:

$$\begin{aligned} \tilde{\varphi}^*([\cdot Q_{A''}^+(x_0), A(\cdot)]_-)(\chi d\mathcal{C}, \mathbf{h}d\mathcal{C}) &= [\chi^{d\mathcal{C}} Q_{A''}^+(x_0), A(\mathbf{h}d\mathcal{C})]_- \\ &= \iint_{\mathcal{C} \times \mathcal{C}} [k_A^+(x) \mathcal{J}^A_{A''}(x, x_0) n_a(x), A(y)]_- \chi(x) \mathbf{h}(y) d\mathcal{C}_x d\mathcal{C}_y \\ &= \iint_{\mathcal{C} \times \mathcal{C}} (-i)(\Psi^+ \gamma^a (-i\nabla_x - m))_A [A(x), A(y)]_- \mathcal{J}^A_{A''}(x, x_0) n_a(x) \chi(x) \mathbf{h}(y) d\mathcal{C}_x d\mathcal{C}_y \end{aligned} \quad (80)$$

since A commutes with B and Ψ

$$\begin{aligned} &= \iint_{\mathcal{C} \times \mathcal{C}} (-1)(\Psi^+(x) \gamma^a \gamma^b)_A \mathcal{J}^A_{A''} n_a \nabla_b [A(x), A(y)]_- (\chi(x) \mathbf{h}(y)) \\ &\quad + im(\Psi^+(x) \gamma^a)_A \mathcal{J}^A_{A''} n_a [A(x), A(y)]_- \chi(x) \mathbf{h}(y) d\mathcal{C}_x d\mathcal{C}_y \end{aligned}$$

We are dealing with free fields, hence on the GNS Hilbertspace we have the identity:

$$[A(x), A(y)]_- = iE(x, y)\mathbf{1},$$

where $\mathbf{1}$ denotes the identity on the Hilbertspace. It was shown in Proposition 4.13 that the restriction of E to $\mathcal{C}$ vanishes, whereas the restriction of its derivative in direction v^a is given by $-v^a n_a$ times the Dirac delta distribution by Corollary 4.15. Therefore our result is

$$(80) = i \int_{\mathcal{C}} \Psi^+_A \mathcal{J}^A_{A''}(x, x_0) \mathbf{h}(x) d\mathcal{C}_x,$$

since $\gamma^a \gamma^b n_a n_b = g^{ab} n_a n_b = +1$ and the y -integration can be carried out explicitly. Note that the result is independent of χ since we assumed $\chi|_{\text{supp}h} \equiv 1$. The pseudoscalar case (replace ${}^A V_{x_0}$ by ${}^B V_{x_0}$) is -up to a factor $(-i\gamma^5)$ - identical to the previous one:

$$\begin{aligned} [\chi^{d\mathcal{C}} Q_{A''}^+(x_0), B(\mathbf{h}d\mathcal{C})]_- &= \iint_{\mathcal{C} \times \mathcal{C}} (-i)(\Psi^+ \gamma^a (-i\nabla_x - m)(-i\gamma^5))_A [A(x), A(y)]_- \\ &\quad \mathcal{J}^A_{A''}(x, x_0) n_a(x) \chi(x) \mathbf{h}(y) d\mathcal{C}_x d\mathcal{C}_y \\ &= \int_{\mathcal{C}} \Psi^+_A(x) \gamma^5 \mathcal{J}^A_{A''}(x, x_0) \mathbf{h}(x) d\mathcal{C}_x \end{aligned} \quad (81)$$

Finally for the spinorial case we get:

$$\{\Psi^{A'}(\mathbf{h}_{A'}d\mathcal{C}), \chi^{d\mathcal{C}}Q_{A''}^+(x_0)\}_+ = \iint_{\mathcal{C} \times \mathcal{C}} (-i)\{\Psi^{A'}(y), \Psi^+_B(x)\}_+ (\not{x}(-i\nabla_x - m)(A - i\gamma^5 B)(x))^B_A \mathcal{J}^A_{A''}(x, x_0) \mathbf{h}_{A'}(y) \chi(x) d\mathcal{C}_x d\mathcal{C}_y$$

Since the anti-commutator of Ψ and Ψ^+ is given by (Note the minus sign!)

$$\{\Psi^{A'}(y), \Psi^+_B(x)\}_+ = -iS^{A'}_B(y, x)\mathbf{1}$$

one obtains

$$\begin{aligned} &= \iint_{\mathcal{C} \times \mathcal{C}} (-1) \mathbf{h}_{A'}(y) S^{A'}_B(y, x) \\ &\quad (\not{x}(-i\nabla_x - m)(A - i\gamma^5 B)(x))^B_A \mathcal{J}^A_{A''}(x, x_0) \chi(x) d\mathcal{C}_y d\mathcal{C}_x \\ &= \int_{\mathcal{C}} (-1) \mathbf{h}_A(x) i \underbrace{(\not{x} \not{x})}_{=1} (-i\nabla_x - m)(A - i\gamma^5 B)(x) \mathcal{J}^A_{A''} \chi(x) d\mathcal{C}_x \end{aligned} \quad (82)$$

where the result of Proposition 4.18 was used.

$$= \int_{\mathcal{C}} i \mathbf{h}_A(x) ((i\nabla_x + m)(A - i\gamma^5 B)(x))^A_B \mathcal{J}^B_{A''}(x, x_0) d\mathcal{C}_x$$

The last equation is valid since we assume $\chi \equiv 1$ on those points where $\mathbf{h}_A$ does not vanish. Let us consider now:

$$\begin{aligned} &[\chi^{d\mathcal{C}}Q_{A''}^+(x_0), \nabla_a A(\mathbf{g}^a d\mathcal{C})]_- \\ &= \iint_{\mathcal{C} \times \mathcal{C}} (-i)(\Psi^+ \not{x}(-i\nabla_x - m))_A i(\nabla_{a'} E(x, y)) \chi(x) \mathbf{g}^{a'}(y) \mathcal{J}^A_{A''}(x, x_0) d\mathcal{C}_x d\mathcal{C}_y \\ &= \iint_{\mathcal{C} \times \mathcal{C}} \Psi^+_B(x) \not{x} \left(\underbrace{-i\gamma^b h_b^c h_{a'}^{c'} \nabla_c \nabla_{c'} E}_I - \underbrace{i \not{x} n^c h_{a'}^{c'} \nabla_c \nabla_{c'} E}_II \right. \\ &\quad \left. - \underbrace{i\gamma^b n_{a'} h_b^c n^{c'} \nabla_c \nabla_{c'} E}_III - \underbrace{i \not{x} n_{a'} n^c n^{c'} \nabla_c \nabla_{c'} E}_IV \right. \\ &\quad \left. - \underbrace{m h_{a'}^{c'} \nabla_{c'} E}_V - \underbrace{m n_{a'} n^{c'} \nabla_{c'} E}_VI \right)^B_A \\ &\quad \chi(x) \cdot \mathbf{g}^{a'}(y) \mathcal{J}^A_{A''} d\mathcal{C}_x d\mathcal{C}_y \end{aligned} \quad (83)$$

For the six terms (I-VI) we find the following results (As usual $\tilde{\varphi}$ denotes the natural embedding of $\mathcal{C} \times \mathcal{C}$ into $M \times M$):

$$\text{I:} \quad \tilde{\varphi}^* \left(h_b^c h_{a'}^{c'} \nabla_c \nabla_{c'} E \right) \left(\tilde{\chi}^b, \tilde{\mathbf{g}}^{a'} \right) = 0 \quad \forall \tilde{\chi}^a, \tilde{\mathbf{g}}^a \in D(T\mathcal{C}) \quad (84)$$

by Corollary 4.15.

$$\begin{aligned} \text{II:} \quad \tilde{\varphi}^* \left(n^c h_{a'}^{c'} \nabla_c \nabla_{c'} E \right) (\chi d\mathcal{C}, \mathbf{g}^{a'} d\mathcal{C}) &= h_{a'}^{c'} D_{c'} \tilde{\varphi}^* (n^c \nabla_c E) (\chi d\mathcal{C}, \mathbf{g}^{a'} d\mathcal{C}) \\ &= +\chi(D_c(h_a^c \mathbf{g}^a d\mathcal{C}), \end{aligned} \quad (85)$$

since $\tilde{\varphi}^*(n^c \nabla_c E)(x, y) = -\delta(x, y)$ by Proposition 4.13. For the third term let us consider the whole integral expression:

$$\begin{aligned}
 \text{III:} \quad & -i\tilde{\varphi}^* \left((\Psi^+ \not{n} \gamma^b \mathcal{J})_{A''} (n_{a'} h_b^c n^{c'} \nabla_c \nabla_{c'} E) \right) (\chi d\mathcal{C}, g^{a'} d\mathcal{C}) \\
 & = \left(-i\tilde{\varphi}^* ((\Psi^+ \not{n} \gamma^b \mathcal{J})_{A''}) \cdot h_b^c D_c \tilde{\varphi}^*(n^{c'} \nabla_{c'} E) \right) (\chi d\mathcal{C}, n_{a'} g^{a'} d\mathcal{C}) \\
 & = +iD_c (h_b^c \tilde{\varphi}^* ((\Psi^+ \not{n} \gamma^b \mathcal{J})_{A''})) \cdot \tilde{\varphi}^*(n^{c'} \nabla_{c'} E) (\chi d\mathcal{C}, n_{a'} g^{a'} d\mathcal{C}) \\
 & = +ih_b^c D_c \tilde{\varphi}^* ((\Psi^+ \not{n} \gamma^b \mathcal{J})_{A''}) (n_a g^a d\mathcal{C})
 \end{aligned}$$

Note that D_c is the covariant derivative on $\mathcal{C}$, hence $D_b h_{cd} = 0$ and we can apply Leibnitz rule on the derivative of products. The boundary term $D_c(\cdot)$ vanishes since we assume $\chi \equiv 1$ on the support of $g^{a'}$. We also used that the normal derivative of E with respect to the second entry is the Dirac delta distribution (compare VI:).

$$= +i \int_{\mathcal{C}} n_a h_b^c \nabla_c (\Psi^+ \not{n} \gamma^b \mathcal{J})_{A''} g^a d\mathcal{C},$$

since $h_b^c D_c v^b = h_b^c \nabla_c v^b$ for all vector fields v^b on $\mathcal{C}$. The fourth terms vanishes by Huygens principle, since for the (distributional) solution u_y of the Klein Gordon Equation with the following initial conditions on $\mathcal{C}$

$$u_y \upharpoonright_{\mathcal{C}} = \delta(\cdot, y) \quad (n^a \nabla_a u_y) \upharpoonright_{\mathcal{C}} = 0$$

we obtain by Proposition 4.12:

$$D'(M) \ni u_y(\cdot) = n^{c'} \nabla_{c'} E(\cdot, y)$$

$$\text{IV:} \quad \tilde{\varphi}^*(n^c n^{c'} \nabla_c \nabla_{c'} E) = 0 \tag{86}$$

The fifth term vanishes by Proposition 4.13:

$$\text{V:} \quad \tilde{\varphi}^*(h_a^c \nabla_c E) = 0.$$

The result of IV: confirms

$$\text{VI:} \quad \tilde{\varphi}^*(n^{c'} \nabla_{c'} E)(x, y) = +\delta(x, y). \tag{87}$$

Inserting these expressions in (83) yields:

$$(83) = \int_{\mathcal{C}} +ih_a^c D_c (\Psi^+ \mathcal{J})_{A''} g^a + ih_b^c D_c ((\Psi^+ \not{n} \gamma^b \mathcal{J})_{A''}) n_a g^a - m(\Psi^+ \not{n} \mathcal{J})_{A''} n_a g^a d\mathcal{C} \tag{88}$$

Since the result for B differs by a factor $(-i\gamma^5)$ only, we obtain for the corresponding kernel:

$$[\chi^{d\mathcal{C}} Q_{A''}^+(x_0), \nabla_a B(x)]_- = -i \left(+iD_a (\Psi^+ \gamma^5 \mathcal{J})_{A''} + iD_b (\Psi^+ \not{n} \gamma^b \gamma^5 \mathcal{J})_{A''} n_a - m(\Psi^+ \not{n} \gamma^5 \mathcal{J})_{A''} n_a \right)$$

Finally for the spinorial case consider the map

$$\begin{aligned}
& \{\nabla_a \Psi^A(g_A^a d\mathcal{C}), \mathfrak{X}^{d\mathcal{C}} Q_{A''}^+(x_0)\}_+ \\
&= \iint_{\mathcal{C} \times \mathcal{C}} \{\nabla_a \Psi^A(x), -i(\Psi^+ \not{\nabla}_y - m)_{A'}(A - i\gamma^5 B)^{A'}_{B'}(y)\}_+ \\
&\quad g_A^a(x) \chi(y) \mathcal{J}^{B'}_{A''}(y, x_0) d\mathcal{C}_x d\mathcal{C}_y \\
&= \iint_{\mathcal{C} \times \mathcal{C}} (-i) \nabla_a \underbrace{\{\Psi^A(x), \Psi^+_{A'}(y)\}_+}_{=-iS^A_{A'}(x, y)} (\not{\nabla}_y - m)(A - i\gamma^5 B)(y) \mathcal{J}^{A'}_{A''} \\
&\quad g_A^a(x) \chi(y) d\mathcal{C}_x d\mathcal{C}_y \\
&= - \iint_{\mathcal{C} \times \mathcal{C}} (\not{\nabla}_y - m)(A - i\gamma^5 B) \mathcal{J}^{A'}_{A''} \cdot (\nabla_a S^A_{A'}(x, y)) \\
&\quad g_A^a(x) \chi(y) d\mathcal{C}_x d\mathcal{C}_y
\end{aligned}$$

Inserting $S^A_{A'}(x, y) \equiv E^A_{B'}(x, y)(+i\overleftarrow{\not{\nabla}}_y - m)^{B'}_{A'}$, which is valid since $S^A_{A'}$ is the fundamental solution of the Dirac equation (Compare Proposition 4.16), we get

(89)

$$\begin{aligned}
&= - \iint_{\mathcal{C} \times \mathcal{C}} (\not{\nabla}_y - m)(A - i\gamma^5 B) \mathcal{J}^{A'}_{A''} \\
&\quad \times \nabla_a E^A_{B'}(+i\overleftarrow{\not{\nabla}}_y - m)^{B'}_{A'} g_A^a(x) \chi(y) d\mathcal{C}_x d\mathcal{C}_y
\end{aligned}$$

Similar to the calculation done above, we treat derivatives tangential and normal to $\mathcal{C}$ individually:

$$\begin{aligned}
&= - \iint_{\mathcal{C} \times \mathcal{C}} (\not{\nabla}_y - m)(A - i\gamma^5 B) \mathcal{J}^{A'}_{A''} \\
&\quad \left(\underbrace{(h_a^c h_{b'}^{c'} \nabla_c \nabla_{c'} E^A_{B'})(+i\gamma^{b'})^{B'}_{A'}}_I + \underbrace{(h_a^c n_{b'} n^{c'} \nabla_c \nabla_{c'} E^A_{B'})(+i\gamma^{b'})^{B'}_{A'}}_{II} \right. \\
&\quad + \underbrace{(n_a n^c h_{b'}^{c'} \nabla_c \nabla_{c'} E^A_{B'})(+i\gamma^{b'})^{B'}_{A'}}_{III} + \underbrace{(n_a n^c n_{b'} n^{c'} \nabla_c \nabla_{c'} E^A_{B'})(+i\gamma^{b'})^{B'}_{A'}}_{IV} \\
&\quad \left. - m(\underbrace{(h_a^c \nabla_c E^A_{A'})}_V + \underbrace{(n_a n^c \nabla_c E^A_{A'})}_{VI}) \right) g_A^a(x) \chi(y) d\mathcal{C}_x d\mathcal{C}_y
\end{aligned}$$

The first term (I:) vanishes since we can apply Lemma 4.20.

$$\begin{aligned}
\text{I:} \quad & \tilde{\varphi}^*(h_a^c h_{a'}^{c'} \nabla_c \nabla_{c'} E^A_{A'}) (\tilde{\chi}_A^a, \tilde{\mathbf{g}}^{a'A'}) = \tilde{\varphi}^*(E^A_{A'}) (D_c(h_a^c \tilde{\chi}_A^a), D_{c'}(h_{a'}^{c'} \tilde{\mathbf{g}}^{a'A'})) \\
&= 0 \quad \forall \tilde{\chi}_A^a \in D(D_{\mathcal{C}} M \otimes T^* \mathcal{C}), \tilde{\mathbf{g}}^{a'A'} \in D(D_{\mathcal{C}}^* M \otimes T^* \mathcal{C})
\end{aligned}$$

For the second term let us consider the whole integral expression:

$$\begin{aligned} \text{II:} \quad & - \iint_{\mathcal{C} \times \mathcal{C}} (i \not{\nabla}_y ((-i \not{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J})^{A'}_{A''} h_a{}^b n^{c'} \nabla_c \nabla_{c'} E^A_{A'} g_A^a(x) \chi(y) d\mathcal{C}_x d\mathcal{C}_y \\ & = + \iint_{\mathcal{C} \times \mathcal{C}} (i ((-i \not{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J})^{A'}_{A''} n^{c'} \nabla_{c'} E^A_{A'} D_c(h_a{}^c g_A^a) \chi(y) d\mathcal{C}_x d\mathcal{C}_y \end{aligned}$$

by Lemma 4.20

$$= + \int_{\mathcal{C}} (i ((-i \not{\nabla}_x - m)(A - i\gamma^5 B)) \mathcal{J})^A_{A''} D_c(h_a{}^c g_A^a) d\mathcal{C}_x,$$

since $\chi \equiv 1$ on the support of g_A^a and the normal derivative of $E^A_{A'}$ with respect to the *second* entry restricted to $\mathcal{C}$ is plus the Dirac delta distribution by Lemma 4.19

$$\begin{aligned} & =: - \int_{\mathcal{C}} h_a{}^c D_c(i ((-i \not{\nabla}_x - m)(A - i\gamma^5 B)) \mathcal{J})^A_{A''} g_A^a d\mathcal{C}_x \\ & = \varphi^* \left(-i h_a{}^c \nabla_c \left(i ((-i \not{\nabla}_x - m)(A - i\gamma^5 B)) \mathcal{J} \right)^A_{A''} \right) (g_A^a) \end{aligned}$$

The third term yields:

$$\begin{aligned} \text{III:} \quad & - \iint_{\mathcal{C} \times \mathcal{C}} (i \gamma^{b'} \not{\nabla} ((-i \not{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J})^{A'}_{A''} n^c h_b{}^{c'} \nabla_c \nabla_{c'} E^A_{A'} n^a g_A^a(x) \chi(y) d\mathcal{C}_x d\mathcal{C}_y \\ & = + \iint_{\mathcal{C} \times \mathcal{C}} D_{c'}(h_b{}^{c'} \chi(y) \gamma^{b'} i \not{\nabla} ((-i \not{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J})^{A'}_{A''} n^c \nabla_c E^A_{A'} n^a g_A^a(x) d\mathcal{C}_x d\mathcal{C}_y \end{aligned}$$

since $h_b{}^c \nabla_c \equiv h_b{}^c D_c$ for cospinors. This allows the necessary integration by parts. The boundary term vanishes, since χ and g_A^a have compact support.

$$= - \int_{\mathcal{C}} \gamma^b h_b{}^c D_c(i \not{\nabla} ((-i \not{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J})^A_{A''} n_a g_A^a d\mathcal{C},$$

since the normal derivative of $E^A_{A'}$ with respect to the first entry restricted to $\mathcal{C}$ is minus the Dirac delta distribution by Lemma 4.19. Recall that χ is one on the support of g_A^a and note that the metric h_{ab} is covariantly constant for D_c . Moreover, the same result holds for the Dirac γ -matrices, i.e., $D_c \gamma^b = h_c{}^d h^b{}_d \nabla_d \gamma^b = 0$.

$$= \varphi^* \left(-i n_a \gamma^b h_b{}^c \nabla_c (i \not{\nabla} ((-i \not{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J})^A_{A''} \right) (g_A^a)$$

The forth term is shown to vanish by using the same argument as in the scalar case: Consider the unique solution $u_y^A{}_{B'} : D(DM) \rightarrow \Gamma(D^*M)$ of the SUSY-coupled Klein Gordon operator with distributional initial data $u_y^A{}_{B'} \upharpoonright_{\mathcal{C}}$ is the Dirac delta distribution and $n^a \nabla_a u_y^A{}_{B'} \upharpoonright_{\mathcal{C}}$ vanishes. Inserting these initial data into Eqn. (73), we get

$$u_y^A{}_{B'} = (n^{c'} \nabla_{c'} E)^A_{B'}$$

which verifies the statement, i.e.,

$$\text{IV:} \quad \tilde{\varphi}^*(n^c n^{c'} \nabla_c \nabla_{c'} E^A_{B'}) = 0.$$

By Lemma 4.19 we get for the fifth term:

$$\text{V:} \quad \tilde{\varphi}^*(h_a{}^c \nabla_c E^A_{A'}) = 0$$

Finally, since the normal derivative of $E^A_{A'}$ with respect to the first entry restricted to $\mathcal{C}$ is minus the Dirac delta distribution, the sixth term is

$$\begin{aligned} \text{VI:} \quad & \iint_{\mathcal{C} \times \mathcal{C}} +m(\not{\nabla} ((-i \not{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J})^{A'}_{A''} n^c \nabla_c E^A_{A'} n_a g_A^a \chi(y) d\mathcal{C}_x d\mathcal{C}_y \\ & = \int_{\mathcal{C}} -m(\not{\nabla} ((-i \not{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J})^A_{A''} n_a g_A^a d\mathcal{C}_x \\ & = \varphi^* \left(-m n_a (\not{\nabla} ((-i \not{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J})^A_{A''} \right) (g_A^a) \end{aligned} \tag{90}$$

To summarize:

$$\begin{aligned}
 (89) = & \varphi^* \left(-i h_a^c \nabla_c \left(((-i \overleftarrow{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J} \right)^A_{A''} \right. \\
 & - i n_a \gamma^b h_b^c \nabla_c (\not{n}((-i \overleftarrow{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J})^A_{A''} \\
 & \left. - m n_a (\not{n}((-i \overleftarrow{\nabla} - m)(A - i\gamma^5 B)) \mathcal{J})^A_{A''} \right) (g_A^a)
 \end{aligned} \tag{91}$$

□

2.5. Traces for the ‘supercharge’. In this subsection we calculate the traces of the anti-commutator of the ‘supercharge’ with various point-separated currents. The results of the following Proposition will be used in the next section to calculate the expectation value of our energy-momentum tensor (Definition 4.22 below) in a locally SUSY Hadamard product state. We use the results of Proposition 4.8 to calculate the following traces:

PROPOSITION 4.21. *Consider two points x and y on $\mathcal{C}$. The corresponding vector and spinor indices are denoted by a, A and a', A' respectively and a semicolon is a shorthand for covariant differentiation. Then we have the following three identities in the sense of distributional kernels on $\mathcal{C}$.*

$$\begin{aligned}
 & \text{Tr} \left(\gamma^{aD}_A \left\{ i((A - i\gamma^5 B)(i \overleftarrow{\nabla} - m)\gamma^b)^A_C \mathcal{J}^C_{C'} \Psi^{C'}, {}^c G_B^+ \right\}_+ \right) \\
 = & 2 \left(g^b_{b'} (\{A^{;a}, A^{;b'}\}_+ + \{B^{;a}, B^{;b'}\}_+) + g^a_{a'} (\{A^{;b}, A^{;a'}\}_+ + \{B^{;b}, B^{;a'}\}_+) \right. \\
 & - g^{ab} g^c_{c'} (\{A^{;c}, A^{;c'}\}_+ + \{B^{;c}, B^{;c'}\}_+) + m^2 g^{ab} (\{A, A'\}_+ + \{B, B'\}_+) \\
 & + 2g^{d'}_d \epsilon^{abcd} (A^{;c} B^{;d'} - B^{;c} A^{;d'}) \\
 & \left. + i \text{Tr}(\gamma^b \mathcal{J}[\Psi', \Psi^{+;a}]_-) + iK \text{Tr}((\gamma^a \not{n} \gamma^b \mathcal{J})[\Psi', \Psi^+]_-) \right)
 \end{aligned} \tag{92}$$

$$\begin{aligned}
 & \text{Tr} \left(\gamma^{aD}_A \left\{ ((\gamma^b \overleftarrow{\nabla} - \overleftarrow{\nabla} \gamma^b)(A - i\gamma^5 B))^A_C \mathcal{J}^C_{C'} \Psi^{C'}, {}^c G_B^+ \right\}_+ \right) \\
 = & 4 \left(-g^{ab} g^c_{c'} (\{A^{;c}, A^{;c'}\}_+ + \{B^{;c}, B^{;c'}\}_+) + g^a_{a'} (A^{;a'} A^{;b} + B^{;a'} B^{;b}) \right. \\
 & + g^b_{b'} (A^{;a} A^{;b'} + B^{;a} B^{;b'}) + 2g^{d'}_d \epsilon^{abcd} (A^{;c} B^{;d'} - B^{;c} A^{;d'}) \\
 & \left. - i \text{Tr}(\gamma^a G^{[bc]} \mathcal{J}[\Psi', \Psi^{+;c}]_-) \right)
 \end{aligned} \tag{93}$$

$$\begin{aligned}
 & \text{Tr} \left(\gamma^{aD}_A \mathcal{J}^A_{A'} \left\{ (A' - i\gamma^5 B')^{A'}_{C'} \mathcal{J}^{C'}_C ((\gamma^b \overleftarrow{\nabla} - \overleftarrow{\nabla} \gamma^b) \Psi)^C \mathcal{J}^C_{C'}, {}^c G_B^+ \right\}_+ \right) \\
 = & 4(\{A^{;ab}, A'\}_+ + \{B^{;ab}, B'\}_+) + 4(g^{ab} m^2 + \xi R n^a n^b)(\{A, A'\}_+ + \{B, B'\}_+ \\
 & - 4i \text{Tr}(\mathcal{J} \gamma^a G^{[bc]} [\nabla_c \Psi, \Psi'^+]_-) \\
 & - 2K \text{Tr}(\gamma^a G^{[bc]} n_c \{ \overleftarrow{\nabla}(A - i\gamma^5 B), (A' - i\gamma^5 B') \}_+),
 \end{aligned} \tag{94}$$

where Tr means the trace over the suppressed spinor indices, K denotes the extrinsic curvature of $\mathcal{C}$, $G^{[AB]} = \frac{1}{4}(\gamma^a \gamma^b - \gamma^b \gamma^a)$, ϵ^{abcd} is the totally anti-symmetric ϵ tensor on (M, g_{ab}) and ξ denotes the coupling of the Klein Gordon equation for the fields A and B .

PROOF. We start with the first equation. Write the anti-commutator as a sum of commutators and anti-commutators in the basic fields with ${}^c G_B^+$.

$$\begin{aligned} & \text{Tr} \left(\gamma^{aD}{}_A \left\{ i((A - i\gamma^5 B)(i\overleftarrow{\nabla} - m)\gamma^b)^A{}_C \mathcal{J}^C{}_{C'} \Psi^{C'}, {}^c G_B^+ \right\}_+ \right) \\ &= \frac{1}{2} \text{Tr} \left(\gamma^{aD}{}_A i((A - i\gamma^5 B)(i\overleftarrow{\nabla} - m)\gamma^b)^A{}_C \mathcal{J}^C{}_{C'} \left\{ \Psi^{C'}, {}^c G_B^+ \right\} \right) \end{aligned} \quad (92 \text{ A})$$

$$+ \frac{1}{2} \left[{}^c G_D^+ \gamma^{aD}{}_A, i((A - i\gamma^5 B)(i\overleftarrow{\nabla} - m)\gamma^b)^A{}_C \right]_- \mathcal{J}^C{}_{C'} \Psi^{C'} \quad (92 \text{ B})$$

$$+ \frac{1}{2} \text{Tr} \left(\mathcal{J}^C{}_{C'} \left\{ \Psi^{C'}, {}^c G_D^+ \right\}_+ \gamma^{aD}{}_B i((A - i\gamma^5 B)(i\overleftarrow{\nabla} - m)\gamma^b)^B{}_A \right) \quad (92 \text{ C})$$

$$- \frac{1}{2} \text{Tr} \left(\mathcal{J}^C{}_{C'} \Psi^{C'} \left[({}^c G^+ \gamma^a)_A, i((A - i\gamma^5 B)(i\overleftarrow{\nabla} - m)\gamma^b)^A{}_B \right]_- \right) \quad (92 \text{ D})$$

Consider the part (92A):

$$\begin{aligned} & \frac{1}{2} \text{Tr} \left(\gamma^{aD}{}_A i((A - i\gamma^5 B)(i\overleftarrow{\nabla} - m)\gamma^b)^A{}_C \mathcal{J}^C{}_{C'} \left\{ \Psi^{C'}, {}^c G_B^+ \right\}_+ \right) \\ &= \frac{1}{2} \text{Tr} \left(\gamma^{aD}{}_A i((A - i\gamma^5 B)(i\overleftarrow{\nabla} - m)\gamma^b)^A{}_C \mathcal{J}^C{}_{C'} (+i)((i\overleftarrow{\nabla} + m)(A - i\gamma^5 B))^{C'}{}_{B'} \mathcal{J}^{B'}{}_B \right) \\ &= \frac{1}{2} \text{Tr} \left(i\gamma^{aD}{}_A \left(\gamma^c \nabla_c (A + i\gamma^5 B) \gamma^b \right)^A{}_C \mathcal{J}^C{}_{C'} (-i) \left(\gamma^{d'} \nabla_{d'} (A' - i\gamma^{5'} B') \right)^{C'}{}_{B'} \mathcal{J}^{B'}{}_B \right. \\ & \quad + i\gamma^{aD}{}_A \left(\gamma^c \nabla_c (A + i\gamma^5 B) \gamma^b \right)^A{}_C \mathcal{J}^C{}_{C'} (-m) \left((A' - i\gamma^{5'} B') \right)^{C'}{}_{B'} \mathcal{J}^{B'}{}_B \\ & \quad - m\gamma^{aD}{}_A ((A - i\gamma^5 B) \gamma^b)^A{}_C \mathcal{J}^C{}_{C'} (-i) \left(\gamma^{d'} \nabla_{d'} (A' - i\gamma^{5'} B') \right)^{C'}{}_{B'} \mathcal{J}^{B'}{}_B \\ & \quad \left. - m\gamma^{aD}{}_A ((A - i\gamma^5 B) \gamma^b)^A{}_C \mathcal{J}^C{}_{C'} (-m) \left((A' - i\gamma^{5'} B') \right)^{C'}{}_{B'} \mathcal{J}^{B'}{}_B \right) \\ &= \frac{1}{2} \text{Tr} (\gamma^a \gamma^c \gamma^b \gamma^d) g^{d'}{}_d (\nabla_c A \nabla_{d'} A' + \nabla_c B \nabla_{d'} B') \\ & \quad + \frac{1}{2} i \text{Tr} (\gamma^a \gamma^c \gamma^b \gamma^d \gamma^5) g^{d'}{}_d (\nabla_c B \nabla_{d'} A' - \nabla_c A \nabla_{d'} B') + \frac{1}{2} \text{Tr} (\gamma^a \gamma^b) m^2 (AA' + BB') \end{aligned}$$

due to the cyclicity of the trace and since $\mathcal{J}^C{}_{C'} \gamma^{d'C'}{}_{A'} \mathcal{J}^{A'}{}_A = g^{d'}{}_d \gamma^{dC}{}_A$ by Eqn. (117), $(\gamma^5)^2 = 1$ and the trace over an odd number of γ -matrices vanishes.

$$\begin{aligned} &= +2 \left(g^b{}_{b'} (\nabla^a A \nabla^{b'} A' + \nabla^a B \nabla^{b'} B') - g^{ab} g^c{}_{c'} (\nabla_c A \nabla^{c'} A' + \nabla_c B \nabla^{c'} B') \right. \\ & \quad + g^a{}_{a'} (\nabla^b A \nabla^{a'} A' + \nabla^b B \nabla^{a'} B') + g^{d'}{}_d \epsilon^{abcd} (\nabla_c A \nabla_{d'} B' - \nabla_c B \nabla_{d'} A') \\ & \quad \left. + m^2 g^{ab} (AA' + BB') \right), \end{aligned}$$

since the parallel transport commutes with contraction. The various traces for the γ -matrices can be found in Appendix C section 1. For the next part (92B) one obtains:

$$\begin{aligned}
& \frac{1}{2} \left[{}^c G_D^+ \gamma^{aD}{}_A, i((A - i\gamma^5 B)(i\overleftarrow{\nabla} - m)\gamma^b)^A{}_C \right]_- \mathcal{J}^C{}_{C'} \Psi^{C'} \\
&= \frac{1}{2} i \left([{}^c G_D^+, i\nabla_c A]_- (\gamma^{aD}{}_A \gamma^c{}_C) - i[{}^c G_D^+, i\nabla_c B]_- (\gamma^{aD}{}_A (\gamma^5 \gamma^c)^A{}_C) \right. \\
&\quad \left. - m[{}^c G_D^+, A]_- \gamma^{aD}{}_C + im[{}^c G_D^+, B]_- (\gamma^5 \gamma^a)^D{}_C \right) \mathcal{J}^C{}_{C'} \Psi^{C'} \\
&= - \left((i(D_c \Psi^+) + in_c(D_d \Psi^+) \not{n} \gamma^d - mn_c \Psi^+)_D (\gamma^a \gamma^c \gamma^b \mathcal{J} \Psi')^D - m \Psi^+ \gamma^a \gamma^b \mathcal{J} \Psi' \right) \\
&\quad - in_c \Psi^+ (D_d \not{n}) \gamma^d (\gamma^a \gamma^c \gamma^b \mathcal{J} \Psi')^{D'},
\end{aligned}$$

since $(\gamma^5)^2 = 1$. Note that $D_d(\not{n})\gamma^d$ equals K , which is the extrinsic curvature scalar.

$$\begin{aligned}
&= - \left(i(\nabla_c \Psi^+) \gamma^a \gamma^c \gamma^b \mathcal{J} \Psi' - m \Psi^+ \gamma^a \gamma^b \mathcal{J} \Psi' \right) - iK \Psi^+ \gamma^a \not{n} \gamma^b \mathcal{J} \Psi' \\
&= -2i \nabla^a \Psi^+ \gamma^b \mathcal{J} \Psi' - iK \Psi^+ \gamma^a \not{n} \gamma^b \mathcal{J} \Psi'
\end{aligned}$$

where we used the Dirac equation $\Psi^+(i\overleftarrow{\nabla} + m) = 0$ for Ψ^+ .

The result for the third part (92C) is:

$$\begin{aligned}
&+ \frac{1}{2} \text{Tr} \left(\mathcal{J}^C{}_{C'} \left\{ \Psi^{C'}, {}^c G_D^+ \right\}_+ \gamma^{aD}{}_A i((A - i\gamma^5 B)(i\overleftarrow{\nabla} - m)\gamma^b)^A{}_B \right) \\
&= + \frac{1}{2} \text{Tr} \left(-i \mathcal{J} \gamma^{c'} \nabla_{c'} (A' - i\gamma^{5'} B') \mathcal{J} \gamma^a (i) \gamma^d \nabla_d (A + i\gamma^5 B) \gamma^b \right. \\
&\quad - i \mathcal{J} \gamma^{c'} \nabla_{c'} (A' - i\gamma^{5'} B') \mathcal{J} \gamma^a (-m) (A - i\gamma^5 B) \gamma^b \\
&\quad - m \mathcal{J} (A' - i\gamma^{5'} B') \mathcal{J} \gamma^a (i) \gamma^c \nabla_c (A + i\gamma^5 B) \gamma^b \\
&\quad \left. - m \mathcal{J} (A' - i\gamma^{5'} B') \mathcal{J} \gamma^a (-m) (A - i\gamma^5 B) \gamma^b \right) \\
&= + 2 \left(g^a{}_{a'} (\nabla^{a'} A' \nabla^b A + \nabla^{a'} B' \nabla^b B) - g^{ab} g^c{}_{c'} (\nabla_{c'} A' \nabla^c A + \nabla_{c'} B' \nabla^c B) \right. \\
&\quad + g^b{}_{b'} (\nabla'^b A' \nabla^a A + \nabla'^b B' \nabla^a B) \\
&\quad \left. + g^{d'}{}_d \epsilon^{abcd} (\nabla_c A \nabla_{d'} B' - \nabla_c B \nabla_{d'} A') + m^2 g^{ab} (A' A + B' B') \right),
\end{aligned}$$

since up to a commutation of x and y the same terms as in (92A) occur. Finally for the term (92D) one finds:

$$\begin{aligned}
&- \frac{1}{2} \text{Tr} \left(\mathcal{J}^C{}_{C'} \Psi^{C'} \left[({}^c G^+ \gamma^a)_A, i((A - i\gamma^5 B)(i\overleftarrow{\nabla} - m)\gamma^b)^A{}_B \right]_- \right) \\
&= - \text{Tr} \left(\mathcal{J} \Psi' (-2i) \nabla^a \Psi^+ \gamma^b - iK \mathcal{J} \Psi' \Psi^+ \gamma^a \not{n} \gamma^b \right),
\end{aligned}$$

since all terms of (92B) can be used again. To summarize the results:

$$\begin{aligned}
(92) = & +2 \left(g^b_{b'} (\nabla^a A \nabla^{b'} A' + \nabla^a B \nabla^{b'} B') - g^{ab} g^c_{c'} (\nabla_c A \nabla^{c'} A' + \nabla_c B \nabla^{c'} B') \right. \\
& + g^a_{a'} (\nabla^b A \nabla^{a'} A' + \nabla^b B \nabla^{a'} B') + g^{d'}_d \epsilon^{abcd} (\nabla_c A \nabla_{d'} B' - \nabla_c B \nabla_{d'} A') \\
& \left. + m^2 g^{ab} (AA' + BB') \right) \\
& - 2i \text{Tr}(\gamma^b (\nabla^a \Psi^+ \mathcal{J} \Psi') - (iK \Psi^+ \gamma^a \not{n} \gamma^b \mathcal{J} \Psi')) \\
& + 2 \left(g^a_{a'} (\nabla^{a'} A' \nabla^b A + \nabla^{a'} B' \nabla^b B) - g^{ab} g^c_{c'} (\nabla_{c'} A' \nabla^c A + \nabla_{c'} B' \nabla^c B) \right. \\
& + g^b_{b'} (\nabla'^b A' \nabla^a A + \nabla'^b B' \nabla^a B) + g^{d'}_d \epsilon^{abcd} (\nabla_c A \nabla_{d'} B' - \nabla_c B \nabla_{d'} A') \\
& \left. + m^2 g^{ab} (A' A + B' B') \right) \\
& + 2i \text{Tr}(\mathcal{J} \Psi' \nabla^a \Psi^+ \gamma^b) + \text{Tr}(iK \mathcal{J} \Psi' \Psi^+ \gamma^a \not{n} \gamma^b) \\
= & 2 \left(g^b_{b'} (\{A^{;a}, A^{;b'}\}_+ + \{B^{;a}, B^{;b'}\}_+) + g^a_{a'} (\{A^{;b}, A^{;a'}\}_+ + \{B^{;b}, B^{;a'}\}_+) \right. \\
& - g^{ab} g^c_{c'} (\{A_{;c}, A^{;c'}\}_+ + \{B_{;c}, B^{;c'}\}_+) + m^2 g^{ab} (\{A, A'\}_+ + \{B, B'\}_+) \\
& + 2g^{d'}_d \epsilon^{abcd} (A^{;c} B^{d'} - B^{;c} A^{d'}) + i \text{Tr}(\gamma^b \mathcal{J} [\Psi', \Psi^{+;a}]_-) \\
& \left. + iK \text{Tr}((\gamma^a \not{n} \gamma^b \mathcal{J}) [\Psi', \Psi^+]_-) \right)
\end{aligned}$$

We continue with the same calculation for the second Equation:

$$\begin{aligned}
& 4\text{Tr} \left(\gamma^{aD}{}_A \left\{ (G^{[bc]} \nabla_c (A - i\gamma^5 B) \mathcal{J} \Psi')^A, {}^c G^+{}_B \right\}_+ \right) \\
& = 2\text{Tr} \left((\gamma^a G^{[bc]} \nabla_c (A - i\gamma^5 B))^D{}_C \mathcal{J}^C{}_{C'} \left\{ \Psi^{C'}, {}^c G^+{}_B \right\}_+ \right) \tag{93 A}
\end{aligned}$$

$$+ 2 \left[{}^c G^+{}_B \gamma^{aB}{}_A, (G^{[bc]} \nabla_c (A - i\gamma^5 B))^A{}_C \right] \mathcal{J}^C{}_{C'} \Psi^{C'} \tag{93 B}$$

$$+ 2\text{Tr} \left(\mathcal{J}^C{}_{C'} \left\{ \Psi^{C'}, {}^c G^+{}_B \right\}_+ (\gamma^a G^{[bc]} \nabla_c (A - i\gamma^5 B))^A{}_D \right) \tag{93 C}$$

$$- 2\text{Tr} \left(\mathcal{J}^C{}_{C'} \Psi^{C'} \left[{}^c G^+{}_B \gamma^B{}_A, (G^{[bc]} \nabla_c (A - i\gamma^5 B))^A{}_D \right] \right) \tag{93 D}$$

For the part (93A) one calculates:

$$\begin{aligned}
& 2\text{Tr} \left((\gamma^a G^{[bc]} \nabla_c (A - i\gamma^5 B))^D{}_C \mathcal{J}^C{}_{C'} \left\{ \Psi^{C'}, {}^c G^+{}_B \right\}_+ \right) \\
& = 2\text{Tr} \left(\gamma^a G^{[bc]} \nabla_c (A - i\gamma^5 B) \mathcal{J} (+i)((i\not{\nabla} + m)(A' - i\gamma^5 B')) \mathcal{J} \right) \\
& = -2\text{Tr}(\gamma^a G^{[bc]} \gamma^d) g_d{}^{d'} (\nabla_c A \nabla_{d'} A' + \nabla_c B \nabla_{d'} B') \\
& \quad + 2i \text{Tr}(\gamma^a G^{[bc]} \gamma^d \gamma^5) g_d{}^{d'} (\nabla_c A \nabla_{d'} B' - \nabla_c B \nabla_{d'} A') \\
& = -4 \left(g^{ab} g^c_{c'} (\nabla_c A \nabla^{c'} A' + \nabla_c B \nabla^{c'} B') - g^b_{b'} (\nabla^a A \nabla^{b'} A' + \nabla^a B \nabla^{b'} B') \right) \\
& \quad + 4\epsilon^{abcd} g^{d'}_d (\nabla_c A \nabla_{d'} B' - \nabla_c B \nabla_{d'} A')
\end{aligned}$$

The next part (93B) gives:

$$\begin{aligned}
& + 2 \left[{}^C G_B^+ \gamma^{aB}{}_A, (G^{[bc]} \nabla_c (A - i\gamma^5 B))^A{}_C \right] \mathcal{J}^C{}_{C'} \Psi^{C'} \\
& = 2 \left(\left[{}^C G_B^+, \nabla_c A \right]_- (\gamma^a G^{[bc]}) - i \left[{}^C G_B^+, \nabla_c B \right]_- (\gamma^a G^{[bc]} \gamma^5) \right) \mathcal{J} \Psi' \\
& = 2 \left((iD_c \Psi^+ + in_c (D_d \Psi^+) \not{D} \gamma^d - mn_c \Psi^+ \not{D} + iK \Psi^+ n_c) \gamma^a G^{[bc]} \right. \\
& \quad \left. - i(iD_c \Psi^+ + in_c (D_d \Psi^+) \not{D} \gamma^d - mn_c \Psi^+ \not{D} + iK \Psi^+ n_c) \gamma^5 \gamma^a G^{[bc]} \gamma^5 \right) \mathcal{J} \Psi'
\end{aligned}$$

Recall the identity $(D_d \not{D}) \gamma^d = K$.

$$= 4i(\nabla_c \Psi^+) \gamma^a G^{[bc]} \mathcal{J} \Psi',$$

For the last equality we used the fact that Ψ^+ is a solution of the Dirac equation and that $\not{D}$ anti-commutes with $h^a{}_b \gamma^b$. The third term is again relatively easy to compute:

$$\begin{aligned}
& 2\text{Tr} \left(\mathcal{J}^C{}_{C'} \left\{ \Psi^{C'}, {}^C G_B^+ \right\}_+ (\gamma^a G^{[bc]} \nabla_c (A - i\gamma^5 B))^B{}_D \right) \\
& = 2\text{Tr} \left(\mathcal{J}(i)((i\not{D} - m)(A' - i\gamma^5 B')) \mathcal{J}(\gamma^a G^{[bc]} \nabla_c (A - i\gamma^5 B))^B{}_D \right) \\
& = -2\text{Tr}(\gamma^d \gamma^a G^{[bc]}) g_d{}^{d'} (\nabla_{d'} A' \nabla_c A + \nabla_{d'} B' \nabla_c B) \\
& \quad + 2i\text{Tr}(\gamma^a G^{[bc]} \gamma^d \gamma^5) g_d{}^{d'} (\nabla_c A \nabla_{d'} B' - \nabla_c B \nabla_{d'} A'),
\end{aligned}$$

since all other traces vanish

$$\begin{aligned}
& = -4 \left(g^{ab} g_c{}^{c'} (\nabla_{c'} A' \nabla^c A + \nabla_{c'} B' \nabla^c B) - g^a{}_{a'} (\nabla^{a'} A' \nabla^b A + \nabla^{a'} B' \nabla^b B) \right) \\
& \quad + 4\epsilon^{abcd} g_d{}^{d'} (\nabla_c A \nabla_{d'} B' - \nabla_c B \nabla_{d'} A')
\end{aligned}$$

The forth term (93D) gives:

$$\begin{aligned}
& - 2\text{Tr} \left(\mathcal{J} \Psi' \left[{}^C G_B^+ \gamma^{aB}{}_A, (G^{[bc]} \nabla_c (A - i\gamma^5 B))^A{}_D \right]_- \right) \\
& = -4i\text{Tr}(\gamma^a G^{[bc]} \mathcal{J} \Psi' \nabla_c \Psi^+),
\end{aligned}$$

as an immediate consequence of the calculations of part (93B). To summarize the second equation:

$$\begin{aligned}
(93) & = -4 \left(g^{ab} g_c{}^{c'} (\nabla_{c'} A \nabla^{c'} A' + \nabla_{c'} B \nabla^{c'} B') - g^b{}_{b'} (\nabla^a A \nabla^{b'} A' + \nabla^a B \nabla^{b'} B') \right. \\
& \quad \left. - \epsilon^{abcd} g_d{}^{d'} (\nabla_c A \nabla_{d'} B' - \nabla_c B \nabla_{d'} A') \right) \\
& \quad + 4i(\nabla_c \Psi^+) \gamma^a G^{[bc]} \mathcal{J} \Psi' \\
& \quad - 4 \left(g^{ab} g_c{}^{c'} (\nabla_{c'} A' \nabla^c A + \nabla_{c'} B' \nabla^c B) - g^a{}_{a'} (\nabla^{a'} A' \nabla^b A + \nabla^{a'} B' \nabla^b B) \right. \\
& \quad \left. - \epsilon^{abcd} g_d{}^{d'} (\nabla_c A \nabla_{d'} B' - \nabla_c B \nabla_{d'} A') \right) \\
& \quad - 4i\text{Tr}(\gamma^a G^{[bc]} \mathcal{J} \Psi' \nabla_c \Psi^+) \\
& = +4 \left(-g^{ab} g_c{}^{c'} (\{A^{;c}, A^{;c'}\}_+ \{B^{;c}, B^{;c'}\}_+) + g^a{}_{a'} (A^{;a'} A^{;b} + B^{;a'} B^{;b}) \right. \\
& \quad + g^b{}_{b'} (A^{;a} A^{;b'} + B^{;a} B^{;b'}) + 2\epsilon^{abcd} g_d{}^{d'} (\nabla_c A \nabla_{d'} B' - \nabla_c B \nabla_{d'} A') \\
& \quad \left. - i\text{Tr}(\gamma^a G^{[bc]} \mathcal{J} [\Psi', \Psi^+{}_{;c}]_-) \right)
\end{aligned}$$

Finally for the third Equation we consider:

$$\begin{aligned} & \text{Tr} \left(\gamma^{aD} \mathcal{J}^A{}_{A'} \{ (A' - i\gamma^{5'} B')^{A'}{}_{C'} \mathcal{J}^{C'}{}_C ((\gamma^b \not{\nabla} - \not{\nabla} \gamma^b) \Psi)^C, {}^C G_B^+ \}_+ \right) \\ &= 2\text{Tr} \left((\gamma^a \mathcal{J} (A' - i\gamma^{5'} B') \mathcal{J} G^{[bc]})^D{}_C \{ \nabla_c \Psi^C, {}^C G_B^+ \}_+ \right) \end{aligned} \quad (94 \text{ A})$$

$$+ 2 \left[{}^C G_B^+ \gamma^{aB}{}_{A'}, (G^{[bc]} \mathcal{J} (A' - i\gamma^{5'} B') \mathcal{J})^A{}_C \right]_- \nabla_c \Psi^C \quad (94 \text{ B})$$

$$+ 2\text{Tr} \left(\{ \nabla_c \Psi^C, {}^C G_B^+ \}_+ (\gamma^a G^{[bc]} \mathcal{J} (A' - i\gamma^{5'} B') \mathcal{J})^B{}_D \right) \quad (94 \text{ C})$$

$$- 2\text{Tr} \left(\nabla_c \Psi^C \left[{}^C G_B^+ \gamma^{aB}{}_{A'}, (G^{[bc]} \mathcal{J} (A' - i\gamma^{5'} B') \mathcal{J})^A{}_D \right]_- \right) \quad (94 \text{ D})$$

Consider the part (94A) and note that the bi-spinors of parallel transport cancel each other.

$$\begin{aligned} & 2\text{Tr} \left((\gamma^a (A' - i\gamma^5 B') G^{[bc]})^D{}_C \{ \nabla_c \Psi^C, {}^C G_B^+ \}_+ \right) \\ &= 2\text{Tr} \left(\gamma^a (A' - i\gamma^5 B') G^{[bc]} \left(-iD_c((-i\not{\nabla} - m)(A - i\gamma^5 B)) + in_c \not{\gamma}^d D_d((-i\not{\nabla} - m)(A - i\gamma^5 B)) \right. \right. \\ & \quad \left. \left. - mn_c \not{\gamma}((-i\not{\nabla} - m)(A - i\gamma^5 B)) - in_c K(-i\not{\nabla} - m)(A - i\gamma^5 B) \right) \right), \end{aligned}$$

since $\not{\gamma} \not{\gamma} = 1$ and $\not{\gamma}$ commutes with $\gamma^a h_a{}^b$

$$\begin{aligned} &= -2\text{Tr} \left(\gamma^a (A' - i\gamma^5 B') G^{[bc]} \left(D_c \not{\nabla} (A - i\gamma^5 B) - n_c \not{\gamma}^d D_d \not{\nabla} (A - i\gamma^5 B) \right. \right. \\ & \quad \left. \left. - m^2 n_c \not{\gamma} (A - i\gamma^5 B) + K n_c \not{\nabla} (A - i\gamma^5 B) \right) \right), \end{aligned}$$

since traces over an odd number of γ -matrices are zero.

$$\begin{aligned} &= -2\text{Tr} \left(\gamma^a G^{[bc]} (A' - i\gamma^5 B') (\nabla_c \not{\nabla} + \xi R n_c \not{\gamma}) (A - i\gamma^5 B) \right) \\ & \quad - 2K \text{Tr} \left(\gamma^a G^{[bc]} (A' - i\gamma^5 B') n_c \not{\nabla} (A - i\gamma^5 B) \right) \end{aligned}$$

For the last equality we used the fact that A and B are solutions of the Klein Gordon equation with coupling ξ , i.e., $(\square + (m^2 + \xi R))A = 0$ and $(\square + (m^2 + \xi R))B = 0$. It was also used that $\not{\nabla} \not{\nabla} = \square$ for scalars.

$$\begin{aligned} &= 4(A' \nabla^a \nabla^b A + B' \nabla^a \nabla^b B) + 4(m^2 g^{ab} + \xi R n^a n^b)(A' A + B' B) \\ & \quad - 2K \text{Tr} \left(\gamma^a G^{[bc]} (A' - i\gamma^5 B') n_c \not{\nabla} (A - i\gamma^5 B) \right) \end{aligned}$$

The last equality follows, since $\text{Tr}(\gamma^a G^{[bc]} \gamma^d) = 2(g^{ab} g^{cd} - g^{ac} g^{bd})$, whereas the corresponding terms with a *single* γ^5 gives a result totally anti-symmetric in the indices $abcd$ which vanishes due to the symmetry of $\nabla_c \nabla_d$ and $n_c n_d$ respectively. The calculation for the next part (94B) results in:

$$\begin{aligned} & + 2 \left[{}^C G_B^+ \gamma^{aB}{}_{A'}, (G^{[bc]} (A' - i\gamma^5 B'))^A{}_C \right]_- \nabla_c \Psi^C \\ &= 2 \left(\left[{}^C G_B^+, A' \right]_- (\gamma^a G^{[bc]})^B{}_C - i \left[{}^C G_B^+, A' \right]_- (\gamma^a G^{[bc]} \gamma^5)^B{}_C \right) \nabla_c \Psi^C \\ &= 4i \Psi'^+ \mathcal{J} \gamma^a G^{[bc]} \nabla_c \Psi \end{aligned}$$

The third part (94C) yields as a consequence of our results for (94A):

$$\begin{aligned}
& + 2\text{Tr}\left(\{\nabla_c \Psi^C, {}^c G_B^+\}_+ (\gamma^a G^{[bc]}(A' - i\gamma^5 B'))^B{}_D\right) \\
& = 4((\nabla^a \nabla^b A)A' + (\nabla^a \nabla^b B)B') + 4(m^2 g^{ab} + \xi R n^a n^b)(AA' + BB') \\
& \quad - 2K\text{Tr}\left(\overline{\nabla}(A - i\gamma^5 B)\gamma^a G^{[bc]}(A' - i\gamma^5 B')n_c\right)
\end{aligned}$$

The result for the forth term follows immediately from the second:

$$\begin{aligned}
& - 2\text{Tr}\left(\nabla_c \Psi^C \left[{}^c G_B^+ \gamma^{aB}{}_A, (G^{[bc]} \mathcal{J}(A' - i\gamma^5 B') \mathcal{J})^A{}_D\right]_-\right) \\
& = -4i\text{Tr}\left(\mathcal{J} \gamma^a G^{[bc]} \nabla_c \Psi \Psi'^+\right)
\end{aligned}$$

To summarize these results:

$$\begin{aligned}
(94) & = 4(A' \nabla^a \nabla^b A + B' \nabla^a \nabla^b B) + 4(m^2 g^{ab} + \xi R n^a n^b)(A' A + B' B) \\
& \quad - 2K\text{Tr}\left(\gamma^a G^{[bc]}(A' - i\gamma^5 B')n_c \overline{\nabla}(A - i\gamma^5 B)\right) \\
& \quad + 4i\text{Tr}\left(\mathcal{J} \gamma^a G^{[bc]} \Psi'^+ \nabla_c \Psi\right) \\
& = 4((\nabla^a \nabla^b A)A' + (\nabla^a \nabla^b B)B') + 4(m^2 g^{ab} + \xi R n^a n^b)(AA' + BB') \\
& \quad - 2K\text{Tr}\left(\overline{\nabla}(A - i\gamma^5 B)\gamma^a G^{[bc]}(A' - i\gamma^5 B')n_c\right) \\
& \quad - 4i\text{Tr}\left(\mathcal{J} \gamma^a G^{[bc]} \nabla_c \Psi \Psi'^+\right) \\
& = 4(\{A^{;ab}, A'\}_+ + \{B^{;ab}, B'\}_+) + 4(g^{ab} m^2 + \xi R n^a n^b)(\{A, A'\}_+ + \{B, B'\}_+) \\
& \quad - 4i\text{Tr}(\mathcal{J} \gamma^a G^{[bc]} [\nabla_c \Psi, \Psi'^+]_-) \\
& \quad - 2K\text{Tr}\left(\gamma^a G^{[bc]} n_c \{\overline{\nabla}(A - i\gamma^5 B), (A' - i\gamma^5 B')\}_+\right)
\end{aligned}$$

□

3. The energy momentum tensor in terms of ‘supercurrents’ and ‘supercharges’

In this section we define a candidate for a point-separated energy momentum tensor of our locally SUSY model by an analogon of Eqn. (58). Let $(\mathcal{H}, D, A, B, \Psi, \Psi^+, \Omega)$ denote our locally SUSY quantum field theoretical model on the spacetime (M, g_{ab}) . We define two point separated ‘supercurrents’:

$$\begin{aligned}
{}_1 j^{aA}(x, y) & = \frac{1}{2} \left(i(A - i\gamma^5 B)(i\overleftarrow{\nabla} - m)\gamma^a \mathcal{J} \Psi' \right. \\
& \quad \left. - 2\xi \left\{ (4G^{[ab]} \nabla_b (A - i\gamma^5 B)) \mathcal{J} \Psi' + \mathcal{J}(A' - i\gamma^5 B') \mathcal{J} 4G^{[ab]} \nabla_b \Psi \right\} \right)
\end{aligned} \tag{95}$$

$$\begin{aligned}
{}_2 j^{a'A'}(x, y) & = \frac{1}{2} \left(i(A' - i\gamma^5 B')(i\overleftarrow{\nabla} - m)\gamma^{a'} \mathcal{J} \Psi \right. \\
& \quad \left. - 2\xi \left\{ (4G^{[a'b']} \nabla_{b'} (A' - i\gamma^5 B')) \mathcal{J} \Psi + \mathcal{J}(A - i\gamma^5 B) \mathcal{J} 4G^{[a'b']} \nabla_{b'} \Psi' \right\} \right)
\end{aligned} \tag{96}$$

Note that ${}_1 j^{aA}(x, y)$ are two well defined operator valued distributions on $\mathcal{H}$. The constant ξ is undetermined at this stage and will be fixed below to the SUSY coupling $1/4$. Setting

$$j^{aA}(x, y) := {}_1 j^{aA}(x, y) + g^a{}_{a'} \mathcal{J}^A{}_{A'} {}_2 j^{a'A'}(x, y),$$

it follows, using Proposition 4.1 and the fact that $g^a{}_{a'}$ and $\mathcal{J}^A{}_{A'}$ are smooth, that the restriction of j^{aA} to the diagonal exists and equals the improved supercurrent of Eqn. (38). Moreover, since $[\nabla_a g^a{}_{a'}] = [\nabla_a \mathcal{J}^A{}_{A'}] = 0$, where the square brackets denote the coincidence limit, it easy to see that $j^{aA}(x, y)$ is ‘asymptotically conserved’ on spacetimes with vanishing scalar curvature, i.e.,

$$[\nabla_a j^{aA}(x, y)] = \nabla_a [j^{aA}(x, y)] = \nabla_a j^{aA}(x) = 0 \quad \text{iff } R = 0.$$

Recall that the last equality was shown in Proposition 3.2 for classical fields propagating on a spacetime with $R = 0$. This result extends without modification to the quantized version of the theory.

DEFINITION 4.22. Let $\mathcal{C}$ be a smooth maximal spacelike hypersurface in the vacuum spacetime (M, g_{ab}) , i.e., $\mathcal{C}$ has extrinsic curvature $K = 0$. Then the point separated energy momentum tensor $\tilde{T}^{ab}(x, y)$ of a locally SUSY quantum field theoretical model $(\mathcal{H}, D, A, B, \Psi, \Psi^+, \Omega)$ on this spacetime is defined as the operator valued distribution on $\mathcal{C} \times \mathcal{C}$ with kernel (Compare Eqn. (30))

$$\tilde{T}^{ab}(x, y) := \frac{1}{8} \left(\text{Tr} \left(\gamma^{(a} \{ {}_1 j^{b)A}(x, y), {}^c G_{B^+}^+ \} + \right) + g^a{}_{a'} g^b{}_{b'} \text{Tr} \left(\gamma^{(a'} \{ {}_2 j^{b')A'}(x, y), {}^c G_{B'^+}^+ \} + \right) \right), \quad (97)$$

where ${}_{1,2}j^{aA}$ are given by Eqn. (95) and Eqn. (96) respectively with $\xi = 1/4$.

REMARK . The r.h.s. of Eqn. (97) is a well defined operator valued distribution on $\mathcal{C} \times \mathcal{C}$ by Proposition 4.8, even if our spacetime does not have vanishing scalar curvature, $\xi \neq 1/4$ and $\mathcal{C}$ is not maximal.

We will now use the results of Proposition 4.21 to express the r.h.s. of Eqn. (97) as commutators and anti-commutators respectively of the basic fields. Let us relax the assumption on the spacetime (M, g_{ab}) , the hypersurface $\mathcal{C}$ and the coupling ξ for a moment.

THEOREM 4.23. *The point separated energy momentum tensor $\tilde{T}^{ab}(x, y)$ defined by Eqn. (97) is an operator valued distribution on $\mathcal{C} \times \mathcal{C}$, even if the spacetime does not have vanishing scalar curvature, $\xi \neq 1/4$ and $\mathcal{C}$ is not maximal. Its kernel can be written in the basic fields as:*

$$\begin{aligned} \tilde{T}^{ab}(x, y) = & \frac{1-2\xi}{4} g^b{}_{b'} (\{A^{;a}, A^{;b'}\}_+ + \{B^{;a}, B^{;b'}\}_+) + \frac{1-2\xi}{4} g^a{}_{a'} (\{A^{;b}, A^{;a'}\}_+ + \{B^{;b}, B^{;a'}\}_+) \\ & - \frac{1-4\xi}{4} g^{ab} g^c{}_{c'} (\{A_{;c}, A^{;c'}\}_+ + \{B_{;c}, B^{;c'}\}_+) + \frac{1-4\xi}{4} g^{ab} m^2 (\{A, A'\}_+ + \{B, B'\}_+) \\ & - \frac{\xi}{2} \left((\{A^{;ab}, A'\}_+ + \{B^{;ab}, B'\}_+) + g^a{}_{a'} g^b{}_{b'} (\{A, A^{;a'b'}\}_+ + \{B, B^{;a'b'}\}_+) \right) \\ & - \xi^2 R n^a n^b (\{A, A'\}_+ + \{B, B'\}_+) \\ & + \frac{i}{8} \text{Tr} \left(\mathcal{J} \gamma^{(a} (g^b)_{b'} [\Psi, \Psi^{+;b'}]_- - [\Psi^{;b}, \Psi^{+'}]_- \right) \\ & + \frac{\xi}{16} K \text{Tr} \left(\gamma^{(a} G^{[b]c]} n_c \{ \nabla (A - i\gamma^5 B), (A' - i\gamma^5 B') \}_+ \right) \\ & + \frac{\xi}{16} K g^a{}_{a'} g^b{}_{b'} \text{Tr} \left(\gamma^{(a'} G^{[b']c']} n_{c'} \{ (A - i\gamma^5 B), \nabla' (A' - i\gamma^5 B') \}_+ \right) \\ & + \frac{i}{8} K \text{Tr} \left(\gamma^{(a} \not{n} \gamma^b) \mathcal{J} [\Psi', \Psi^+]_- \right) + \frac{i}{8} K g^b{}_{b'} g^a{}_{a'} \text{Tr} \left(\gamma^{(a'} \not{n} \gamma^{b')} \mathcal{J} [\Psi, \Psi^{+'}]_- \right) \end{aligned} \quad (98)$$

Now set $\xi = 1/4$ and assume $K = R = 0$. The result is

$$\begin{aligned} \tilde{T}^{ab}(x, y) = & \frac{1}{8} g^b{}_{b'} (\{A^{;a}, A^{;b'}\}_+ + \{B^{;a}, B^{;b'}\}_+) + \frac{1}{8} g^a{}_{a'} (\{A^{;b}, A^{;a'}\}_+ + \{B^{;b}, B^{;a'}\}_+) \\ & - \frac{1}{8} \left((\{A^{;ab}, A'\}_+ + \{B^{;ab}, B'\}_+) + g^a{}_{a'} g^b{}_{b'} (\{A, A^{;a'b'}\}_+ + \{B, B^{;a'b'}\}_+) \right) \\ & + \frac{i}{8} \text{Tr} \left(\mathcal{J} \gamma^{(a} (g^b)_{b'} [\Psi, \Psi^{+;b'}]_- - [\Psi^{;b}, \Psi^{+'}]_- \right) \quad \text{iff } \xi = 1/4 \text{ and } K = R = 0 \end{aligned} \quad (99)$$

As an immediate consequence we get

COROLLARY 4.24. *The point separated energy momentum tensor $\tilde{T}^{ab}(x, y)$ with $\xi = 1/4$ agrees with the Definition of the point separated energy momentum tensor T^{ab} of section 3, Eqn. (37), provided $K = R^{ab} = R = 0$.*

This compatibility result will allow us in the next section to calculate the ‘vacuum’ expectation value $\langle \Omega | \tilde{T}^{ab} \Omega \rangle$ explicitly using the results of Christensen [Chr76, Chr78]. Note that we need R^{ab} here, i.e., vanishing scalar curvature is not sufficient. We close this section with the proof of the Theorem:

PROOF OF THEOREM 4.23. For the proof insert the Definitions and use Proposition 4.21. After some lengthly but elementary calculation one obtains:

$$\begin{aligned}
\tilde{T}^{ab}(x, y) = & \frac{1}{4}g^b_{b'}(\{A^{;a}, A^{;b'}\}_+ + \{B^{;a}, B^{;b'}\}_+) + \frac{1}{4}g^a_{a'}(\{A^{;b}, A^{;a'}\}_+ + \{B^{;b}, B^{;a'}\}_+) \\
& - \frac{1}{4}g^{ab}g^c_{c'}(\{A_{;c}, A^{;c'}\}_+ + \{B_{;c}, B^{;c'}\}_+) + \frac{1}{4}g^{ab}m^2(\{A, A'\}_+ + \{B, B'\}_+) \\
& - \frac{\xi}{2}\left((\{A^{;ab}, A'\}_+ + \{B^{;ab}, B'\}_+) + g^a_{a'}g^b_{b'}(\{A, A^{;a'b'}\}_+ + \{B, B^{;a'b'}\}_+)\right) \\
& - \xi(g^{ab}m^2 + \xi Rn^a n^b)(\{A, A'\}_+ + \{B, B'\}_+) \\
& + \xi g^{ab}g^c_{c'}(\{A_{;c}, A^{;c'}\}_+ + \{B_{;c}, B^{;c'}\}_+) - \frac{\xi}{2}g^a_{a'}(\{A^{;b}, A^{;a'}\}_+ + \{B^{;b}, B^{;a'}\}_+) \\
& - \frac{\xi}{2}g^b_{b'}(\{A^{;a}, A^{;b'}\}_+ + \{B^{;a}, B^{;b'}\}_+) \\
& + \frac{i}{8}\text{Tr}\left(\gamma^{(a}\mathcal{J}[\Psi', \Psi^{+;b})]_-\right) + \frac{i}{8}g^a_{a'}g^b_{b'}\text{Tr}\left(\gamma^{(a'}\mathcal{J}[\Psi, \Psi^{+;b')}]_-\right) \\
& + \frac{\xi}{2}i\text{Tr}\left(\gamma^{(a}G^{[b]c]}\mathcal{J}[\Psi', \Psi^{+}_{;c}]_-\right) + \frac{\xi}{2}g^a_{a'}g^b_{b'}i\text{Tr}\left(\gamma^{(a'}G^{[b')c']}\mathcal{J}[\Psi, \Psi^{+}_{;c'}]_-\right) \\
& + \frac{\xi}{2}i\text{Tr}\left(\mathcal{J}\gamma^{(a}G^{[b]c]}\mathcal{J}[\Psi_{;c}, \Psi^{+'}]_-\right) + \frac{\xi}{2}g^a_{a'}g^b_{b'}i\text{Tr}\left(\mathcal{J}\gamma^{(a'}G^{[b')c']}\mathcal{J}[\Psi'_{;c'}, \Psi^{+}]_-\right) \\
& + \frac{\xi}{16}K\text{Tr}\left(\gamma^{(a}G^{[b]c]}n_c\{\mathcal{N}(A - i\gamma^5 B), (A' - i\gamma^5 B')\}_+\right) \\
& + \frac{\xi}{16}Kg^a_{a'}g^b_{b'}\text{Tr}\left(\gamma^{(a'}G^{[b')c']}\mathcal{N}_{c'}\{(A - i\gamma^5 B), \mathcal{N}'(A' - i\gamma^5 B')\}_+\right) \\
& + \frac{i}{8}K\text{Tr}\left(\gamma^{(a}\not{n}\gamma^b)\mathcal{J}[\Psi', \Psi^{+}]_-\right) + \frac{i}{8}Kg^b_{b'}g^a_{a'}\text{Tr}\left(\gamma^{(a'}\not{n}\gamma^{b')}\mathcal{J}[\Psi, \Psi^{+'}]_-\right)
\end{aligned}$$

By applying Lemma C.2 of Appendix C one verifies:

$$\begin{aligned}
\frac{\xi}{2}i\text{Tr}\left(\gamma^{(a}G^{[b]c]}\mathcal{J}[\Psi', \Psi^{+}_{;c}]_-\right) &= -\frac{\xi}{2}i\text{Tr}\left(\mathcal{J}\gamma^{(a}G^{[b]c]}\mathcal{J}[\Psi_{;c}, \Psi^{+'}]_-\right) \\
\frac{\xi}{2}g^a_{a'}g^b_{b'}i\text{Tr}\left(\gamma^{(a'}G^{[b')c']}\mathcal{J}[\Psi, \Psi^{+}_{;c'}]_-\right) &= -\frac{\xi}{2}g^a_{a'}g^b_{b'}i\text{Tr}\left(\mathcal{J}\gamma^{(a'}G^{[b')c']}\mathcal{J}[\Psi'_{;c'}, \Psi^{+}]_-\right)
\end{aligned}$$

and

$$+\frac{i}{8}\text{Tr}\left(\gamma^{(a}\mathcal{J}[\Psi', \Psi^{+;b})]_-\right) = -\frac{i}{8}\text{Tr}\left(\gamma^{(a}\mathcal{J}[\Psi^{;b}, \Psi^{+'}]_-\right).$$

For the first two equalities one should note that $\gamma^{(a}G^{[b]c]} = \frac{1}{4}(2g^{ab}\gamma^c - g^{cb}\gamma^a - g^{ac}\gamma^b)$. Finally use the fact that the γ -matrices are covariantly constant, i.e., $g^a_{a'}\mathcal{J}\gamma^{a'}\mathcal{J} = \gamma^a$. This proves the Theorem. $\square$

4. The ‘vacuum’ expectation value

In this section the results of Christensen’s covariant geodesic point separation [Chr76, Chr78] are used to calculate the ‘vacuum’ expectation value of our point-splitting energy momentum tensor $\tilde{T}^{ab}$. It is emphasized that we suppose the assumption of Definition 4.22, unless otherwise stated, throughout this section, i.e., a vacuum spacetime ($R^{ab} = 0$) with maximal spacelike hypersurface $\mathcal{C}$ ($K = 0$). Let us start with a short introduction to the covariant geodesic point separation: Consider the ‘vacuum’ expectation value of $\tilde{T}^{ab}$. Using the explicit form for $\tilde{T}^{ab}$ (Eqn. (99)), one sees that $\langle \Omega | \tilde{T}^{ab}(x, y) | \Omega \rangle$ depends on the ‘vacuum’ expectation value of the anti-commutator, or commutator respectively, of the basic fields, together with their derivatives, only. Recall that Ω arises from a quasifree locally SUSY product state ω and $\langle \Omega | \{A(x), A(y)\}_+ | \Omega \rangle$, $\langle \Omega | \{B(x), B(y)\}_+ | \Omega \rangle$ and $\langle \Omega | [\Psi(x), \Psi^+(y)]_- | \Omega \rangle$ are the real and imaginary parts respectively of the Hadamard distributions ω^2 introduced in chapter 3 section 4 (Eqn. (44)). Using the same notation as in that section, we denote the corresponding singular kernels by ${}^AG_\epsilon^{T,P}$, ${}^BG_\epsilon^{T,P}$ and $\Psi\mathcal{G}_\epsilon^{T,P}$. We remind the reader of the fact that in the fermionic case $\Psi\mathcal{G}_\epsilon^{T,P}$ is the singular kernel of the auxiliary two point distribution $\tilde{\omega}^2$. Since both scalar kernels are identical, we may drop the indices and write G for the scalar and $\mathcal{G}^B_{A'}$ for the fermionic case. As a result

$\langle \tilde{T}^{ab} \rangle_\omega$ can be written as⁵:

$$\begin{aligned}
\langle \tilde{T}^{ab} \rangle_\omega &= \frac{1}{4} \left(g^b_{b'} G^{;ab'} + g^a_{a'} G^{;a'b} - g^a_{a'} g^b_{b'} G^{;a'b'} \right) \\
&\quad - \frac{1}{8} \text{Tr} \left(\mathcal{J} \gamma^{(a} \gamma^c \left(g^b)_{b'} \mathcal{G}_{;c}{}^{b'} - \mathcal{G}_{;c}{}^{b)} \right) \right) \\
&\quad + \frac{1}{8} \left(g^b_{b'} ({}^A H^{(p)} + {}^B H^{(p)})^{;ab'} + g^a_{a'} ({}^A H^{(p)} + {}^B H^{(p)})^{;a'b} \right. \\
&\quad \left. - ({}^A H^{(p)} + {}^B H^{(p)})^{;ab} - g^a_{a'} g^b_{b'} ({}^A H^{(p)} + {}^B H^{(p)})^{;a'b'} \right) \\
&\quad - \frac{1}{8} \text{Tr} \left(\mathcal{J} \gamma^{(a} \gamma^c \left(g^b)_{b'} \Psi H^{(p)}_{;c}{}^{b'} - \Psi H^{(p)}_{;c}{}^{b)} \right) \right) \\
&=: \langle \tilde{T}^{ab} \rangle_{\omega, \text{sing}} + \langle \tilde{T}^{ab} \rangle_{\omega, \text{finite}},
\end{aligned} \tag{100}$$

i.e., as a sum of two parts $\langle \tilde{T}^{ab} \rangle_{\omega, \text{sing}}$ and $\langle \tilde{T}^{ab} \rangle_{\omega, \text{finite}}$. The first summand is constructed solely with the singular kernels G and $\mathcal{G}$, while the second consists of the remaining regular kernels $H^{(p)}$ and their derivatives only. The label ‘finite’ was chosen for the latter, since the corresponding distribution on $\mathcal{C} \times \mathcal{C}$ can be restricted to the diagonal and the result is a finite *function* $\langle \tilde{T}^{ab} \rangle_{\omega, \text{finite}}$. (Recall that $H^{(p)}$ can be made sufficiently smooth by choosing p large enough. It follows, that the coincidence limit exist.) On the other hand, due to the divergences in both, G and $\mathcal{G}$, the coincidence limit of $\langle \tilde{T}^{ab} \rangle_{\omega, \text{sing}}$ does *not* exist in general, i.e., if we use Eqn. (100), but relax our assumptions on the spacetime and the hypersurface. Renormalization is necessary in this case. It is worth noting that all state dependent information is contained in $\langle \tilde{T}^{ab} \rangle_{\omega, \text{finite}}$, since G and $\mathcal{G}$ are purely geometric objects. We digress for a moment to describe the renormalization prescription according to Adler & al. [ALN77] or Wald [Wal78]: Subtract from G the Hadamard solution G^0 , which is defined by setting the (arbitrary) function H_0 in the expansion of $H^{(p)}$ (see Appendix B) equal to zero and plug in the result into $\tilde{T}^{ab}(x, y)$. Then, since G and G^0 have equal singularities, the coincidence limit $x \rightarrow y$ will exist and, after performing the same operation on $\mathcal{G}$, defines a finite expression for $\langle \tilde{T}^{ab}(x) \rangle_{\omega, \text{sing, ren.}}$. Unfortunately, as is stated in Wald loc.cit., this prescription is not unique, since the length scale, by which σ is measured has not been specified. In the massive case the Compton wave length of the particle can be used to introduce such a length scale and the ambiguity can be resolved. However, for a massless theory the ambiguity remains due to the fact that these models do not possess a natural length scale. Moreover, it was pointed out by Wald loc.cit. that an unambiguous prescription for the renormalized stress tensor can not be given without introducing a new fundamental length scale. Another problem, which one encounters using this prescription is related to the fact that not all functions H_0 lead to *symmetric* Hadamard bisolutions. Namely, $\langle \tilde{T}^{ab}(x) \rangle_{\omega, \text{sing, ren.}}$ fails to be locally conserved in general. To obtain a locally conserved object one must in a last step add the local curvature term $-\frac{1}{32\pi^2} [v_1] g^{ab}$, where $[v_1]$ is the coincidence limit of the second expansion coefficient of $v^{(p)}$ (See Eqn. (111) in Appendix B). Apart from these deficiencies the geodesic point separation method turned out to be a valuable tool for explicit calculations. For some of the results obtained by this method the reader is referred to the books of Birell & Davis [BD82] or Wald [Wal84] respectively and the original literature cited therein. The result of this section is that in our case a renormalization is not necessary (at least for the massive theory), since the divergences of the bosonic and fermionic parts of the model already cancel each other. Unfortunately there remain direction dependent terms, which must be dealt with by an ‘elementary averaging’ procedure, in order to obtain a tensorial, i.e., direction independent, result. To establish this result we have to isolate the divergences of the fermionic and bosonic parts of $\langle \tilde{T}^{ab} \rangle_\omega$ respectively in a covariant way. This can be done using a covariant Taylor expansion for all terms in $\langle \tilde{T}^{ab} \rangle_\omega$, i.e., for G , $\mathcal{G}$ and their derivatives. The latter is a method frequently used in classical general relativity. See for instance DeWitt’s classical book [DeW65]. For the convenience of the reader we describe the basic idea in the next paragraph: Covariant Taylor expansion, as well as the usual Taylor expansion, uses a power series to describe a smooth function near some initial point. For the covariant Taylor expansion one uses the fact that every smooth bi-scalar $K(x, x')$ on a Lorentz manifold, i.e., a smooth function depending on x and x' , for x sufficiently close to x' can be written as a power series in σ and its covariant derivatives up to order q with coefficients $k(x)$ depending on x only, plus a remainder, which is of order $\mathcal{O}(\sigma^{(q+1)})$. The coefficients $t(x)$ are completely determined by the coincidence limits of K and its derivatives at x and one writes

$$K(x, x') \equiv k(x) + k_a(x) \sigma^{;a}(x, x') + \frac{1}{2!} k_{ab} \sigma^{;ab}(x, x') + \dots \mathcal{O}(\sigma^{(q+1)}). \tag{101}$$

The relation between the coefficients and K can be established by repeatedly differentiating both sides of Eqn. (101) with respect to x' and performing the coincidence limit afterwards. The results, as well as the various coincidence limits

⁵Note that the auxiliary two-point distribution $\mathcal{G} + H^{(p)}$ in the fermionic case is *even* in the γ -matrices [DeW65], which implies that $\text{Tr}(\gamma^a (\mathcal{G} + H^{(p)}))$ vanishes.

of σ , which are needed, can be found for instance in [Chr76, Chr78]. Unfortunately we can not expand the kernels G and $\mathcal{G}$ directly according to Eqn. (101), since their coincidence limits do not exist. On the other hand the van Vleck Morette determinant $\Delta^{1/2}$, $v^{(p)}$ and σ are smooth in any convex normal neighborhood and thus *their* covariant Taylor expansions exist to any order. (Note that we do not assume the convergence of the infinite series without remainder). Therefore the only problematic terms are those like $(\sigma^{-1})^{;ab}(x, x')$, which diverge as x' approaches x . In order to expand them, we carry out the differentiation, which yields

$$(\sigma^{-1})^{;ab} = 2\sigma^{-3}\sigma^{;a}\sigma^{;b} - \sigma^{-2}\sigma^{;ab}. \quad (102)$$

We can now expand $\sigma^{;ab}$ and $\sigma^{;a}$ and plug in the result into Eqn. (102). The result, which contains terms proportional to σ^{-2} and $\sigma^{-1/2}$ can be found e.g. in [Chr76]. It should be noted that this expansion is only valid if the points x and x' are not null related, i.e., for $\sigma(x, x') \neq 0$. On the other hand, the distribution $\langle \tilde{T}^{ab}(x, y) \rangle_{\omega, \text{sing}}$ is defined on $\mathcal{C} \times \mathcal{C}$, hence the only singularities occur at the diagonal, while everywhere else the corresponding kernel is smooth. We may therefore expand G , $\mathcal{G}$ and their derivatives by substituting for $\Delta^{1/2}$, $v^{(p)}$, σ^{-1} , $\ln(\sigma)$ and their derivative the corresponding power series, form $\langle \tilde{T}^{ab} \rangle_{\omega, \text{sing}}$ and collect together terms like powers of $\sigma^{;a}$. This tremendous task, which involves much algebra and the calculation of many coincidence limits, has fortunately already been done by Christensen. In his PhD thesis he performed the calculation for the scalar field and extended the result later to fermions ([Chr76] and [Chr78]). He did not use the representation (111) for the Hadamard distributions. Instead he considered a special Hadamard distribution, usually denoted by G_{DS} , which was originally obtained by DeWitt [DeW65] using Schwinger's proper time formalism. The kernel of this DeWitt-Schwinger Hadamard distribution has an asymptotic expansion of the form [Chr76]:

$$\begin{aligned} G_{DS}(x, x') &= \frac{\Delta^{1/2}}{4\pi^2} \left\{ a_0 \left[\frac{1}{\sigma} + m^2 L \left(1 + \frac{1}{4} m^2 \sigma + \dots \right) - \frac{1}{2} m^2 - \frac{5}{16} m^2 \sigma + \dots \right] \right. \\ &\quad - a_1 \left[L \left(1 + \frac{1}{2} m^2 \sigma + \dots \right) - \frac{1}{2} m^2 \sigma \right] \\ &\quad + a_2 \sigma \left[L \left(\frac{1}{2} + \frac{1}{8} m^2 \sigma + \dots \right) - \frac{1}{4} - \dots \right] + \dots \\ &\quad \left. + \frac{1}{2m^2} [a_2 + \dots] + \frac{1}{2m^4} [a_3 + \dots] + \dots \right\}, \end{aligned} \quad (103)$$

where $L = (\gamma + \frac{1}{2} \ln |\frac{1}{2} m^2 \sigma|)$, γ is Euler's constant, $\Delta^{1/2}$ is the van Vleck Morette determinant and the a_n are smooth functions satisfying certain recursion relations with boundary condition $a_0(x, x) = 1$. Note that Eqn. (103) is equally valid for the scalar and fermionic case, but in the former $\Delta^{1/2}$ and a_n are bi-scalars, while in the latter they are bi-spinors. The omitted terms are either of order σ^2 or finite of order $1/m^6$. It was pointed out by Adler, Lieberman and Ng [ALN77], and also by Castagnino and Harari [CH84] that G_{DS} can be identified with a Hadamard series in the DeWitt-Brehme form, i.e., the form given in Appendix B written as an asymptotic series, by setting the undetermined function H_0 to

$$\begin{aligned} {}^{DS}H_0 &= \Delta^{-1/2} \left\{ m^2 (\ln m^2 - \ln 2 + 2\gamma - 1) \right. \\ &\quad \left. - a_1 (2\gamma - \ln 2 + \ln m^2) + \frac{1}{m^2} a_2 + \frac{1}{m^4} a_3 + \dots \right\} \end{aligned} \quad (104)$$

Moreover, the second authors together with Nuñez claimed in [CHN82] that G_{DS} is the unique kernel which covariantly generalizes to curved spacetimes the singularity structure of the Minkowski space Hadamard distribution together with its derivatives with respect to m^2 . Christensen continues by inserting into Eqn. (103) as well as into its derivatives the covariant Taylor expansions for the derivatives of σ , for $\Delta^{1/2}$ and a_n and for their derivatives. He forms the stress energy tensor for a scalar and a Fermi field using an analogon of Eqn. (99), collects terms like powers of $\sigma^{;a}$ and observes quartic (σ^{-2}), quadratic (σ^{-1}), linear ($1/\sigma^{;a}$) and logarithmic ($\ln(m^2 \sigma)$) divergences for both fields. Inserting his results into Eqn. (99), one sees that all divergences cancel on a vacuum solution of the Einstein equations. One ends up with:

$$\begin{aligned} \langle \tilde{T}^{ab}(x, y) \rangle_{DS} &= \frac{1}{4\pi^2} \left(-\frac{1}{192} R^{cdef} R_{cdef} \frac{\sigma^{;a}\sigma^{;b}}{2\sigma} + \frac{1}{3} R^{(ac}{}_{d}{}^e R_{f cge} \frac{\sigma^{;b)}\sigma^{;d}\sigma^{;f}\sigma^{;g}}{(2\sigma)^2} \right. \\ &\quad + \frac{1}{24} R^{cde(a} R_{cdef} \frac{\sigma^{;b)}\sigma^{;f}}{2\sigma} \\ &\quad - \frac{1}{24} [R^{acbd} R_{cedf} + R^{acd}{}_{e} R^b{}_{dcf} + R^{acd}{}_{e} R^b{}_{cdf}] \frac{\sigma^{;e}\sigma^{;f}}{2\sigma} \\ &\quad \left. + \frac{1}{96} R^{cde}{}_{f} R_{cdeg} \frac{\sigma^{;f}\sigma^{;g}}{2\sigma} \left(g^{ab} - 2 \frac{\sigma^{;a}\sigma^{;b}}{2\sigma} \right) \right), \end{aligned} \quad (105)$$

where the subscript DS reminds of the fact that the special Hadamard distribution G_{DS} was used.

REMARK .

- This result is not exact, but of order $1/m^6$. However all missing terms have only finite contributions in the coincidence limit (See [Chr76]).
- The result does not depend on the length scale chosen. A change of length scale ($\sigma \rightarrow \lambda^2 \sigma'$) yields simple factors in front of the quartic, quadratic and linear divergent terms associated to the single fields, but produces a finite ($\ln(\lambda^2 m^2)$) contribution to be added to the logarithmic divergent terms. On the other hand the fact that the divergent terms cancel each other in the situation considered here, implies that those finite terms cancel as well.

It is easy to see that Eqn. (105) remains finite if y approaches x , but unfortunately the result is direction dependent, i.e., it depends on how the points were ‘separated’. To remedy this defect, we use an elementary averaging procedure described by Adler et al. in [ALN77]. It consists in the following replacements:

$$\begin{aligned}\sigma^{;a}\sigma^{;b} &\rightarrow \frac{1}{2}g^{ab}\sigma \\ \sigma^{;a}\sigma^{;b}\sigma^{;c}\sigma^{;d} &\rightarrow \frac{1}{6}\sigma^2(g^{ab}g^{cd} + g^{ac}g^{bd} + g^{ad}g^{cd})\end{aligned}$$

Note that these replacements correspond to a decomposition of a direction dependent expression, e.g., $\sigma^{;a}\sigma^{;b}$, according to irreducible representations of the Lorentz-group, keeping the scalar part, e.g., $\frac{1}{4}g^{ab}\sigma^{;c}\sigma^{;c}$, only. Let us denote the averaged result for $\langle \tilde{T}^{ab} \rangle_{DS}$ by $\langle \tilde{T}^{ab} \rangle_{\overline{DS}}$. Then the coincidence limit is well defined and we obtain as a final result:

$$\begin{aligned}\langle \tilde{T}^{ab} \rangle_{\overline{DS}} &= \frac{1}{4\pi^2} \left(+\frac{1}{288}R^{acde}R^b_{cde} + \frac{1}{288}R^{acde}R^b_{edc} - \frac{1}{96}R^{acde}R^b_{dce} \right). \\ &= 0 \quad \text{if } R^{ab} = 0,\end{aligned}\tag{106}$$

since $R^{acde}(R^b_{cde} + R^b_{dec} + R^b_{ecd}) = 0 = R^{acde}R^b_{cde} + 2R^{acde}R^b_{dec}$ (See Adler et al. [loc. cit.]). It is worth noting that this result already implies the finiteness of $\langle \tilde{T}^{ab} \rangle_\omega$ in the coincidence limit⁶ for every locally SUSY Hadamard product state ω , since all Hadamard distributions differ by a *smooth* function only. Thus for our state ω , which we assume to be characterized by its smooth part $H^{(p)}$:

$$\langle \tilde{T}^{ab} \rangle_\omega = \langle \tilde{T}^{ab} \rangle_{\overline{DS}} + \langle \tilde{T}^{ab} \rangle_{(H^{(p)} - DS H^{(p)})} \equiv \langle \tilde{T}^{ab} \rangle_{\overline{DS}} + \langle \tilde{T}^{ab} \rangle_{H^{(p)}} - \langle \tilde{T}^{ab} \rangle_{DS H^{(p)}}$$

where $\langle \tilde{T}^{ab} \rangle_{H^{(p)}}$ denotes Eqn. (100) with G and $\mathcal{G}$ respectively omitted.

REMARK . DeWitt has shown in [DeW75] that $\langle \tilde{T}^{ab} \rangle_{\overline{DS}}$ can be obtained by functionally differentiating an effective action with respect to the metric. Hence this expression –as is the classical stress-energy tensor– is covariantly conserved by construction. On the other hand for the smooth part $\langle \tilde{T}^{ab} \rangle_{DS H^{(p)}}$ one might encounter $\nabla_a \langle \tilde{T}^{ab} \rangle_{DS H^{(p)}} \neq 0$, since this part does not contain the finite contributions from $\langle \tilde{T}^{ab} \rangle_{\overline{DS}}$. However being an auxiliary expression only, which was introduced to simplify the calculations of $\langle \tilde{T}^{ab} \rangle_\omega$, this property of $\langle \tilde{T}^{ab} \rangle_{DS H^{(p)}}$ does not affect the covariant conservation of $\langle \tilde{T}^{ab} \rangle_\omega$ as a whole.

LEMMA 4.25. Let $H^{(p)} = \sum_{i=0}^p H_i \sigma^i$ denote the smooth⁷ part of the scalar and fermionic Hadamard series respectively. Then

$$\begin{aligned}\langle \tilde{T}^{ab}(x, x) \rangle_{H^{(p)}} &\equiv [\langle \tilde{T}^{ab} \rangle_{H^{(p)}}] \\ &= \frac{1}{4} \left(2[H_0^{;a};^b + 2[H_0^{;b};^a - [H_0]^{;ab} - 4[H_0^{;ab}] - 4[H_1]g^{ab} \right) \\ &\quad - \frac{1}{8} \left\{ \gamma^{(a} \gamma^{c} \left([H_{0;c}]^{;b)} - 2[H_{0;c}^b] - 2[H_1]g_c^b \right) \right\},\end{aligned}$$

where $[\cdot]$ as usual denotes the coincidence limit.

PROOF. Write $H^{(p)}$ in its series expansion:

$$H^{(p)} = \sum_{i=1}^p H_i \sigma^i$$

and note that $\langle \tilde{T}^{ab} \rangle_{H^{(p)}}$ contains second derivatives only. Differentiating $H^{(p)}$ two times yields:

$$H^{(p)}_{;ab} = H_{0;ab} + H_{1;ab}\sigma + H_{1;a}\sigma_{;b} + H_{1;a}\sigma_{;b} + H_{1;a}\sigma_{;b} + O(\sigma).$$

⁶After elementary averaging.

⁷Strictly speaking $H^{(p)}$ is p -times differentiable.

For the Lemma we are only interested in terms which do not vanish in the coincidence limit. These terms are $H_{0;ab}$ and $H_1\sigma_{;ab}$, since it is one of the defining properties of σ that its coincidence limit and that of its covariant derivative vanishes. Thus for the coincidence limit of $\langle \tilde{T}^{ab}(x, y) \rangle_{H(p)}$ one gets:

$$\begin{aligned} [\langle \tilde{T}^{ab} \rangle_{H(p)}] &= 2 \cdot \frac{1}{8} \left([g^b_{c'}] ([H_0^{ac'}] + [H_1][\sigma^{;ac'}]) + [g^a_{c'}] ([H_0^{c'b}] + [H_1][\sigma^{;c'b}]) \right. \\ &\quad \left. - ([H_0^{;ab}] + [H_1][\sigma^{;ab}]) - [g^a_{c'}][g^b_{d'}] ([H_0^{c'd'}] + [H_1][\sigma^{;c'd'}]) \right) \\ &\quad - \frac{1}{8} \text{Tr} \left\{ [\mathcal{J}] \gamma^{(a} \gamma^c ([g^b)_{d'}] ([H_{0;c}{}^{d'} + [H_1][\sigma_{;c}{}^{d'}]) - ([H_{0;c}{}^b] + [H_1][\sigma_{;c}{}^b]) \right\} \\ &= \frac{1}{4} \left(2[H_0^{;a}]^b + 2[H_0^{;b}]^a - [H_0]^{;ab} - 4[H_0^{;ab}] - 4[H_1]g^{ab} \right) \\ &\quad - \frac{1}{8} \left\{ \gamma^{(a} \gamma^c ([H_{0;c}]^b) - 2[H_{0;c}{}^b] - 2[H_1]g_c{}^b \right\} \end{aligned} \quad (107)$$

since $[\sigma^{;ab}] = [\sigma^{;a'b'}] = g^{ab}$, $[\sigma^{;ab'}] = -g^{ab}$ and $[g^b_{c'}] = \delta^b_{c'}$. We used Synge's Theorem ([Chr76]) which states that

$$[H_{a_1, \dots, a_n, b'_1, \dots, b'_m; c'}] = -[H_{a_1, \dots, a_n, b'_1, \dots, b'_m; c}] + [H_{a_1, \dots, a_n, b'_1, \dots, b'_m}]_{;c},$$

for any bi-tensor $H_{a_1, \dots, a_n, b'_1, \dots, b'_m}$ whose coincidence limits are known. $\square$

Now we apply this Lemma to the smooth DeWitt-Schwinger term, ${}^{DS}H(p)$. The necessary coincidence limits for ${}^{DS}H_0$ and its derivatives, as well as for ${}^{DS}H_1$, are collected in Appendix B. The result is:

$$\begin{aligned} \langle \tilde{T}^{ab}(x, x) \rangle_{{}^{DS}H(p)} &= (-1) \left(-\frac{\delta}{90} R^{cdea} R_{cde}{}^b + g^{ab} \left(\frac{2\delta-3}{720} W^2 - \frac{1}{4} m^4 (\delta-1) \right) \right. \\ &\quad \left. - \frac{1}{8} \text{Tr} \left(\gamma^{(a} \gamma^c \left\{ +\frac{\delta}{24} G_{[kd]} G_{[ef]} (R^{kdg}{}_c R^{ef}{}_g{}^b) + R^{kdgb} R^{ef}{}_{gc} \right\} + \frac{\delta}{45} R^{kde}{}_c R_{kde}{}^b \right) \right. \\ &\quad \left. - 2g_c{}^b \left(\frac{2\delta-3}{192} G_{[kd]} G_{[ef]} R^{kdgh} R^{ef}{}_{gh} + \frac{2\delta-3}{720} W^2 - \frac{m^4}{4} (\delta-1) \right) \right\} \\ &= \frac{\delta}{90} R^{cdea} R_{cde}{}^b - g^{ab} \left(\frac{2\delta-3}{720} W^2 - \frac{1}{4} m^4 (\delta-1) \right) \\ &\quad + g^{ab} \left(\frac{2\delta-3}{720} W^2 - \frac{1}{4} m^4 (\delta-1) \right) - g^{ab} W^2 \frac{2\delta-3}{768} - \frac{\delta}{90} R^{cdea} R_{cde}{}^b \\ &\quad - \frac{\delta}{192} \text{Tr} (\gamma^{(a} \gamma^c G_{[kd]} G_{[ef]} (R^{kdg}{}_c R^{ef}{}_g{}^b) + R^{kdgb} R^{ef}{}_{gc}) \\ &= \frac{2\delta-3}{768} W^2 g^{ab} - \frac{\delta}{48} (R^{acde} R^b{}_{ced} + 2R^{acde} R^b{}_{dec}), \end{aligned} \quad (108)$$

where $W^2 = R^{abcde} R_{abcde}$, $\delta = \ln m^2 - \ln 2 + 2\gamma$ and the trace formulas of Appendix C were used. As an example let us now consider the Schwarzschild and Kerr solution of Einstein's equation.

5. Two examples: The Schwarzschild and Kerr solution

5.1. The Schwarzschild solution. One of the simplest relativistic star models is the Schwarzschild solution of Einstein's equation. It is an exact vacuum solution which describes the geometry outside a static spherical symmetric relativistic system. Moreover, it has been shown implicitly by Bartnik, Chruściel and Murchdha [BCO90] that every point in this spacetime possesses a maximal spacelike hypersurface passing through it. With these two properties the Schwarzschild solution provides a good background for an application of our method. We are going to calculate the 'expectation value' of the stress energy tensor of a locally SUSY model with respect to the DeWitt Schwinger pseudostate ($\langle \tilde{T}^{ab} \rangle_{\overline{DS}}$) together with the corresponding smooth part $\langle \tilde{T}^{ab} \rangle_{{}^{DS}H(p)}$. The actual computations are performed using the Maple V Symbolic Computation Program by Waterloo Maple Software [C+91b, C+91a]. (A commented listing and sample run can be found in the Appendix D. Using the Schwarzschild coordinates t, r, θ, ϕ , the line element of this spacetime is known to be of the form:

$$ds^2 = \left(1 - \frac{2M}{r^2}\right) dt^2 - \left(1 - \frac{2M}{r^2}\right)^{-1} dr^2 - r^2 d\theta^2 - r^2 \sin^2 \theta d\phi^2,$$

where M can be interpreted as the total mass of the system as measured from infinity. In these coordinates the only non vanishing components of the Riemann tensor are

$$\begin{aligned} R_{0101} &= \frac{2M}{r^3} & R_{0202} &= -\frac{(r-2M)M}{r^2} & R_{0303} &= \frac{(r-2M)\sin^2\theta}{r^2} \\ R_{1212} &= \frac{\sin^2\theta}{r-2M} & R_{2323} &= -2rM\sin^2\theta \end{aligned}$$

and those which follow from symmetry. Contracting the Riemann tensor with itself yields:

$$R^{abcd}R_{abcd} = 48\frac{M^2}{r^6},$$

the Kretschman invariant. Inserting these results in Eqn. (106) we obtain:

$$\langle \tilde{T}^{ab}(x) \rangle_{DS} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

a result which clearly is covariantly constant. For the smooth part $\langle \tilde{T}^{ab} \rangle_{DSH(p)}$ the result is more complicated:

$$\begin{aligned} \langle \tilde{T}^{ab} \rangle_{DSH(p)} &= \left(\frac{1}{16} \frac{M^2 (10r\delta - 3r - 16\delta M)}{r^5 (r-2M)^2}, -\frac{1}{16} \frac{(2\delta-3)M^2}{r^6}, \right. \\ &\quad \left. -\frac{1}{16} \frac{(2\delta-3)M^2}{r^7 (r-2M)}, \frac{1}{16} \frac{(2\delta-3)M^2}{r^7 (-r+r\cos(\theta)^2+2M-2M\cos(\theta)^2)} \right) \\ &\quad \left(-\frac{1}{16} \frac{(2\delta-3)M^2}{r^6}, -\frac{1}{16} \frac{(6r\delta+4\delta M+3r-6M)M^2 (r-2M)}{r^8} \right. \\ &\quad \left. , \frac{1}{16} \frac{(r-2M)(2\delta-3)M^2}{r^9}, -\frac{1}{16} \frac{M^2 (2r\delta-4\delta M-3r+6M)}{r^9 (-1+\cos(\theta)^2)} \right) \\ &\quad \left(-\frac{1}{16} \frac{(2\delta-3)M^2}{r^7 (r-2M)}, \frac{1}{16} \frac{(r-2M)(2\delta-3)M^2}{r^9}, \right. \\ &\quad \left. -\frac{1}{16} \frac{M^2 (-2\delta+3+8\delta r^2)}{r^{10}}, -\frac{1}{16} \frac{(2\delta-3)M^2}{r^{10} (-1+\cos(\theta)^2)} \right) \\ &\quad \left(\frac{1}{16} \frac{(2\delta-3)M^2}{r^7 (-r+r\cos(\theta)^2+2M-2M\cos(\theta)^2)}, \right. \\ &\quad \left. -\frac{1}{16} \frac{M^2 (2r\delta-4\delta M-3r+6M)}{r^9 (-1+\cos(\theta)^2)}, -\frac{1}{16} \frac{(2\delta-3)M^2}{r^{10} (-1+\cos(\theta)^2)}, \right. \\ &\quad \left. \frac{1}{16} \frac{M^2 (2\delta-3-8\delta r^2+8\delta r^2\cos(\theta)^2)}{r^{10} (1-2\cos(\theta)^2+\cos(\theta)^4)} \right) \end{aligned}$$

REMARK . It is explicitly shown in the Appendix that this expression is *not* covariantly constant. This only expresses Wald's observation [Wal78] that the divergent part in the Adler, Lieberman and Ng renormalization prescription is not covariantly constant in general.

5.2. The Kerr solution. The Kerr solution of the Einstein equation describes the geometry near a rotating mass. The results of Bartnik and al. loc. cit. apply to this Lorentz manifold, too, i.e., every point in the Kerr spacetime possesses a maximal spacelike hypersurface passing through it. We are going to calculate the 'expectation value' of the stress energy tensor of our model with respect to the DeWitt Schwinger pseudostate ($\langle \tilde{T}^{ab} \rangle_{DS}$), by using a slightly modified version of the Maple V program, which was used above for the Schwarzschild case. The reader is referred to Appendix D for a commented listing and a sample run. It should be noted that this example is already complex enough so that the smooth part, $\langle \tilde{T}^{ab} \rangle_{DSH(p)}$, can't be handled by the program in reasonable times. Using the Schwarzschild coordinates t, r, θ, ϕ , the line element of the Kerr spacetime is known to be of the form:

$$ds^2 = dt^2 - (r^2 + a^2) \sin^2\theta d\phi^2 - \frac{2Mr}{\rho^2} (dt + a \sin^2\theta d\phi)^2 - \rho^2 (d\theta^2 + \frac{dr^2}{r^2 - 2Mr + a^2}),$$

where $\rho^2 := r^2 + a^2 \cos^2 \theta$. The parameter a corresponds to the angular momentum of the rotating system. The calculation yields the manifestly conserved result:

$$\langle \tilde{T}^{ab}(x) \rangle_{DS} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Discussion and outlook

For an analogon of the free Wess-Zumino model on a vacuum solution of the Einstein equations, a relation between conserved ‘supercurrents’ and the point-separated stress energy tensor, similar to the one on Minkowski space, was established. The assumption on the scalar curvature of the underlying manifold, i.e., $R^{ab} = R = 0$, is necessary to yield a *conserved* current. This in turn ensures –at least in the flat case– that the corresponding stress tensor is covariantly constant. On the other hand if one relaxes this assumption, our definition of the stress tensor, Eqn. (97), would still make sense, yielding Eqn. (98). However, the result in terms of the basic fields would *not* reproduce the true improved stress tensor, Eqn.(37), i.e., Eqn. (98) and Eqn. (37) differ by terms which vanish on vacuum solutions of the Einstein equations only. These difference terms prevent Eqn. (98) from being covariantly conserved. This means that Eqn. (98) can not be interpreted as a physically sensible stress tensor in this case. Moreover, the divergent parts of the true improved stress energy tensor, according to Christensen’s calculations, do not cancel if $R^{ab} \neq 0$.⁸ Both facts considerable limit the use of the Bose-Fermi cancellation mechanism in the framework of quantum field theory on curved spacetime, since a ‘fine tuning’ of the underlying geometry is necessary. Nevertheless the examples of chapter 4 section 5 show that the Schwarzschild and Kerr solutions together with the DeWitt-Schwinger pseudostate⁹ of a locally SUSY model are self consistent solutions of the semiclassical Einstein equation. A more realistic state on the first manifold is the Hartle-Hawking vacuum [HH76], which represents a black hole of mass M in unstable thermal equilibrium with a bath of black body radiation of local temperature $T_{\text{loc}} = (8\pi M)^{-1}(1 - 2M/r)^{-1/2}$. In a future work this state and its generalization to Fermi fields could be used to calculate the corresponding expectation value of our stress tensor. One could further investigate whether the Hawking radiation associated to this state gets modified by the Bose-Fermi symmetry of the model. This might also answer the question whether the Hartle Hawking vacuum gives a new self consistent solution of the semiclassical Einstein equation. On the other hand, independent of the result, the ‘fine tuning’ problem mentioned above suggests that such a solution, as well as the one presented in this work, is not stable, i.e., every small perturbation of the metric which changes the Ricci tensor immediately yields a divergent result for the stress tensor, making the semiclassical Einstein equation ill defined. It should be noted that this statement is qualitatively different compared to similar statements for the semiclassical Einstein equation with a *renormalized* stress tensor on the r.h.s. In the latter case the equation remains well defined if the metric changes and the usual phase space methods can be used to check stability. (See [CGNP86] for partial results on the stability of the DeSitter universe with a neutral scalar field.) The second assumption necessary to reproduce the true improved stress energy tensor, is the setting of the coupling constant ξ to $1/4$. This setting is motivated by the observation that the ‘squared’ Dirac operator when applied to spinors equals the SUSY coupled Klein-Gordon operator times the identity. Moreover, according to Christensen’s calculations, any other setting of ξ would give a divergent result for the expectation value of the true improved stress energy tensor. For the physical implications of this setting one should note that the field equations on a Ricci flat manifold are independent of ξ . Thus on these manifolds SUSY coupling yields –via Dimock’s construction– the *same* algebras of local observables as the usual minimal ($\xi = 0$) or conformal ($\xi = 1/6$) setting. It follows that the physics does not depend on the choice of ξ , which is satisfactory. On the other hand the parameter ξ enters into the classical definition of the improved stress energy tensor in a non trivial way, since in order to obtain it, the Lagrangian is varied with respect to the metric, i.e., the variations are not restricted to the Ricci flat case. This means that from a Lagrangian point of view, SUSY coupling may be interpreted as a renormalization: For instance, in the minimal ($\xi = 0$) setting, it consists in adding the counterterm $+(m^2 + \frac{1}{4}R)^{1/2} - m$ to the scalar masses. Obviously an analogous statement holds for the conformal setting. Recall that the latter would be appropriate for a massless model. Unfortunately for such models, the DeWitt-Schwinger approach can not be used a priori, since it contains a power series expansion in $1/m$. On the other hand Christensen showed in [Chr78] that all his calculations remain valid in the massless limit, if one sets $m = 0$ everywhere except in the logarithmic divergent term. There one has to substitute the mass m by some ‘cutoff’ mass μ . In our setting the logarithmic divergent parts cancel each other and we could remove the cutoff. Moreover, our final result for the DeWitt-Schwinger pseudostate (Eqn. (106)) is independent of m , which means that it is valid in the massless limit without any modifications. The reader should recall that one does not have a natural length scale in the massless case, which would, if renormalization were necessary, result in an ambiguity of the stress tensor. It is worth

⁸This result can also be verified using the Hadamard condition of Kay and Wald instead of Christensen’s DeWitt-Schwinger ansatz [Köh93].

⁹It is not clear whether the DeWitt-Schwinger prescription leads to a *state* in our case, i.e., whether positivity holds.

noting that in our model, due to its Bose-Fermi symmetry, this ambiguity is not present (compare chapter 4 section 4). For a successful cancellation of the divergencies, as a third assumption, equal masses for the three fields contained in the model were needed. As is well known from experiments, this assumption is strongly violated in the real world. For supersymmetric models on Minkowski space, which do face the same problem, the best way to solve this deficiency seems to be using a mechanism of spontaneous symmetry breaking. However as far as the author knows, such a mechanism has not yet been implemented into a quantum field theory on a curved spacetime. This might be due to the fact that it is up to now very difficult to do perturbation theory on manifolds. The study of more realistic models should therefore presumably be done on a simple spacetime first, for instance Schwarzschild, since the corresponding equations of motion are too complicated otherwise. It is worth noting that in our model, even for different masses, the worst divergencies of the point separated stress energy tensor cancel each other, as long as the metric remains Ricci flat, i.e., $R^{ab} = 0$. According to Christensen, we are left with a logarithmically divergent part only. Its *numerical* contribution to the ‘vacuum expectation value’ of the stress tensor is expected to be small, even if the ‘separated’ points almost coincide, i.e., if their distance is comparable to the Planck length. Recall that this distance is believed to be the border of validity of our semiclassical approximation, meaning that smaller distances presumably do not make sense in the framework. Thus there should exist a regime, where the logarithmic divergent part can be neglected in explicit calculations. Up to now we only considered the expectation value of the stress tensor in some state. For a supersymmetric model on flat space, the stress tensor itself, due to its relation to the conserved supercurrent and the supercharge, can be defined as an operator valued distribution on the vacuum Hilbertspace. Unfortunately one can not apply the same line of argument to a locally SUSY model on a general Ricci flat manifold. Although it is true that the latter possesses a conserved ‘supercurrent’, the definition of a ‘supercharge’ as an *operator* on some GNS-Hilbertspace failed, mainly because on a general manifold there do not exist covariantly constant classical spinor fields¹⁰. Without such a constant field, the integral of the ‘supercurrent’, which is necessary for obtaining a charge is not defined. In order to integrate the current, one parallel transports its spinor index to some reference point. The result is a bispinor-tensor, which fails to be conserved. This in turn implies, that no charge *operator* on the GNS-Hilbertspace of any Hadamard state exists (see Theorem 6.2.10 of [Lop91]). On the other hand the existence of the stress energy tensor for a supersymmetric model on Minkowski space as an operator valued distribution can equally well be established using Wick products of the basic fields together with the fact that the vacuum expectation value of the point separated stress tensor vanishes in the coincidence limit. Namely, using the following definition for the Wick ordered stress energy tensor:

$$: T^{ab} : (x) := \lim_{x_1 \rightarrow x} (T^{ab}(x, x_1) - \langle 0 | T^{ab}(x, x_1) | 0 \rangle), \quad (109)$$

where $: \cdot :$ denotes Wick ordering and $|0\rangle$ is the Minkowski vacuum, shows that

$$T^{ab}(x) := \lim_{x \rightarrow x} T^{ab}(x, x_1)$$

is a well defined operator valued distribution that differs from $: T^{ab} :$ by a finite function times the unit operator. Note that Eqn. (109) is the usual definition for Wick ordering (compare e.g. [SW64]), since T^{ab} is quadratic in the basic fields and that the coincidence limit

$$\lim_{x_1 \rightarrow x} \langle 0 | T^{ab}(x, x_1) | 0 \rangle$$

exists. Thus showing that Wick products of our basic fields on a manifold with respect to some Hadamard state can be defined as new Wightman fields would immediately yield a stress tensor as an operator valued distribution on the corresponding GNS-Hilbertspace after elementary averaging. Let us sketch the main idea for the proof of the first statement. Let A denote a free scalar Wightman field on the GNS-Hilbertspace $(\mathcal{H}, D, A, \Omega)$ of a quasifree Hadamard state. The kernel

$$: A^2 : (x, y) := A(x) \cdot A(y) - \langle \Omega | A(x) \cdot A(y) | \Omega \rangle$$

yields a well defined operator valued distribution on $\mathcal{H}$. Its restriction to the diagonal is the definition of the Wick square of A , denoted by $: A^2 : (x)$. To show that $: A^2 : (x, y)$ admits such a restriction, one proceeds as in the proof of Proposition 4.1, i.e., one considers

$$\| \cdots A(z_1) \cdots : A^2 : (x_1, y_1) \cdots A(z_l) \cdots : A^2 : (x_m, y_m) \cdots A(z_n) \Omega \| \quad (110)$$

and asks, whether the restriction of Eqn. (110) to the diagonal $x_1 = y_1, \dots, x_m = y_m$ exists. Note that Eqn. (110) splits into sums of products of two-point distributions, since Ω arises from a quasifree state. Fortunately all terms containing two-point distributions of the form $\langle \Omega | A(x_i) A(y_i) | \Omega \rangle$ occur twice thanks to the definition of $: A^2 : (x, y)$ and cancel each other. For the remaining terms, for instance

$$\cdots \langle \Omega | A(x_i) A(x_j) | \Omega \rangle \cdots \langle \Omega | A(y_i) A(y_j) | \Omega \rangle \cdots,$$

¹⁰It would be interesting to classify all Ricci flat spacetimes, which possesses constant spinors, i.e., which are Kähler manifolds in four dimensions (See [BT87]).

setting $x_i = y_i$ yields products of two-point distributions, which are well defined, since the state Ω satisfies the new general wave front set spectrum condition (Definition 3.29) by assumption (compare chapter 3 section 6). We may therefore use the sequential continuity of the restriction mapping to establish that Eqn. (110) can be restricted to the diagonal. For the exact proof of this statement and its generalization to arbitrary Wick powers, we refer the reader to [BFK94], which is in preparation. Note that the argument in fact uses only that the state under consideration is quasifree and satisfies the general wave front set spectrum condition (GWSSC). Hence the result extends to the other basic fields of our local SUSY model. On the other hand the GWSSC would be of limited interest without the methods of the microlocal analysis developed by Hörmander and Duistermaat [Hör71, DH72] which were extensively used in this work. Their mathematical tools together with the wave front set characterization of globally Hadamard states [Rad92] and our GWSSC might turn out to be very powerful for various constructive problems of quantum field theory on curved spacetime. For example the method developed in chapter 4 section 1 for the ‘supercurrent’ was successfully used to construct other new Wightman fields on globally hyperbolic manifolds [Köh94]. It was already mentioned above that its generalization to Wick powers of free fields is in preparation [BFK94]. Even the formulation of a perturbation theory on manifolds in terms of Wightman fields seems possible if one uses the new GWSSC. Work in this direction is in progress. As a possible application of the latter one could consider quantum electrodynamics on globally hyperbolic manifolds and try to calculate vacuum polarization effects in strong gravitational fields, e.g., corrections to the life time of excited atoms near massive objects. The new GWSSC might be even useful for a better understanding of the usual spectrum condition on Minkowski space. First results in this direction, concerning the compatibility of both conditions can already be found in Radzikowski’s thesis, but the feeling is that there are still many deep relations to discover [Rad]. Another thing that should be done is to clarify the relation between the new GWSSC and the scaling limit criterion of Fredenhagen and Haag [FH90]. In connection with the latter research is done on an *algebraic* characterization of wave front sets associated to states.

ACKNOWLEDGEMENT . It is a pleasure to thank my supervisor K. Fredenhagen for his guidance and constant support. I am indebted to him for many insights I received during the course of this work. His valuable comments and remarks helped me many times to see problems clearer and provided substantial motivations. A better way of supervising a thesis is hard to imagine. This work also profited from helpful and stimulating conversations with D. Buchholz, G. Mack and H. Nicolai. It would have been impossible without M. Radzikowski, who generously made the results of his PhD thesis available for me prior to publication. My friends and colleagues contributed through discussions on topics in general relativity and functional analysis. In particular I would like to mention W. Junker, S. Ohlmeyer and R. Brunetti. Special thanks are due to R. Verch for many helpful hints and to B. Rosenow for a careful reading of the manuscript. The financial support given by the ‘Studienstiftung des Deutschen Volkes’ and the ‘DFG’ as part of the ‘Graduiertenkolleg für theoretische Elementarteilchenphysik’ and as part of the project ‘Lokale Quantenphysik und Gravitation’ is gratefully acknowledged.

APPENDIX A

Algebraic topology

For the convenience of the reader this appendix contains some basic terminology of algebraic topology. A well written and quite short introduction into this subject can be found in the book by R. S. Ward and R. O. Wells [WW90]. The book by M. Nakahara [Nak90] is well suited for this purpose, too. We will restrict ourselves here to the definition and properties of *homology groups*, since homotopy groups, despite of their close relation to the former, are not necessary for the understanding of Theorem 2.20. Algebraic topology is a branch of mathematics which tries to characterize topological spaces by their topological invariants. One of the oldest and maybe also most useful topological invariant is the Euler characteristic $\chi(X)$ of polyeders X . It is given by

$$\chi(X) := (\text{number of vertices}) - (\text{number of edges}) + (\text{number of faces})$$

Homology groups are roughly speaking generalizations of this characteristic. To make this statement more precise we need some basic terminology.

1. Basic Terminology

DEFINITION A.1. The set

$$\Delta^k := \{(x^1, \dots, x^k) \in \mathbb{R}^k \mid x^i \geq 0, x^1 + \dots + x^k \leq 1\}$$

is called *standard simplex*.

Note that Δ^k is a point for $k = 0$, a line for $k = 1$, a triangle for $k = 2$, $\dots$.

DEFINITION A.2. Let M denote a topological space. A continuous mapping

$$\sigma: \Delta^k \rightarrow M$$

is called a (*singular*) k -simplex in M .

One may think of every k -simplex as being generated by a convex linear combination of k points $P_0, \dots, P_{k-1} \in \mathbb{R}^k$. Let $\mathcal{R}$ be a commutative ring.

DEFINITION A.3. The set

$$S_k(M, \mathcal{R}) := \{\sigma \mid \sigma = \sum_{\substack{\mu \\ \text{finite}}} a^\mu \sigma_\mu, a^\mu \in \mathcal{R}, \sigma_\mu \text{ singular } \mu\text{-simplex}\}$$

is called the *module* of singular k -chains of M with dimension k .

Note that the sum is assumed to denote formal linear combination.

DEFINITION A.4. The i -th face F_k^i of a (standard) k -simplex is given by the mapping

$$F_k^i: \Delta^k \rightarrow \Delta^{k-1} := \text{Span}(P_0, \dots, \hat{P}_i, \dots, P_k),$$

where all symbols under a $\hat{}$ are omitted.

Let σ be a singular k -simplex. Its faces are given by the composition with this mapping.

DEFINITION A.5. To every k -simplex we associate a $(k-1)$ -simplex $\partial\sigma$ called the *boundary* of σ by

$$(\partial\sigma) := \sum_{i=0}^k (-1)^i \sigma \circ F_k^i.$$

We remark that this boundary operator ∂ is null potent and can be extended canonically to singular k -chains.

DEFINITION A.6. A singular k -chain $\sigma \in S_k(M, \mathcal{R})$ is called a *cycle* if and only if $\partial\sigma = 0$ and a *boundary* if and only if there exists a singular $(k-1)$ -chain $\sigma' \in S_k(M, \mathcal{R})$ such that $\sigma = \partial\sigma'$.

In the homology group all cycles which differ by a boundary are identified. More precisely:

DEFINITION A.7. Let

$$S_k(M, \mathcal{R}) \supset Z_k(M, \mathcal{R}) = \{\text{Set of all } k\text{-cycles}\}$$

and

$$S_k(M, \mathcal{R}) \supset B_k(M, \mathcal{R}) = \{\text{Set of all } k\text{-boundaries}\},$$

then the k -th homology group of M over the ring $\mathcal{R}$ is given by the quotient of cycles and boundaries

$$H_k(M, \mathcal{R}) := Z_k(M, \mathcal{R})/B_k(M, \mathcal{R}).$$

Using objects which are dual to the simplices *cohomology groups* are defined analogously.

DEFINITION A.8. A homomorphism c from $S_k(M, \mathcal{R})$ into $\mathbb{R}$ is called (*singular*) k -*cochain*. The set of all (formal) linear combinations of such cochains will be denoted by $S^k(M, \mathcal{R})$.

DEFINITION A.9. A *coboundary operator* $\delta: S^k \rightarrow S^{k+1}$ is given by transposition of ∂ with respect of the dual pairing.

EXAMPLE A.10. Let ω be a k -form on a manifold M , $\sigma \in S_k(M, \mathbb{R})$ a singular k -chain. Define

$$c(\sigma) := \int_{\sigma} \omega = a^{\mu} \int_{\sigma_{\mu}} \omega = a^{\mu} \int_{\Delta^k} \sigma_{\mu}^*(\omega),$$

where a^{μ} are the coefficients of $\sigma = a^{\mu} \sigma_{\mu}$. By Stokes Theorem we find on the other hand,

$$c(\sigma) = \int_{\partial\sigma} d\omega = a^{\mu} \int_{\Delta^{k-1}} (\partial\sigma_{\mu})^*(d\omega)$$

and conclude that the coboundary operator is given by the exterior derivative in this case.

Cocycles and coboundaries are defined in analogy to cycles and boundaries respectively. The corresponding sets are denoted by $Z^k(M, \mathcal{R})$ and $B^k(M, \mathcal{R})$ respectively.

DEFINITION A.11. The quotient

$$H^k(M, \mathcal{R}) := Z^k(M, \mathcal{R})/B^k(M, \mathcal{R})$$

is called the k -th *cohomology group* of M over the ring $\mathcal{R}$.

For the Definition of Čech-cohomology groups we need some sheaf theory.

2. Sheaf theory

Sheafs were introduced by J. Leray in the late forties of this century during his research on hyperbolic differential equations. Twenty years later they became a valuable tool in the algebraic topology and algebraic geometry. The main reason for this is presumably the fact that sheafs can be used for non topological problems too: Every sheaf contains local *geometrical* information; sheaf cohomology is roughly speaking a tool that relates local sections of sheafs to global ones. Let M be a topological space. A ‘presheaf’ S of abelian groups¹ on M is an assignment of open subsets U of M to an abelian group $S(U)$

$$U \rightarrow S(U),$$

satisfying the following supplementary condition: For all subsets $V \subset U \subset M$ there exists a restriction homomorphism

$$r_V^U: S(U) \rightarrow S(V)$$

with the property that for all $W \subset V \subset U$

$$r_V^U \circ r_W^V = r_W^U \quad \text{and} \quad r_U^U = 1$$

REMARK . A bundle for example consists of a family of geometrical objects characterized by points (fibers). The corresponding objects in a presheaf are characterized by open subsets.

DEFINITION A.12. A *sheaf* is a presheaf satisfying two additional properties:

- Let (U_{α}) be an open covering of M , $s_{\alpha} \in S(U_{\alpha})$ and

$$\forall \alpha, \beta: r_{U_{\alpha} \cap U_{\beta}}^{U_{\alpha}}(s_{\alpha}) = r_{U_{\alpha} \cap U_{\beta}}^{U_{\beta}}(s_{\beta}),$$

then there exists a $s \in S(U)$, such that all s_{α} are restrictions of s :

$$\forall \alpha: r_{U_{\alpha}}^U(s) = s_{\alpha}.$$

¹In chapter 2 section 3 these groups were identical to $\mathbb{Z}_2$.

- s is unique, i.e., let (U_α) and U as above be given and let $s, s' \in S(U)$, such that $r_{U_\alpha}^U(s) = r_{U_\beta}^U(s')$ is satisfied then $s \equiv s'$.

3. Sheaf cohomology

For the Definition of a sheaf cohomology we will proceed similarly as in section 1. Let M denote a topological space and let (U_α) be an open covering of M . A k -simplex is an ordered set of $(k + 1)$ open subsets of (U_α) with non void intersections.

DEFINITION A.13. Let $\sigma = (U_0, \dots, U_k)$ be a $(k + 1)$ -simplex. Its *support* is defined by

$$|\sigma| := U_0 \cap \dots \cap U_k.$$

DEFINITION A.14. A k -cochain of (U_α) is a continuous mapping

$$f: \sigma \mapsto S(|\sigma|).$$

The set of all k -cochains will be denoted by $S^k(U_\alpha, S)$.

Note that $S^k(U_\alpha, S)$ is an abelian group under pointwise composition.

DEFINITION A.15. A *coboundary operator* δ is defined by:

$$\begin{aligned} \delta: S^k(U_\alpha, S) &\rightarrow S^{k+1}(U_\alpha, S) \\ f &\mapsto \delta f: \sigma \mapsto \prod_{i=0}^{k+1} \left(r_{\sigma}^{|\sigma_i|} f(\sigma_i) \right)^{(-1)^i}, \end{aligned}$$

where $\sigma_i := (U_0, \dots, \hat{U}_i, \dots, U_{k+1})$ and r denotes the restriction mapping.

It is easy to see that this operator is null potent. We denote by $Z^k(U_\alpha, S)$ and $B^k(U_\alpha, S)$ the set of all cocycles and coboundaries respectively.

REMARK . In chapter 2 section 3 we considered the special case $S(U) \equiv S \equiv \mathbb{Z}_2$. The two elements of this abelian group are usually denoted by $+1$ and -1 and their group product is the usual multiplication. Moreover, in this special case every element is its own inverse, such that the coboundary operator defined in chapter 2 section 3 is just a special case of Definition A.15 above.

DEFINITION A.16. The quotient

$$H^k(U_\alpha, S) := Z^k(U_\alpha, S) / B^k(U_\alpha, S)$$

is called k -th Čech-cohomology group of M with coefficients in the sheaf S with respect to the covering (U_α) .

In a last step the construction will be made independent of the covering using an inductive limit procedure. Let $(I, \leq)$ be a partial ordered set, $(S_\alpha)_{\alpha \in I}$ a family of modules indexed by I and let

$$f_\beta^\alpha: S_\alpha \rightarrow S_\beta \quad \forall \alpha, \beta \in I$$

be a homomorphism with the property

$$f_\alpha^\alpha = \mathbf{1}$$

and

$$f_\beta^\alpha \circ f_\gamma^\beta = f_\gamma^\alpha, \quad \text{if } \alpha \geq \beta \geq \gamma.$$

DEFINITION A.17. The *inductive limit* of $(S_\alpha)_{\alpha \in I}$ denoted by

$$S = \lim_{\alpha \in I} \text{ind} S_\alpha$$

are the equivalence classes in the disjoint union

$$\hat{S} = \bigcup_{\alpha \in I} S_\alpha$$

with respect to the equivalence relation

$$\hat{S} \ni x_\alpha \sim x_\beta \in \hat{S} : \Leftrightarrow \exists \gamma \text{ such that } \alpha \geq \gamma, \beta \geq \gamma \text{ and } f_\gamma^\alpha(x_\alpha) = f_\gamma^\beta(x_\beta)$$

DEFINITION A.18. The k -th cohomology group of M with coefficients in S is the inductive limit of $H^k(U_\alpha, S)$, where the partial ordering is given by the refinement of coverings.

$$H^k(M, S) := \lim_{U_\alpha} \text{ind} (Z^k(U_\alpha, S) / B^k(U_\alpha, S))$$

APPENDIX B

The Hadamard recursion relations

For the purpose of proving existence and uniqueness results for elliptic and hyperbolic differential operators on manifolds, Hadamard gave 1923 a prescription for constructing bi-distributional solutions for the corresponding differential equations [Had23]. For the Klein-Gordon operator with variable mass $(\square + m^2(x))$ in four dimensions his algorithm is motivated by writing

$$W^{T,p}(x, x') = \lim_{\epsilon \rightarrow 0} \frac{1}{(2\pi)^2} \left(\frac{\Delta^{1/2}(x, x')}{\sigma(x, x') + 2i\epsilon t + \epsilon^2} + v^{(p)}(x, x') \ln[\sigma(x, x') + 2i\epsilon t + \epsilon^2] + H^{(p)}(x, x') \right) \quad (111)$$

with

$$v^{(p)} = \sum_{i=1}^p v_i(x, x') \sigma^i \quad (112)$$

and

$$H^{(p)} = \sum_{i=1}^p H_i(x, x') \sigma^i \quad (113)$$

where $\sigma(x, x')$ is the square of the geodesic distance, $\Delta^{1/2}(x, x') := g^{-1/2}(x) \det(\sigma_{;ab}(x, x')) g^{-1/2}(x')$ is the van Vleck Morette determinant and $t = T(x) - T(x')$ is some global time function. A semicolon denotes covariant differentiation and σ^i abbreviates $\sigma^{;i}$. The normalization factor $(2\pi)^{-1}$ is chosen so as to agree with the two-point distribution for the scalar field on Minkowski space. One assumes $v^{(p)}$ to be smooth and $H^{(p)}$ to be at least p -times differentiable. To proceed one substitutes the ansatz (111) into the Klein-Gordon equation and equates the coefficients of the explicitly appearing powers of σ and $\ln(\sigma)$ to zero. This yields to a family of ordinary differential equations for $v^{(p)}$ and $H^{(p)}$, which can be written recursively as:

$$0 = (n+1)(n+2)v_{n+1} + (n+1)v_{n+1;a}\sigma^{;a} - (n+1)v_{n+1}\Delta^{-1/2}\Delta^{1/2}_{;a}\sigma^{;a} + \frac{1}{2} \left(\square + m^2 + \frac{1}{4}R \right) v_n \quad (114)$$

$$0 = (n+1)(n+2)H_{n+1} + (n+1)H_{n+1;a}\sigma^{;a} - (n+1)H_{n+1}\Delta^{-1/2}\Delta^{1/2}_{;a}\sigma^{;a} + \frac{1}{2} \left(\square - \frac{1}{4}R + m^2 \right) H_n + (2n+3)v_{n+1} + v_{n+1;\mu}\sigma^{;\mu} - \Delta^{-1/2}\Delta^{1/2}_{;\mu}\sigma^{;\mu}v_{n+1} \quad (115)$$

together with the boundary condition

$$0 = v_{0;a} + \left(1 - \Delta^{-1/2}\Delta^{1/2}_{;b}\sigma^{;b} \right) v_0 + \frac{1}{2} \left(\square + m^2 + \frac{1}{4}R \right) \Delta^{1/2}, \quad (116)$$

provided $m^2(x) \equiv \frac{1}{4}R + m^2$. These relations are called *Hadamard recursion relations* (See also [DeW65]). The boundary condition Eqn. (116) completely fixes $v^{(p)}$, while $H^{(p)}$ remains arbitrary. However the latter gets uniquely fixed if H_0 is specified. It should be noted that $\Delta^{1/2}$ and $v^{(p)}$ are symmetric and hence solve similar equations when $(\square + m^2(x))$ is replaced by $(\square_{x'} + m^2(x'))$ in the considerations above.

REMARK . It is explicitly shown in [Wal78] that $H^{(p)}$ need not to be symmetric even if H_0 is. Thus in order to obtain a symmetric bi-solution (which is necessary for Hadamard bi-distributions arising from states) the initial functions H_0 are constraint (See [CH84] for the constraint in the scalar case). On the other hand the ‘DeWitt-Schwinger’ approach, which was used by Christensen, is effectively a choice of H_0 which yield to a symmetric $H^{(p)}$ [Wal], but it is not suitable for massless theories. Moreover almost all results in this approach have to be understood as being valid asymptotically only.

For the spinor case the analogous treatment for the auxiliary bi-distribution $\mathcal{G}^B_{A'}$ yields:

$$v^{(p)B}_{A'} = \sum_{i=1}^p v_i^B{}_{A'}(x, x') \sigma^i$$

and

$$H^{(p)B}_{A'} = \sum_{i=1}^p H_i^B{}_{A'}(x, x') \sigma^i.$$

For the corresponding coefficient functions the relations (114) and (115) are formally equally valid –except that now the covariant derivatives are understood as acting on Spinors [NO84]. For the boundary condition one finds

$$0 = v_{0;a} + \left(1 - \Delta^{-1/2} \Delta^{1/2}{}_{;b} \sigma^{;b}\right) v_0 + \frac{1}{2} \left(\square - \frac{1}{4} R + m^2\right) (\Delta^{1/2} \mathcal{J}),$$

where $\mathcal{J}$ is the bi-spinor of parallel transport defined by the relations $\mathcal{J}(x, x) = \mathbf{1}$ and $\sigma^{;a} \mathcal{J}_{;a}(x, x') = 0$ (Compare section 3). The function $v^{(p)}$ is again fixed by the geometry, whereas H_0 remains arbitrary. It is worth noting that the Remark written above applies verbatim to the spinor case, too. We close this chapter with some coincidence limits for the smooth DeWitt-Schwinger term ${}^{DS}H^{(p)}$.

1. The coincidence limit of the smooth DeWitt-Schwinger term

In this section some coincidence limits of the smooth DeWitt-Schwinger term ${}^{DS}H^{(p)}$ and its derivatives are collected. The results were used in Eqn. (108). The various coincidence limits of the DeWitt-Schwinger coefficients a_n and $\mathcal{A}_n$ respectively were taken from [Chr76, Chr78]. Consider ${}^{DS}H_0$ given by Eqn. (104) up to order $1/m^2$ and abbreviate $\delta = (\ln(m^2) - \ln 2 + 2\gamma)$, where γ is Euler's constant. Then for the scalar case

$${}^{DS}H_0 = (\Delta^{1/2})^{-1} m^2 (\delta - 1) - (\Delta^{1/2})^{-1} \delta a_1,$$

where a_1 is the second DeWitt-Schwinger coefficient defined in Eqn. (103) (see also [CH84]). We get the following coincidence limits:

$$\begin{aligned} [{}^{DS}H_0] &= m^2 (\delta - 1) \\ [{}^{DS}H_0^{;a}]^{;b} &= -\delta [a_1^{;a}]^{;b} = 0 \\ [{}^{DS}H_0^{;ab}] &= -\delta [a_1^{;ab}] = -\frac{\delta}{90} R^{cdea} R_{cde}{}^b \\ [{}^{DS}H_0]^{;ab} &= 0, \end{aligned}$$

since $[\delta^{1/2}] = 1$, $[\delta^{1/2;a}] = [\delta^{1/2}]^{;ab} = 0$ on vacuum spacetimes ($R^{ab} = 0$) and the coincidence limits of a_1 were used. The coincidence limit of ${}^{DS}H_1$ can be calculated using the Hadamard recursion relation. We obtain:

$$\begin{aligned} [{}^{DS}H_1] &= -\frac{1}{4} [{}^{DS}H_{0;h}{}^h] - \frac{1}{4} m^2 [{}^{DS}H_0] - \frac{3}{4} [a_2] \\ &= \frac{2\delta - 3}{720} R^{abcd} R_{abcd} - \frac{1}{4} m^4 (\delta - 1) \end{aligned}$$

For the fermionic case one finds the following coincidence limits

$$\begin{aligned} [{}^{DS}H_0] &= m^2 (\delta - 1) \\ [{}^{DS}H_0^{;a}]^{;b} &= -\delta [\mathcal{A}_1^{;a}]^{;b} = 0 \\ [{}^{DS}H_0^{;ab}] &= -\delta [\mathcal{A}_1^{;ab}] = -\frac{\delta}{48} G_{[cd]} G_{[ef]} (R^{cdga} R^{ef}{}^b{}_g + R^{cdgb} R^{ef}{}^a{}_g) - \frac{\delta}{90} R^{cdea} R_{cde}{}^b \\ [{}^{DS}H_{0;h}{}^h] &= \delta \left(-\frac{1}{24} G_{[cd]} G_{[ef]} R^{cdgh} R^{ef}{}_{gh} - \frac{1}{90} R^{abcd} R_{abcd} \right), \end{aligned}$$

where now $\mathcal{A}_n$ denotes the fermionic DeWitt-Schwinger coefficient and the Bianchi identities were used to show that terms like $R^{cdea}{}_{;e}{}^b$ vanish. For the coincidence limit of ${}^{DS}H_1$ we use the Hadamard recursion relation again:

$$\begin{aligned} [{}^{DS}H_1] &= -\frac{1}{4} [{}^{DS}H_{0;h}{}^h] - \frac{1}{4} m^2 [{}^{DS}H_0] - \frac{3}{4} [\mathcal{A}_2] \\ &= \frac{2\delta - 3}{192} G_{[cd]} G_{[ef]} R^{cdgh} R^{ef}{}_{gh} + \frac{2\delta - 3}{720} R^{abcd} R_{abcd} - \frac{m^4}{4} (\delta - 1) \end{aligned}$$

APPENDIX C

Useful formulas

This chapter contains some well known formulas that were frequently used during the calculations in this work. At the beginning some identities for the Dirac γ -matrices are collected. Some of the Fierz rearrangement formulas valid for classical Majorana spinors are presented next. We close this chapter with a proof of a rearrangement formula for the trace of the commutator of two Majorana spinor fields which is also valid in a quantized theory.

1. γ -matrix identities

In this section some useful and well known properties of the Dirac γ -matrices are collected (see e.g. [IZ80]). We begin with the basic relation and the definition of the γ^5 matrix:

$$\{\gamma^a, \gamma^b\}_+ = 2g^{ab}$$

$$\gamma^5 := \gamma^a \gamma^b \gamma^c \gamma^d \epsilon_{abcd},$$

where g^{ab} is the metric, ϵ_{abcd} denotes the totally antisymmetric ϵ tensor. An immediate consequence of this definition is the fact that γ^5 anti-commutes with the other γ -matrices and has unit square. For every product of the γ -matrices the trace of the resulting matrix is well defined and one finds:

$$\begin{aligned} \text{Tr}(\gamma^a \gamma^b) &= 4g^{ab} \\ \text{Tr}(\gamma^a \gamma^b \gamma^5) &= 0 \\ \text{Tr}(\gamma^a \gamma^b \gamma^c \gamma^d) &= 4(g^{ab} g^{cd} - g^{ac} g^{bd} + g^{ad} g^{bc}) \\ \text{Tr}(\gamma^a \gamma^b \gamma^c \gamma^d \gamma^5) &= -4i\epsilon^{abcd} \end{aligned}$$

Moreover, all traces over an odd number of γ -matrices vanish. The symbol $G^{[ab]}$ usually denotes 1/4 times the commutator of γ^a and γ^b . This implies

$$\begin{aligned} \text{Tr}(\gamma^a G^{[bc]} \gamma^d) &= 2(g^{ab} g^{cd} - g^{ac} g^{bd}) \\ \text{Tr}(\gamma^a G^{[bc]} \gamma^d \gamma^5) &= -2i\epsilon^{abcd} \\ \text{Tr}(G^{[kd]} G^{[ef]}) &= (g^{de} g^{kf} - g^{df} g^{ke}) \\ \text{Tr}(\gamma^a \gamma^c G^{[kd]} G^{[ef]}) &= g^{ac} g^{de} g^{kf} - g^{ac} g^{df} g^{ke} + g^{ka} g^{de} g^{cf} - g^{ka} g^{df} g^{ce} \\ &\quad - g^{ad} g^{ke} g^{cf} + g^{ad} g^{kf} g^{ce} + g^{fa} g^{ck} g^{ed} - g^{fa} g^{cd} g^{ek} \\ &\quad - g^{ea} g^{ck} g^{fd} + g^{ea} g^{cd} g^{fk} \end{aligned}$$

It is worth noting that on a curved spacetime the γ -matrices are covariantly constant, which implies the following identity:

$$\mathcal{J}^A_{A'} \gamma^{a'A'}_{B'} \mathcal{J}^{B'}_B = g^{a'}_{a'} \gamma^{aA}_{B}, \quad (117)$$

where $\mathcal{J}^A_{A'}$ and $g^{a'}_a$ are the bi-spinor and bi-tensor of parallel transport respectively (Compare chapter 3 and recall that parallel transport commutes with contraction).

2. Rearrangement of Majorana spinors

The following Fierz rearrangement formulas for classical Majorana spinors are valid. The proof of these well known identities can be found in the literature (see for example [MKW87, 1.3.4 and 1.4.7] or [Wes90, Appendix A]). It is worth noting that these identities are equally valid for a metric with signature $(-, +, +, +)$.

LEMMA C.1 (Fierz rearrangement identities). *For two Majorana spinor fields Ψ and χ the following identities hold.*

$$\Psi^+ \chi = \chi^+ \Psi \quad \Psi^+ \gamma^5 \chi = \chi^+ \gamma^5 \Psi \quad (118a)$$

$$\Psi^+ \gamma^a \chi = -\chi^+ \gamma^a \Psi \quad \Psi^+ \gamma^a \gamma^5 \chi = -\chi^+ \gamma^a \gamma^5 \Psi \quad (118b)$$

$$\Psi^+ \gamma^a \gamma^b \chi = \chi^+ \gamma^b \gamma^a \Psi \quad \Psi^+ \gamma^a \gamma^b \gamma^5 \chi = \chi^+ \gamma^5 \gamma^b \gamma^a \Psi \quad (118c)$$

$$\Psi^+(i\overleftrightarrow{\nabla} - m)\chi = \nabla_a(-\chi^+ i\gamma^a \Psi) + \chi^+(i\overleftrightarrow{\nabla} - m)\Psi \quad (118d)$$

While the identities above do not hold for quantized fields, the following result for the commutators only uses the Majorana condition and is hence also valid for quantized fields.

LEMMA C.2. Let χ^A and ϕ^+_B denote a Majorana spinor and a Majorana cospinor respectively. Then the following identity holds:

$$\text{Tr}(\gamma^{aA}{}_B[\chi^B, \phi^+_D]_-) = -\text{Tr}(\gamma^{aA}{}_B[\phi^B, \chi^+_D]_-),$$

where $[\cdot, \cdot]_-$ denotes the commutator.

PROOF. Write down the Definition of the trace and the commutator:

$$\begin{aligned} \text{Tr}(\gamma^{aA}{}_B[\chi^B, \phi^+_D]_-) &= \chi^A(\gamma^{aT})_A{}^B \phi^+_B - \phi^+_B \gamma^{aB}{}_A \chi^A \\ &= -\chi^A(\gamma^{aT})_A{}^B C_{BC} \phi^C + \phi^C (C^T)_{CB} \gamma^{aB}{}_A \chi^A, \end{aligned}$$

where C denotes the charge conjugation matrix. Recall that for a Majorana spinor $\phi^+_B = -C_{BC} \phi^C$. Using the properties of C we obtain

$$\begin{aligned} &= +\chi^A C_{AB} \gamma^{aB}{}_C \phi^C + \phi^C (\gamma^{aT})_C{}^B C_{BA} \chi^A \\ &= \chi^+_B \gamma^{aB}{}_C \phi^C - \phi^C \gamma^{aB}{}_C \chi^+_B \\ &= -\text{Tr}(\gamma^{aA}{}_B[\phi^A, \chi^+_C]_-) \end{aligned}$$

□

Listing of the Maple V programs

This Appendix contains the commented listing and an example run of the Maple V programs that were used for the calculations in chapter 4 section 5.

1. The Schwarzschild spacetime

```
> # Initialization
> # Read tensor package
> readlib(tensor):
> # and linear Algebra
> with(linalg):
> # Define Variables
> x1:=t:
> x2:=r:
> x3:=theta:
> x4:=phi:
> # Schwarzschild metric
> g11:=1-2*M/r:g22:=-1/g11:g33:=-r^2:g44:=-r^2*sin(theta)^2:
> # Compute Riemann etc.
> tensor():
> # {R_a}{bcd} = Rlhhh
> for a from 1 to 4 do
>   for b from 1 to 4 do
>     for c from 1 to 4 do
>       for d from 1 to 4 do
>         Rlhhh.a.b.c.d := 0:
>         for e from 1 to 4 do for f from 1 to 4 do for g from 1 to 4 do
>           Rlhhh.a.b.c.d := Rlhhh.a.b.c.d +
>             h.b.e * h.c.f * h.d.g * R.a.e.f.g
>         od od od
>       od od od od:
> # Kretschman invariant W2 = R^{abcd} R_{abcd}
> W2:=0:
> for a from 1 to 4 do for b from 1 to 4 do for c from 1 to 4 do
>   for d from 1 to 4 do for e from 1 to 4 do
>     W2 := W2 + h.e.a * R.a.b.c.d * Rlhhh.e.b.c.d:
>   od od od od od:
> # SUSY finite Term
> # Tsusy = 2 * 4 Pi * T_{ab} =
> #
> #           + 1/288 {R_a}{cde} R_{bcde}
> #           + 1/288 {R_a}{cde} R_{bedc}
> #           - 1/96 {R_a}{cde} R_{bdce}
> for a from 1 to 4 do for b from 1 to 4 do
>   Tsusy.a.b := 0:
>   for c from 1 to 4 do for d from 1 to 4 do for e from 1 to 4 do
>     Tsusy.a.b := Tsusy.a.b
>       + 1/288 * Rlhhh.a.c.d.e * R.b.c.d.e
>       + 1/288 * Rlhhh.a.c.d.e * R.b.e.d.c
>       - 1/96 * Rlhhh.a.c.d.e * R.b.d.c.e :
>   od od od
> od od:
> # Tsusyyh = Tsusy^{ab}
> for a from 1 to 4 do for b from 1 to 4 do
>   Tsusyyh.a.b := 0:
>   for c from 1 to 4 do for d from 1 to 4 do
>     Tsusyyh.a.b := Tsusyyh.a.b + h.a.c * h.b.d * Tsusy.c.d
>   od od
> od od:
> # Covariant Derivative
> # Tsusyh = T^{ab}_{;a} := T^{ab}_{;a}
> #           + {\gamma_{ad}}^a T^{db}
> #           + {\gamma_{ad}}^b T^{ad}
> for b from 1 to 4 do
>   Tsusyh.b := 0:
>   for a from 1 to 4 do
```

```

> Tsusyh.b := Tsusyh.b + diff(Tsusyhh.a.b,x.a):
> for d from 1 to 4 do
>   Tsusyh.b := Tsusyh.b +
>     C.a.d.a * Tsusyhh.d.b + C.a.d.b * Tsusyhh.a.d:
>   od
> od
> od:
> # Smooth terms: Tds = (2*\delta - 3)/768 * W2 -
> #                      \delta/48
> #                      ( {R_a}^{cde} * {R_bcd} + 2 {R_a}^{cde} *
> #                        R_{bdec} )
> for a from 1 to 4 do for b from 1 to 4 do
>   Tds.a.b := 0:
>   Tds.a.b := (2 * delta - 3) / 768 * W2:
>   for c from 1 to 4 do for d from 1 to 4 do for e from 1 to 4 do
>     Tds.a.b := Tds.a.b -
>       delta / 48 *
>       ( Rlhhh.a.c.d.e * R.b.c.e.d
>         + 2 * Rlhhh.a.c.d.e * R.b.d.e.c ):
>   od od od
> od od:
> # Tdshh = Tds^{ab}
> for a from 1 to 4 do for b from 1 to 4 do
>   Tdshh.a.b := 0:
>   for c from 1 to 4 do for d from 1 to 4 do
>     Tdshh.a.b := Tdshh.a.b + h.a.c * h.b.d * Tds.c.d
>   od od
> od od:
> # Covariant Derivative
> # Tdsh = T^{ab}_{;a} := T^{ab}_{;a}
> #                      + {\gamma_{ad}}^a T^{db}
> #                      + {\gamma_{ad}}^b T^{ad}
> for b from 1 to 4 do
>   Tdsh.b := 0:
>   for a from 1 to 4 do
>     Tdsh.b := Tdsh.b + diff(Tdshh.a.b,x.a):
>     for d from 1 to 4 do
>       Tdsh.b := Tdsh.b +
>         C.a.d.a * Tdshh.d.b + C.a.d.b * Tdshh.a.d:
>     od
>   od
> od:
> # Print results in pretty form
> #
> # define Matrix
> Tsusyhh:= matrix(4,4): # SUSY
> Tsusyh:= vector(4):
> Tdshh:= matrix(4,4): # Smooth part
> Tdsh:= vector(4):
> # Set Tsusyh to Tsusyhh
> for a to 4 do for b to 4 do
>   Tsusyhh[a,b] := simplify(Tsusyhh.a.b):
> od od:
> # Set Tsusyh to Tsusyh
> for a to 4 do
>   Tsusyh[a] := simplify(Tsusyh.a):
> od:
> # Set Tdsh to Tdshh
> for a to 4 do for b to 4 do
>   Tdshh[a,b] := simplify(Tdshh.a.b):
> od od:
> # Set Tdsh to Tdsh
> for a to 4 do
>   Tdsh[a] := simplify(Tdsh.a):
> od:
> print('Stress energy Tensor T^{ab}:');
> print(Tsusyhh);
> print('{T^{ab}}_{;a}:');
> print(Tsusyh);
> print('Smooth T^{ab}:');
> print(Tdshh);
> print(Divergence);
> print(Tdsh);
> # Output in LaTeX to files
> latex(Tsusyhh,Tsusyhh);
> latex(Tsusyh,Tsusyh);
> latex(Tdshh,Tdshh);
> latex(Tdsh,Tdsh);

```

```
> quit;
Warning: new definition for      norm
Warning: new definition for      trace
```

Stress energy Tensor T^{ab} :

$$\begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$T^{ab}{}_{;a}$:

$$[0000]$$

Smooth T^{ab} :

$$\begin{aligned} & \left[\frac{1}{16} \frac{M^2 (10 r \delta - 3 r - 16 \delta M)}{r^5 (r - 2 M)^2}, -\frac{1}{16} \frac{(2 \delta - 3) M^2}{r^6}, \right. \\ & \quad \left. -\frac{1}{16} \frac{(2 \delta - 3) M^2}{r^7 (r - 2 M)}, \frac{1}{16} \frac{(2 \delta - 3) M^2}{r^7 (-r + r \cos(\theta)^2 + 2 M - 2 M \cos(\theta)^2)} \right] \\ & \left[-\frac{1}{16} \frac{(2 \delta - 3) M^2}{r^6}, -\frac{1}{16} \frac{(6 r \delta + 4 \delta M + 3 r - 6 M) M^2 (r - 2 M)}{r^8} \right. \\ & \quad \left. , \frac{1}{16} \frac{(r - 2 M) (2 \delta - 3) M^2}{r^9}, -\frac{1}{16} \frac{M^2 (2 r \delta - 4 \delta M - 3 r + 6 M)}{r^9 (-1 + \cos(\theta)^2)} \right] \\ & \left[-\frac{1}{16} \frac{(2 \delta - 3) M^2}{r^7 (r - 2 M)}, \frac{1}{16} \frac{(r - 2 M) (2 \delta - 3) M^2}{r^9}, \right. \\ & \quad \left. -\frac{1}{16} \frac{M^2 (-2 \delta + 3 + 8 \delta r^2)}{r^{10}}, -\frac{1}{16} \frac{(2 \delta - 3) M^2}{r^{10} (-1 + \cos(\theta)^2)} \right] \\ & \left[\frac{1}{16} \frac{(2 \delta - 3) M^2}{r^7 (-r + r \cos(\theta)^2 + 2 M - 2 M \cos(\theta)^2)}, \right. \\ & \quad \left. -\frac{1}{16} \frac{M^2 (2 r \delta - 4 \delta M - 3 r + 6 M)}{r^9 (-1 + \cos(\theta)^2)}, -\frac{1}{16} \frac{(2 \delta - 3) M^2}{r^{10} (-1 + \cos(\theta)^2)}, \right. \\ & \quad \left. \frac{1}{16} \frac{M^2 (2 \delta - 3 - 8 \delta r^2 + 8 \delta r^2 \cos(\theta)^2)}{r^{10} (1 - 2 \cos(\theta)^2 + \cos(\theta)^4)} \right] \end{aligned}$$

Divergence

$$\begin{aligned}
& \left[\frac{1}{16} M^2 (-20 \delta M \sin(\theta) + 8 \delta \sin(\theta) r - 2 \delta \cos(\theta) + 30 M \sin(\theta) \right. \\
& \quad - 12 \sin(\theta) r + 3 \cos(\theta)) / (\sin(\theta) r^7 (r - 2 M)) \frac{1}{16} M^2 \left(\right. \\
& \quad 180 M^2 r^2 \cos(\theta)^2 - 88 M^3 r \delta - 132 M^3 r \cos(\theta)^2 - 6 r^2 \\
& \quad + 3 r^2 \cos(\theta)^2 + 84 M r^3 + 88 M^3 r \delta \cos(\theta)^2 \\
& \quad + 72 M^2 \delta r^2 \cos(\theta)^2 - 12 r M \cos(\theta)^2 - 136 M r^3 \delta \cos(\theta)^2 \\
& \quad + 12 r^4 \cos(\theta)^2 + 8 M r \delta \cos(\theta)^2 - 12 r^4 - 84 M r^3 \cos(\theta)^2 \\
& \quad - 72 M^2 \delta r^2 - 24 M^2 + 16 M^2 \delta - 16 M r \delta + 40 r^4 \delta \cos(\theta)^2 \\
& \quad - 8 M^2 \delta \cos(\theta)^2 - 2 \delta r^2 \cos(\theta)^2 + 4 \delta r^2 - 40 r^4 \delta + 136 r^3 M \delta \\
& \quad + 3 \sin(\theta) r^3 \cos(\theta) + 24 r M + 12 M^2 \cos(\theta)^2 \\
& \quad - 2 \sin(\theta) r^3 \cos(\theta) \delta + 8 \sin(\theta) r^2 \cos(\theta) M \delta \\
& \quad - 8 \sin(\theta) r \cos(\theta) M^2 \delta - 12 \sin(\theta) r^2 \cos(\theta) M \\
& \quad + 12 \sin(\theta) r \cos(\theta) M^2 - 180 M^2 r^2 + 132 M^3 r \left. \right) / \left(r^{10} \right. \\
& \quad \left. (-r + r \cos(\theta)^2 + 2 M - 2 M \cos(\theta)^2) \right) - \frac{1}{16} M^2 \left(-8 \delta \sin(\theta) r \right. \\
& \quad + 8 \delta r \sin(\theta) \cos(\theta)^2 + 12 \sin(\theta) r - 12 r \sin(\theta) \cos(\theta)^2 \\
& \quad + 20 \delta M \sin(\theta) - 20 M \delta \sin(\theta) \cos(\theta)^2 - 30 M \sin(\theta) \\
& \quad + 30 M \sin(\theta) \cos(\theta)^2 - 2 \delta \cos(\theta)^3 + 3 \cos(\theta)^3 \left. \right) / \left(\sin(\theta) r^{10} \right. \\
& \quad \left. (-1 + \cos(\theta)^2) \right) \frac{1}{16} M^2 (-20 \delta M \sin(\theta) + 8 \delta \sin(\theta) r \\
& \quad - 2 \delta \cos(\theta) + 30 M \sin(\theta) - 12 \sin(\theta) r + 3 \cos(\theta)) / \left(\sin(\theta) r^{10} \right. \\
& \quad \left. (-1 + \cos(\theta)^2) \right) \left. \right]
\end{aligned}$$

2. The Kerr spacetime

```

> # Initialization
> # Read tensor package
> readlib(tensor):
> # and linear Algebra
> with(linalg):
> # Define Variables
> x1:=t:
> x2:=r:
> x3:=theta:
> x4:=phi:
> rho:=(r^2 + a^2 * cos(theta)^2):
> # Kerr metric
> g11:=1- 2*M*r/rho :
> g14:= - (2*M*r*a / rho ) * sin(theta)^2 : g41:=g14 :
> g22:= - rho / (r^2 + a^2 - 2*M*r) :
> g33:= - rho :
> g44:= - ( r^2 + a^2 ) * sin(theta)^2 - (2*M*r*a^2 /rho ) * sin(theta)^4 :
> # Compute Riemann etc.
> tensor():
> # simplify
> # Riemann
> print('Simplify Riemann ...');
> for a from 1 to 4 do
>   for b from 1 to 4 do
>     print(a,b);
>     for c from 1 to 4 do
>       for d from 1 to 4 do
>         R.a.b.c.d := simplify(R.a.b.c.d)
>       od
>     od
>   od
> od:

```

```

> print('... done');
> # Metric
> for a from 1 to 5 do
>   for b from 1 to 4 do
>     h.a.b := simplify(h.a.b)
>   od
> od:
> # Christoffel
> print('Simplify Christoffels ...');
> for a from 1 to 4 do
>   for b from 1 to 4 do
>     print(a,b);
>     for c from 1 to 4 do
>       C.a.b.c := simplify(C.a.b.c)
>     od
>   od
> od:
> print('... done');
> # Show result
> #display(Ricciscalar,Einstein);
> # {R_a}^{bcd} = Rlhhh
> print('Calculating {R_a}^{bcd} ...');
> for a from 1 to 4 do
>   for b from 1 to 4 do
>     print(a,b);
>     for c from 1 to 4 do
>       for d from 1 to 4 do
>         Rlhhh.a.b.c.d := 0:
>         for e from 1 to 4 do for f from 1 to 4 do for g from 1 to 4 do
>           Rlhhh.a.b.c.d := simplify( Rlhhh.a.b.c.d +
>                                     h.b.e * h.c.f * h.d.g * R.a.e.f.g)
>         od od od
>       od od od od:
>     print('...done');
>   # SUSY finite Term
>   # Tsusy = 2 * 4 Pi * T_{ab} =
>   #
>   #               + 1/288 {R_a}^{cde} R_{bcde}
>   #               + 1/288 {R_a}^{cde} R_{bedc}
>   #               - 1/96 {R_a}^{cde} R_{bdce}
>   print('Calculating SUSY finite terms ...');
>   for a from 1 to 4 do for b from 1 to 4 do
>     Tsusy.a.b := 0: print(a,b);
>     for c from 1 to 4 do for d from 1 to 4 do for e from 1 to 4 do
>       Tsusy.a.b := (Tsusy.a.b
>                   + 1/288 * Rlhhh.a.c.d.e * R.b.c.d.e
>                   + 1/288 * Rlhhh.a.c.d.e * R.b.e.d.c
>                   - 1/96 * Rlhhh.a.c.d.e * R.b.d.c.e ):
>     od od od
>   od od:
>   print('... done');
>   # Tsusyh = Tsusy^{ab}
>   print('Calculating Tsusy^{ab} ...');
>   for a from 1 to 4 do for b from 1 to 4 do
>     Tsusyh.a.b := 0:
>     for c from 1 to 4 do for d from 1 to 4 do
>       Tsusyh.a.b := (Tsusyh.a.b + h.a.c * h.b.d * Tsusy.c.d)
>     od: Tsusyh.a.b := simplify(Tsusyh.a.b): od
>   od od:
>   print('... done');
>   # Covariant Derivative
>   # Tsusyh = T^{ab}_{;a} := U^{ab}_{;a}
>   #               + {\gamma_{ad}}^a T^{db}
>   #               + {\gamma_{ad}}^b T^{ad}
>   print('Calculating T^{ab}_{;a} ...');
>   for b from 1 to 4 do
>     Tsusyh.b := 0:
>     for a from 1 to 4 do
>       print(b,a);
>       Tsusyh.b := (Tsusyh.b + diff(Tsusyh.a.b,x.a)):
>       for d from 1 to 4 do
>         Tsusyh.b := (Tsusyh.b +
>                     C.a.d.a * Tsusyh.d.b + C.a.d.b * Tsusyh.a.d):
>       od
>     od
>   od:
>   print('... done');
>   # Print results in pretty form
>   #

```

```

> # define Matrix
> print('Printing ...');
> Tsusyh:= matrix(4,4): # SUSY
> Tsusyh:= vector(4):
> # Set Tsusyh to Tsusyh
> print('Tsusyh ...');
> for a to 4 do for b to 4 do
>   print(a,b);
>   Tsusyh[a,b] := simplify(Tsusyh.a.b):
> od od:
> print('... done');
> # Set Tsusyh to Tsusyh
> print('Tsusyh ...');
> for a to 4 do
>   print(a);
>   Tsusyh[a] := simplify(Tsusyh.a):
> od:
> print('... done');
> print('Stress energy Tensor T^{ab}:');
> print(Tsusyh);
> print('{T^{ab}}_{;a}:');
> print(Tsusyh);
> # Output in LaTeX to files
> latex(Tsusyh,KerrTsusyh);
> latex(Tsusyh,KerrTsusyh);
> print('... done');
> quit;
Warning: new definition for    norm
Warning: new definition for    trace

```

Simplify Riemann ...

1,1

1,2

⋮

4,4

... done

Simplify Christoffels ...

1,1

⋮

4,4

... done

Calculating $R_a{}^{bcd}$...

1,1

⋮

4,4

... done

Calculating SUSY finite terms ...

1,1

⋮

4,4

... done

Calculating T_{susy}^{ab} ...

... done

Calculating $T^{ab}_{;a}$...

1, 1

$\vdots$

4, 4

... done

Printing ...

$T_{susyhh}...$

1, 1

$\vdots$

4, 4

... done

$T_{susyh}...$

1

$\vdots$

... done

Stress energy Tensor T^{ab} :

$$\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$T^{ab}_{;a}$:

[0000]

...done

Bibliography

- [ALN77] S. L. Adler, J. Lieberman, and Y. J. Ng. Regularization of the stress-energy tensor for vector and scalar particles propagating in a general background metric. *Ann. Phys.*, 106:279–321, 1977.
- [B⁺90] N. N. Bogolubov et al. *General Principles of Quantum Field Theory*. Kluwer Academic Publishers, 1990.
- [Bar54] V. Bargmann. On unitary ray representations of continuous groups. *Ann. of Math.*, 59:1, 1954.
- [Bar84] R. Bartnik. Existence of maximal surfaces in asymptotically flat spacetimes. *Comm. Math. Phys.*, 94:155–175, 1984.
- [BC64] R. L. Bishop and R. J. Crittenden. *Geometry of Manifolds*. Academic press, New York, 1964.
- [BCO90] R. Bartnik, P. T. Chruściel, and N. O’Murchadha. On maximal surfaces in asymptotically flat space-times. *Comm. Math. Phys.*, 130:95–109, 1990.
- [BD82] N. D. Birrell and P. C. W. Davis. *Quantum fields in Curved Space*. Cambridge Univ. Press, Cambridge, 1982.
- [BDLR92] D. Buchholz, S. Doplicher, R. Longo, and J. E. Roberts. A new look at Goldstone’s theorem. *Rev. Math. Phys., Special Issue*, pages 49–83, 1992.
- [BF82] P. Breitenlohner and D. Z. Freedman. Stability in gauged extended supergravity. *Ann. of Phys.*, 144:249–281, 1982.
- [BFK94] I. Brunetti, K. Fredenhagen, and M. Köhler. The wave front set spectrum condition and Wick ordered products of free fields on a manifold. In Preparation, 1994.
- [Bor62] H. J. Borchers. On the structure of the algebra of the field operators. *Nuovo Cimento*, 24:214, 1962.
- [BT87] I. M. Benn and R. W. Tucker. *An introduction to spinors and geometry with applications in physics*. Adam Hilger, Bristol and Philadelphia, 1987.
- [BW92] H. Baumgärtel and M. Wollenberg. *Causal Nets of Operator Algebras*. Akademie Verlag GmbH, Berlin, Germany, 1992.
- [C⁺91a] B. W. Char et al. *Maple V Language reference manual*. Springer Verlag, New York, Berlin, 1991.
- [C⁺91b] B. W. Char et al. *Maple V Library reference manual*. Springer Verlag, New York, Berlin, 1991.
- [Car66] E. Cartan. *Theory of Spinors*. Hermann, Paris, 1966.
- [CB77] Y. Choquet-Bruhat. Hyperbolic differential equations on a manifold. In C. De Witt and J. Wheeler, editors, *Battelle Rencontres*, New York, 1977. Benjamin.
- [CCJ70] C. G. Jr. Callan, S. Coleman, and R. Jackiw. A new improved energy-momentum tensor. *Ann. of Phys.*, 59:42–73, 1970.
- [CGNP86] M. A. Castagnino, E. Gunzig, P. Nardone, and J. P. Paz. Hadamard and minimal renormalization. *Phys. Rev. D*, 34(12):3698–3706, 1986.
- [CH84] M. A. Castagnino and D. D. Harari. Hadamard renormalization in curved space-time. *Ann. of Phys.*, 152:85–104, 1984.
- [CHN82] M. Castagnino, D. Harari, and C. Nuñez. Minimal hypotheses for particle definition in curved space-time. In S. Ferrara and G. F. R. Ellis, editors, *Proceedings of the Europhysics Study conference: Unification of fundamental interactions II*, pages 455–466, New York, 1982. Plenum.
- [Chr76] S. M. Christensen. Vacuum expectation value of the stress tensor in an arbitrary curved background: The covariant point-separation method. *Phys. Rev. D*, 14(10):2490–2501, 1976.
- [Chr78] S. M. Christensen. Regularization, renormalization, and covariant geodesic point separation. *Phys. Rev. D*, 17(4):946–963, 1978.
- [Col84] J. F. Colombeau. *New generalized functions and multiplication of distributions*, volume 84. North Holland Math. Studies, 1984.
- [CW94] P. T. Chruściel and R. M. Wald. Maximal hypersurfaces in stationary asymptotically flat spacetimes. *Comm. Math. Phys.*, 163:561–604, 1994.
- [DB60] B. S. DeWitt and R. W. Brehme. Radiation damping in a gravitational field. *Ann. Phys.*, 9:220–259, 1960.
- [DeW65] B. S. DeWitt. *The Dynamical Theory of Groups and Fields*. Gordon and Breach, New York, 1965.
- [DeW75] B. S. DeWitt. Quantum field theory in curved spacetime. *Phys. Rep.*, 19(6):295–357, 1975.
- [DH72] J. J. Duistermaat and L. Hörmander. Fourier integral operators II. *Acta Math.*, 128:183, 1972.
- [Die72] J. Dieudonné. *Treatise on Analysis*, volume III. Academic Press, New York, 1972.
- [Dim80] J. Dimock. Algebras of local observables on a manifold. *Comm. Math. Phys.*, 77:219–228, 1980.
- [Dim82] J. Dimock. Dirac quantum fields on a manifold. *Trans. Am. Math. Soc.*, 269:133–147, 1982.
- [Dim92] J. Dimock. Quantized electromagnetic field on a manifold. *Rev. Math. Phys.*, 4:223–233, 1992.
- [FH87] K. Fredenhagen and R. Haag. Generally covariant quantum field theory and scaling limit. *Comm. Math. Phys.*, 108:91, 1987.
- [FH90] K. Fredenhagen and R. Haag. On the derivation of Hawking’s radiation associated with the formation of a black hole. *Comm. Math. Phys.*, 127:273–284, 1990.
- [Fit81] V. L. Fitch. Update on CP violation. In *On The Unity of the Fundamental Interactions*, Erice, pages 677–693, 1981.
- [FNW81] S. A. Fulling, F. J. Narcowich, and R. M. Wald. Singularity structure of the two-point function in quantum field theory in curved spacetime, II. *Ann. Phys. (NY)*, 136:243–272, 1981.
- [FSW78] S. A. Fulling, M. Sweeny, and R. M. Wald. Singularity structure of the two-point function in quantum field theory in curved spacetime. *Comm. Math. Phys.*, 63:257–264, 1978.
- [Ful89] S. A. Fulling. *Aspects of Quantum Field Theory in Curved Space-Time*. Cambridge University Press, Cambridge, 1989.
- [FZ75] S. Ferrara and B. Zumino. Transformation properties of the supercurrent. *Nucl. Phys. B*, 87:207–220, 1975.
- [Ger68] R. Geroch. Spinor structure of space-times in general relativity. I. *J. Math. Phys.*, 9:1739–1744, 1968.
- [Ger70] R. Geroch. Spinor structure of space-times in general relativity. II. *J. Math. Phys.*, 11:343–348, 1970.
- [Haa92] R. Haag. *Local quantum physics: Fields, particles, algebras*. Springer, Berlin, Germany, 1992.
- [Had23] J. Hadamard. *Lectures on Cauchy’s problem in Linear Partial Differential Equations*. Yale University, New Haven, 1923.
- [Haw75] S. W. Hawking. Particle creation by black holes. *Comm. Math. Phys.*, 43:199–220, 1975.

- [HE73] S. W. Hawking and G. F. R. Ellis. *The large scale structure of space-time*. Cambridge University Press, Cambridge, 1973.
- [HH76] J. B. Hartle and S. W. Hawking. Path integral derivation of black hole radiance. *Phys. Rev. D*, 21:2188–2203, 1976.
- [HNS84] R. Haag, H. Narnhofer, and U. Stein. On quantum field theory in gravitational background. *Comm. Math. Phys.*, 94:219–238, 1984.
- [Hör71] L. Hörmander. Fourier integral operators I. *Acta Math.*, 127:79, 1971.
- [Hör83] L. Hörmander. *The Analysis of Linear Partial Differential Operators I*. Springer, Berlin, 1983.
- [Hor90] Sergei Sergeevich Horuzhy. *Introduction to Algebraic Quantum Field Theory*. Kluwer, Dordrecht, Netherlands, 1990.
- [IZ80] C. Itzykson and J.-B. Zuber. *Quantum field theory*. McGraw-Hill, New York, 1980.
- [Kay92] B. S. Kay. The principle of locality and quantum field theory on (non globally hyperbolic) curved spacetimes. To be published, 1992.
- [KN63] S. Kobayashi and K. Nomizu. *Foundations of Differential Geometry I*. John Wiley & Sons, 1963.
- [Köh91] M. Köhler. Multiplikation von Distributionen und Skalenlimes. Diplomarbeit, TU-Berlin, 1991.
- [Köh93] M. Köhler. On the leading singularities of the expectation value of the energy momentum tensor of a locally supersymmetric quantum field on a curved space time. Internal Report, 1993.
- [Köh94] M. Köhler. New examples for Wightman fields on a manifold. *DESY-Preprint 94-161; Submitted to Class. and Quant. Grav.*, 1994.
- [KW91] B. S. Kay and R. M. Wald. Theorems on the uniqueness and thermal properties of stationary, nonsingular, quasifree states on spacetimes with a bifurcate Killing horizon. *Phys. Rep.*, 207(2):49–136, 1991.
- [Ler63] J. Leray. *Hyperbolic Differential Equations*. Lecture notes. Institute for Advanced Study, Princeton, N.J., 1963.
- [Łop91] J. Łopuszański. *An Introduction to symmetry and supersymmetry in quantum field theory*. World Scientific, Singapore, 1991.
- [LR90] C. Lüders and J. E. Roberts. Local quasiequivalence and adiabatic vacuum states. *Comm. Math. Phys.*, 134:29–63, 1990.
- [Mil63] J. Milnor. Spin structures on a manifold. *L'Enseig. Math.*, 9:168–173, 1963.
- [MKW87] H. J. W. Müller-Kirsten and A. Wiedemann. *Supersymmetry*. World Scientific, Singapore, 1987.
- [MTW73] C. W. Misner, K. S. Thorne, and J. A. Wheeler. *Gravitation*. Freeman, San Francisco, USA, 1973.
- [Nak90] M. Nakahara. *Geometry, Topology and Physics*. Adam Hilger, Bristol and New York, 1990.
- [Nie81] P. van Nieuwenhuizen. Supergravity. *Phys. Rep.*, 68(4):189–398, 1981.
- [NO84] A.-H. Najmi and A. C. Ottewill. Quantum states and the Hadamard form. II. Energy minimization for spin-1/2 fields. *Phys. Rev. D*, 30(12):2573, 1984.
- [Ohl92] S. Ohlmeyer. Raumzeiten ohne Spinstruktur. Diplomarbeit, Universität Hamburg, II. Institut für Theoretische Physik, 1992.
- [PR84] R. Penrose and W. Rindler. *Spinors and Spacetime I*. Cambridge University Press, 1984.
- [Rad] M. Radzikowski. Private communication. 1994.
- [Rad92] M. J. Radzikowski. *The Hadamard condition and Kay's conjecture in (axiomatic) quantum field theory on curved space-time*. PhD thesis, Princeton University, October 1992.
- [Rom60] P. Roman. *Theory of elementary Particles*. North-Holland Publishing Company, Amsterdam, 1960.
- [RS75] M. Reed and B. Simon. *Methods of Modern Mathematical Physics II*. Academic Press Inc. London, 1975.
- [Ste51] N. Steenrod. *The Topology of Fibre Bundles*. Princeton University Press, 1951.
- [SW64] R. F. Streater and A. S. Wightman. *PCT, Spin & Statistics and all that*. W. A. Benjamin, Inc., New York, 1964.
- [Tay81] M. E. Taylor. *Pseudodifferential Operators*. Princeton University Press, Princeton, New Jersey, 1981.
- [Uhl62] A. Uhlmann. Über die Definition der Quantenfelder nach Wightman und Haag. *Wiss. Zeitschrift Karl Marx Univ.*, 11:213, 1962.
- [Unr76] W. G. Unruh. Notes on black hole evaporation. *Phys. Rev. D*, 14:870–892, 1976.
- [Ver93] R. Verch. Nuclearity, split property, and duality for the Klein-Gordon field in curved spacetime. *Lett. Math. Phys.*, 29:297–310, 1993.
- [Ver94] R. Verch. Local definiteness, primarity and quasiequivalence of quasifree Hadamard quantum states in curved spacetime. *Comm. Math. Phys.*, 160:507–536, 1994.
- [W⁺57] C. S. Wu et al. Experimental test of parity conservation in beta decay. *Phys. Rev.*, 105:1413–1415, 1957.
- [Wal] R. M. Wald. Quantum field theory in curved spacetime. To appear in the Proceedings of the 1992 Les Houches School on Gravitation and Quantization.
- [Wal77] R. M. Wald. The back reaction effect in particle creation in curved spacetime. *Comm. Math. Phys.*, 54:1, 1977.
- [Wal78] R. M. Wald. Trace anomaly of a conformally invariant quantum field in curved spacetime. *Phys. Rev. D*, 17(6):1477–1484, 1978.
- [Wal84] R. Wald. *General Relativity*. The University of Chicago Press, 1984.
- [Wes90] P. West. *Introduction to Supersymmetry and Supergravity*. World Scientific, Singapore, 1990.
- [Wig39] E. Wigner. On unitary representations of the inhomogeneous Lorentz group. *Ann. of Math.*, 40:149, 1939.
- [Wol92] M. Wollenberg. Scaling limits and type of local algebras over curved spacetime. In W. B. Arveson et al., editors, *Operator algebras and topology*. Putman Research notes in Mathematics 270, Harlow: Longman, 1992.
- [WW90] R. S. Ward and R. O. Wells. *Twistor Geometry and Field Theory*. Cambridge University Press, 1990.
- [WZ74] J. Wess and B. Zumino. A lagrangian model invariant under supergauge transformations. *Phys. Lett.*, 49b:52, 1974.